

09/ 964,161

Connecting via Winsock to STN

Welcome to STN International! Enter x:x

LOGINID:sssptal202txn

PASSWORD:

TERMINAL (ENTER 1, 2, 3, OR ?):2

* * * * * Welcome to STN International * * * * *

NEWS 1 Web Page URLs for STN Seminar Schedule - N. America
NEWS 2 "Ask CAS" for self-help around the clock
NEWS 3 JAN 27 Source of Registration (SR) information in REGISTRY updated
and searchable
NEWS 4 JAN 27 A new search aid, the Company Name Thesaurus, available in
CA/CAplus
NEWS 5 FEB 05 German (DE) application and patent publication number format
changes
NEWS 6 MAR 03 MEDLINE and LMEDLINE reloaded
NEWS 7 MAR 03 MEDLINE file segment of TOXCENTER reloaded
NEWS 8 MAR 03 FRANCEPAT now available on STN
NEWS 9 MAR 29 Pharmaceutical Substances (PS) now available on STN
NEWS 10 MAR 29 WPIFV now available on STN
NEWS 11 MAR 29 No connect hour charges in WPIFV until May 1, 2004
NEWS 12 MAR 29 New monthly current-awareness alert (SDI) frequency in RAPRA

NEWS EXPRESS MARCH 31 CURRENT WINDOWS VERSION IS V7.00A, CURRENT
MACINTOSH VERSION IS V6.0c(ENG) AND V6.0Jc(JP),
AND CURRENT DISCOVER FILE IS DATED 13 APRIL 2004
NEWS HOURS STN Operating Hours Plus Help Desk Availability
NEWS INTER General Internet Information
NEWS LOGIN Welcome Banner and News Items
NEWS PHONE Direct Dial and Telecommunication Network Access to STN
NEWS WWW CAS World Wide Web Site (general information)

Enter NEWS followed by the item number or name to see news on that
specific topic.

All use of STN is subject to the provisions of the STN Customer
agreement. Please note that this agreement limits use to scientific
research. Use for software development or design or implementation
of commercial gateways or other similar uses is prohibited and may
result in loss of user privileges and other penalties.

* * * * * STN Columbus * * * * *

FILE 'HOME' ENTERED AT 13:43:56 ON 20 APR 2004

=> file reg

COST IN U.S. DOLLARS

SINCE FILE

TOTAL

ENTRY

SESSION

FULL ESTIMATED COST

0.21

0.21

FILE 'REGISTRY' ENTERED AT 13:44:06 ON 20 APR 2004

USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.

PLEASE SEE "HELP USAGETERMS" FOR DETAILS.

COPYRIGHT (C) 2004 American Chemical Society (ACS)

09/ 964,161

Property values tagged with IC are from the ZIC/VINITI data file provided by InfoChem.

STRUCTURE FILE UPDATES: 19 APR 2004 HIGHEST RN 676225-08-4
DICTIONARY FILE UPDATES: 19 APR 2004 HIGHEST RN 676225-08-4

TSCA INFORMATION NOW CURRENT THROUGH JANUARY 6, 2004



Please note that search-term pricing does apply when conducting SmartSELECT searches.

Crossover limits have been increased. See HELP CROSSOVER for details.

Experimental and calculated property data are now available. For more information enter HELP PROP at an arrow prompt in the file or refer to the file summary sheet on the web at:
<http://www.cas.org/ONLINE/DBSS/registryss.html>

=>

Uploading C:\STNEXP4\QUERIES\09964161.str



chain nodes :
7 8 9 10 11 12 15 16 17
ring nodes :
1 2 3 4 5 6
chain bonds :
4-7 7-8 8-9 8-15 9-10 10-11 10-12 15-16 15-17
ring bonds :
1-2 1-6 2-3 3-4 4-5 5-6
exact/norm bonds :
8-9 9-10 10-11 10-12 15-16 15-17
exact bonds :
4-7 7-8 8-15
normalized bonds :
1-2 1-6 2-3 3-4 4-5 5-6
isolated ring systems :
containing 1 :

G1:O,N

Match level :
1:Atom 2:Atom 3:Atom 4:Atom 5:Atom 6:Atom 7:CLASS 8:CLASS 9:CLASS 10:CLASS
11:Atom 12:CLASS 15:CLASS 16:CLASS 17:CLASS
Generic attributes :
11:
Number of Carbon Atoms : 7 or more
Type of Ring System : Polycyclic

Element Count :
Node 11: Limited
C,C9-13
O,O0-3
S,S0-3
N,N0-5

09/ 964,161

L1 STRUCTURE UPLOADED

=> s l1 ful

FULL SEARCH INITIATED 13:44:24 FILE 'REGISTRY'

FULL SCREEN SEARCH COMPLETED - 643968 TO ITERATE

47.2% PROCESSED 304013 ITERATIONS

943 ANSWERS

62.1% PROCESSED 400000 ITERATIONS

1110 ANSWERS

INCOMPLETE SEARCH (SYSTEM LIMIT EXCEEDED)

SEARCH TIME: 00.00.33

FULL FILE PROJECTIONS: ONLINE **INCOMPLETE**
BATCH **INCOMPLETE**

PROJECTED ITERATIONS: 643968 TO 643968


PROJECTED ANSWERS: 1661 TO 1913

L2 1110 SEA SSS FUL L1

=> d l1

L1 HAS NO ANSWERS

L1 STR



G1 O,N

Structure attributes must be viewed using STN Express query preparation.

=> s l2

SAMPLE SEARCH INITIATED 13:46:09 FILE 'REGISTRY'

SAMPLE SCREEN SEARCH COMPLETED - 32383 TO ITERATE

3.1% PROCESSED 1000 ITERATIONS

3 ANSWERS

INCOMPLETE SEARCH (SYSTEM LIMIT EXCEEDED)

SEARCH TIME: 00.00.01

FULL FILE PROJECTIONS: ONLINE **INCOMPLETE**
BATCH **INCOMPLETE**

PROJECTED ITERATIONS: 636917 TO 658403

PROJECTED ANSWERS: 1351 TO 2533

L3 3 SEA SSS SAM L1

=> file caplus

COST IN U.S. DOLLARS

SINCE FILE
ENTRY

TOTAL
SESSION

09/ 964,161

FULL ESTIMATED COST

156.68

156.89

FILE 'CAPLUS' ENTERED AT 13:46:39 ON 20 APR 2004

USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.

PLEASE SEE "HELP USAGETERMS" FOR DETAILS.

COPYRIGHT (C) 2004 AMERICAN CHEMICAL SOCIETY (ACS)

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications. The CA Lexicon is the copyrighted intellectual property of the American Chemical Society and is provided to assist you in searching databases on STN. Any dissemination, distribution, copying, or storing of this information, without the prior written consent of CAS, is strictly prohibited.

FILE COVERS 1907 - 20 Apr 2004 VOL 140 ISS 17

FILE LAST UPDATED: 19 Apr 2004 (20040419/ED)

This file contains CAS Registry Numbers for easy and accurate substance identification.

=> s l2

L4 261 L2


=> d l4 1- ibib abs fhitr

YOU HAVE REQUESTED DATA FROM 261 ANSWERS - CONTINUE? Y/(N):y

L4 ANSWER 1 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
 ACCESSION NUMBER: 2004-252486 CAPLUS
 TITLE: Preparation of quinoline and naphthyridine derivatives
 as HIV integrase inhibitors
 INVENTOR(S): Murai, Hitoshi; Endo, Takeshi; Kurose, Noriyuki;
 Taishi, Teruhiko; Yoshida, Hiroshi
 PATENT ASSIGNEE(S): Shionogi & Co., Ltd., Japan
 SOURCE: PCT Int. Appl., 396 pp.
 CODEN: PIXXD2
 DOCUMENT TYPE: Patent
 LANGUAGE: Japanese
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2004024693	A1	20040325	WO 2003-JP10212	20030811
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
PRIORITY APPLN. INFO.:				
		JP 2002-235582	A	20020813
		JP 2002-245772	A	20020826
		JP 2003-121726	A	20030425
		JP 2003-270863	A	20030704

GI



AB The title compds. I [wherein B1 = N or (un)substituted CH; R1 = H, (un)substituted alkyl, alkenyl, etc.; R1' = H, halo, NO2, OH, CO2H, (un)substituted alkoxy, alkyl, alkoxy, etc.; A1 = (un)substituted -CH=CH-CH=CH-, -CH=CH-CH=CH-, -CH=CH-CH=CH-, -CH=CH-CH=CH-, -CH=CH-CH=CH-, -CH=CH-CH=CH-, or -CH=CH-CH=CH-] or prodrugs, solvates, or pharmaceutically acceptable salts thereof are prepared as HIV integrase inhibitors. For example, the compound II was prepared in a multi-step synthesis. II showed inhibitory activity with IC50 of 0.071 µg/mL against integrase. Formulations containing I as an active ingredient were also described.

IT 675611-88-8P

L4 ANSWER 2 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
 ACCESSION NUMBER: 2004-203814 CAPLUS
 DOCUMENT NUMBER: 140-253449
 TITLE: Preparation of heterocyclylcarboxamides as oxytocin inhibitors
 INVENTOR(S): Armour, Duncan Robert; Bell, Andrew Simon; Edwards, Paul John; Ellis, David; Hepworth, David; Lewis, Mark Llewellyn; Smith, Christopher Ronald
 PATENT ASSIGNEE(S): Pfizer Limited, UK; Pfizer Inc.
 SOURCE: PCT Int. Appl., 124 pp.
 CODEN: PIXXD2
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2004020414	A1	20040311	WO 2003-IB3705	20030813
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
PRIORITY APPLN. INFO.:				
		GB 2002-19961	A	20020828


AB R1CON[(CH2)xR2]C(R4)[(CH2)yR3](CH2)zR5 [R1 = (substituted) Ph, heteroaryl; R2 = (substituted) Ph, OPh, cycloalkyl, heteroaryl, heterocyclyl, etc.; R3 = (substituted) (fused) Ph, heterocyclyl, heteroaryl, R6, etc.; R4 = H, Me; R5 = CONH2, NH2, OH, R6, NHR6, OR6, CONHR6, (substituted) heteroaryl, etc.; R6 = alkyl; x, y, z = 0-2]. were prepared Thus, 4-chlorobenzylamine, o-tolualdehyde, 2-aminocyclohexanecarboxylic acid, and (4-isocyanocyclohex-3-enyl)benzene (preparation given) were stirred in MeOH/cyclohexane to give a residue which was stirred in aqueous HCl/THF to give 2-amino-N-[2-amino-1-(2-methylphenyl)-2-oxoethyl]-N-(4-chlorobenzyl)nicotinamide. Title compds. at 10 µM gave >70% inhibition of oxytocin.

IT 669087-02-9P

RL: PAC (Pharmacological activity); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)
 (preparation of heterocyclylcarboxamides as oxytocin inhibitors)
 RN 669087-02-9 CAPLUS
 CN 2-Quinolincarboxamide, N-[2-amino-1-(3-methoxyphenyl)-2-oxoethyl]-N-[(4-methylphenyl)methyl]- (9CI) (CA INDEX NAME)


L4 ANSWER 1 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)
 RL: PAC (Pharmacological activity); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)
 (drug candidate; prepn. of quinoline and naphthyridine derivs. as HIV integrase inhibitors)
 RN 675611-88-8 CAPLUS
 CN INDEX NAME NOT YET ASSIGNED

Absolute stereochemistry.



REFERENCE COUNT: 12 THERE ARE 12 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 2 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)



REFERENCE COUNT: 15 THERE ARE 15 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 3 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
 ACCESSION NUMBER: 2004:143094 CAPLUS
 DOCUMENT NUMBER: 140:199743
 TITLE: Preparation of substituted (2S)-(arylamino)-3-(biphenyl-4-yl)propionic acids as antagonists of factor IX for inhibiting the intrinsic pathway of blood coagulation
 INVENTOR(S): Mjalli, Adnan M. M.; Andrews, Robert C.; Guo, Xiao-chuan; Christen, Daniel Peter; Gohimmukula, Devi Reddy; Huang, Guoxiang; Rothlein, Robert; Tyagi, Sameer; Yamasu, Tripura; Behme, Christopher
 PATENT ASSIGNEE(S): Transtech Pharma, Inc., USA
 SOURCE: PCT Int. Appl., 326 pp.
 CODEN: PIXXD2
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2004018444	A2	20040219	WO 2003-US25045	20030808
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU			
RW:	GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG			

PRIORITY APPL. INFO.: US 2002-402272P P 20020809
 OTHER SOURCE(S): MARPAT 140:199743
 AB The title compds. Ar2XCH(Var1)(CH2)CG [I; c = 0-2; G = H, CO2R1, CH2OR1, COR1, CRI; NOR2, an acid isostere (wherein R1, R2 = H, alkyl, aryl, etc.); V = (CH2)bo(CH2)a, (CH2)bNR7(CH2)a, (CH2)bo, (CH2)bNR7, (CH2)a, a bond (a = 0-2; b = 1-2; R7 = H, alkyl, aryl, etc.); X = NR8, COR8, NR8CO, etc. (R8 = H, alkyl, aryl, etc.); Ar1 = (un)substituted aryl, heteroaryl, cycloalkylaryl, etc.; Ar2 = (un)substituted aryl or heteroaryl], useful as antagonists, or more preferably, partial antagonists of factor IX and thus, may be used to inhibit the intrinsic pathway of blood coagulation. were prepared Thus, reacting Me 2-L-amino-3-biphenyl-4-yl-propionate with isoguinoline-3-carboxylic acid followed by hydrolysis afforded 3-L-3-biphenyl-4-yl-(2S)-[[(isoguinoline-3-carbonyl)amino]propionic acid. The compds. I inhibit factor IX with IC50 of less than 30 µM, and are useful in a variety of applications including the management, treatment and/or control of diseases caused in part by the intrinsic clotting pathway utilizing factor IX. Such diseases or disease states include stroke, myocardial infarction, aneurysm surgery, and deep vein thrombosis associated with surgical procedures, long periods of confinement, and acquired or inherited pro-coagulant states. The pharmaceutical composition comprising the compound I is claimed.

IT 660823-98-3P
 RL: PAC (Pharmacological activity); RCT (Reactant); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP

L4 ANSWER 3 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)
 (Preparation); RACT (Reactant or reagent); USES (Uses)
 (prepn. of substituted (2S)-(arylamino)-3-(biphenyl-4-yl)propionic acids as antagonists of factor IX for inhibiting the intrinsic pathway of blood coagulation)
 RN 660823-98-3 CAPLUS
 CN [1,1'-Biphenyl]-4-propanoic acid, α-[[[(7-bromo-3-isoguinolinyl)carbonyl]amino]-, (αS)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




L4 ANSWER 4 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
 ACCESSION NUMBER: 2004:120719 CAPLUS
 DOCUMENT NUMBER: 140:175176
 TITLE: Use of a proteasome inhibitor in the treatment of endothelial dysfunction and/or in a low-dose proteasome inhibitor therapy
 INVENTOR(S): Stangl, Verena; Stangl, Karl; Lorenz, Mario
 PATENT ASSIGNEE(S): Charite-Universitätsmedizin Berlin, Germany
 SOURCE: PCT Int. Appl., 37 pp.
 CODEN: PIXXD2
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2004012732	A2	20040212	WO 2003-EP8495	20030731
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW, AM, AZ, BY, BG, KZ, MD, RU, TJ, TM			
RW:	GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GM, ML, MR, NE, SN, TD, TG			

PRIORITY APPL. INFO.: EP 2002-17234 A 20020731
 AB The invention discloses the use of a proteasome inhibitor for the manufacture of a medicament for the prevention, onset therapy, acute therapy, and/or regression of diseases associated with endothelial dysfunction. The invention also discloses the use of a proteasome inhibitor as a low-dose treatment.

IT 179324-99-5
 RL: PAC (Pharmacological activity); THU (Therapeutic use); BIOL (Biological study); USES (Uses)
 (proteasome inhibitor in treatment of endothelial dysfunction and/or in low-dose proteasome inhibitor therapy)
 RN 179324-99-5 CAPLUS
 CN Boronic acid, [(1R)-3-methyl-1-[[[(2S)-1-oxo-4-phenyl-2-[[2-(quinolinyl)carbonyl]amino]butyl]amino]butyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




L4 ANSWER 5 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
 ACCESSION NUMBER: 2004:102765 CAPLUS
 DOCUMENT NUMBER: 140:270821
 TITLE: Halogenoalkyl Isocyanates as Bifunctional Reagents in an Aza-Wittig/Heterocyclization Reaction on the Solid Phase: Efficient Entry into New Tetracyclic Benzimidazole Systems
 AUTHOR(S): Hoel, Cornelia E.; Nefzi, Adel; Houghten, Richard A.
 CORPORATE SOURCE: Torrey Pines Institute for Molecular Studies, San Diego, CA, 92121, USA
 SOURCE: Journal of Combinatorial Chemistry (2004), 6(2), 220-223
 CODEN: JCCHFF; ISSN: 1520-4766
 PUBLISHER: American Chemical Society
 DOCUMENT TYPE: Journal
 LANGUAGE: English

AB An efficient one-pot procedure for the solid-phase synthesis of new tetracyclic 1,3,5-triazino[1,2-a]benzimidazolium derivs. starting from resin-bound benzimidazoles is described. The synthetic strategy involves an unprecedented one-pot aza-Wittig/heterocyclization/substitution reaction sequence using halogenoalkyl isocyanates. The structure of the tetracyclic ring system was determined by two-dimensional NMR expts. and X-ray anal.

IT 673461-98-8P
 RL: SPN (Synthetic preparation); PREP (Preparation)
 (solid-phase synthesis of 1,3,5-triazino[1,2-a]benzimidazolium derivs. via one-pot aza-Wittig/heterocyclization/substitution reaction sequence using halogenoalkyl isocyanates)
 RN 673461-98-8 CAPLUS
 CN 1H-5H-Imidazo[2',1':4,5][1,3,5]triazino[1,2-a]benzimidazolium, 8-[[[(2-amino-2-oxo-1-phenylethyl)amino]carbonyl]-11-butyl-1-[2-(butylamino)ethyl]-2,3-dihydro-5-oxo-, salt with trifluoroacetic acid (1:1) (9CI) (CA INDEX NAME)

CM 1

CRN 673461-97-7
 CMF C30 H39 N8 O3



CM 2

CRN 14477-72-6
 CMF C2 F3 O2

L4 ANSWER 5 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)




REFERENCE COUNT: 10 THERE ARE 10 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 6 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 2004:56093 CAPLUS
DOCUMENT NUMBER: 140:248526
TITLE: Ochratoxin A: Lack of Formation of Covalent DNA Adducts
AUTHOR(S): Mally, Angela; Zepnik, Herbert; Wanek, Paul; Eder, Erwin; Dingley, Karen; Ihmels, Heiko; Voelkel, Wolfgang; Dekant, Wolfgang
CORPORATE SOURCE: Institut fuer Toxikologie, Universitaet Wuerzburg, Wuerzburg, 97078, Germany
SOURCE: Chemical Research in Toxicology (2004), 17(2), 234-242
CODEN: CRTOC; ISSN: 0893-228X
PUBLISHER: American Chemical Society
DOCUMENT TYPE: Journal
LANGUAGE: English
AB The mycotoxin ochratoxin A (OTA) is a potent nephrotoxin and renal carcinogen in rodents. However, the mechanism of OTA-induced tumor formation is unknown and conflicting results have been obtained regarding the potential of OTA to bind to DNA. OTA is poorly metabolized, and no reactive intermediates capable of interacting with DNA have been detected in vitro or in vivo. Recently, a hydroquinone/quinone redox couple and a carbon-bonded OTA-deoxyguanosine (OTA-dG) adduct formed by electrochem. oxidation and photoreaction of OTA have been reported and suggested to be involved in OTA carcinogenicity. This study was designed to characterize the role of DNA binding and to determine if formation of these derivs. occurs in vivo and in relevant activation systems in vitro using specific and sensitive methods. Horseradish peroxidase activation of OTA and its dechlorinated analog ochratoxin B (OTB) yielded ochratoxin A-hydroquinone (OTHQ), but the postulated OTA-dG adduct was not detectable using LC-MS/MS. In support of this, no OTA-related DNA adducts were observed by 32P-postlabeling. In vivo, only traces of OTHQ were found in the urine of male F344 rats treated with high doses of OTA (2 mg/kg body wt) for 2 wk, suggesting that this metabolite is not formed to a relevant extent. In agreement with the in vitro data, OTA-dG was not detected by LC-MS/MS in liver and kidney DNA extracted from treated animals. In addition, DNA binding of OTB and OTB was assessed in male rats given a single dose of 14C-OTA or 14C-OTB using accelerator mass spectrometry, a highly sensitive method for quantifying extremely low concns. of radiocarbon. The 14C content in liver and kidney DNA from treated animals was not significantly different from controls, indicating that OTA does not form covalent DNA adducts in high yields. In summary, the results presented here demonstrate that DNA binding of OTA is not detectable with sensitive anal. and is unlikely to represent a mechanism for OTA-induced tumor formation.
IT 205034-32-8
RL: FMU (Formation, unclassified); FORM (Formation, nonpreparative)
(DNA binding of ochratoxin A is not detectable with sensitive anal. and is unlikely to represent a mechanism for OTA-induced tumor formation)
RN 205034-32-8 CAPLUS
CN L-Phenylalanine, N-[[[(3R)-3,4-dihydro-5,8-dihydroxy-3-methyl-1-oxo-1H-2-benzopyran-7-yl]carbonyl]- (9CI) (CA INDEX NAME)
Absolute stereochemistry.


L4 ANSWER 6 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)



REFERENCE COUNT: 30 THERE ARE 30 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 7 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN


ACCESSION NUMBER: 2004:45782 CAPLUS
DOCUMENT NUMBER: 140:253536
TITLE: Synthesis and Structure-Activity Relationships of Novel Arylpiperazines as Potent and Selective Agonists of the Melanocortin Subtype-4 Receptor
AUTHOR(S): Richardson, Timothy I.; Ornstein, Paul L.; Briner, Karin; Fisher, Matthew J.; Backer, Ryan T.; Biggers, C. Kelly; Clay, Michael P.; Emmerson, Paul J.; Hertel, Larry W.; Hsiung, Hansen M.; Husain, Saba; Kahl, Steven D.; Lee, Jonathan A.; Lindstrom, Terry D.; Martinelli, Michael J.; Mayer, John P.; Mullaney, Jeffery T.; O'Brien, Thomas P.; Pawlak, Joseph M.; Revell, Kevin D.; Shah, Jikesh; Zgombick, John M.; Herr, R. Jason; Melekhov, Alex; Sampson, Peter B.; King, Chi-Hsin R.
CORPORATE SOURCE: Lilly Research Laboratories, A Division of Eli Lilly and Company Lilly Corporate Center, Indianapolis, IN, 46285, USA
SOURCE: Journal of Medicinal Chemistry (2004), 47(3), 744-755
CODEN: JMCMAR; ISSN: 0022-2623
PUBLISHER: American Chemical Society
DOCUMENT TYPE: Journal
LANGUAGE: English
GI



AB The melanocortin receptors have been implicated as potential targets for a number of important therapeutic indications, including inflammation, sexual dysfunction, and obesity. (3R)-N-[[[(1R)-2-[4-[2-(Aminosulfonyl)phenyl]-1-piperazinyl]-1-[(4-chlorophenyl)methyl]-2-oxoethyl]-1,2,3,4-tetrahydro-3-isquinolinecarboxamide (I, R = H2NSO2) an arylpiperazine attached to dipeptide H-D-Tic-D-p-CI-Ph-OH, as a novel melanocortin subtype-4 receptor (MC4R) agonist through iterative directed screening of nonpeptidyl G-protein-coupled receptor biased libraries. Structure-activity relationship (SAR) studies demonstrated that substitutions at the ortho position of the aryl ring improved binding and functional potency. For example, the o-isopropyl-substituted compound I (R = isopropyl) (Ki = 720 nM) possessed 9-fold better binding affinity compared to the unsubstituted aryl ring (Ki = 6600 nM). Sulfonamide I (R = MeSO2NH) (II) (Ki = 220 nM) fills this space with a polar substituent, resulting in a further 2-fold improvement in binding affinity. The most potent compds. such as the dimethylamine derivative I (R = Me2NCH2) (Ki = 60 nM) contain a basic group at this position. Basic heterocycles such as the imidazole I (R = (1H-imidazol-1-yl)methyl) (Ki = 110 nM) were similarly effective. Good oral bioavailability for sulfonamide II was also demonstrated.
IT 254008-71-2P
RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT

L4 ANSWER 7 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)
 (Reactant or reagent)
 (prepn. and structure-activity relationship of
 ((arylpiperazinyl)[(chlorophenyl)methyl]oxoethyl]tetrahydroisoquinoline
 carboxamide deriva. as selective melanocortin subtype-4 receptor
 agonists)
 RN 252008-71-2 CAPLUS
 CN 2-(1H)-isoquinolinecarboxylic acid, 3-[[[(1R)-1-carboxy-2-(4-chlorophenyl)ethyl]amino]carbonyl]-3,4-dihydro-, 2-(1,1-dimethylethyl)
 ester, (3R)-(9CI) (CA INDEX NAME)

Absolute stereochemistry.



REFERENCE COUNT: 31 THERE ARE 31 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT


L4 ANSWER 8 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
 ACCESSION NUMBER: 2004:41297 CAPLUS
 DOCUMENT NUMBER: 140:105236
 TITLE: Proteasome inhibitors for the treatment of herpesviridae-infected individuals
 INVENTOR(S): Prosch, Susanne; Volk, Hans Dieter; Kruger, Detlev
 PATENT ASSIGNEE(S): Universitätsklinikum Charité der Humboldt-Universität zu Berlin Technologietrans Ferstelle, Germany
 SOURCE: PCT Int. Appl., 47 pp.
 CODEN: PIXXD2
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2004004749	A1	20040115	WO 2003-EP7062	20030702
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MY, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW, AM, AZ, BY, BG, KZ, MD, RU, TJ, TM				
RW: GH, GM, KE, LS, MW, MZ, SD, SI, SZ, TZ, UG, ZM, ZW, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GN, GQ, GW, ML, MR, NE, SN, TD, TG				

PRIORITY APPLN. INFO.: EP 2002-14728 A 20020703
 AB The invention discloses the use of a substance or composition comprising one or more proteasome inhibitors for the manufacture of a medicament for the treatment of an individual infected with a virus selected from the group comprising varicella zoster virus, human cytomegalovirus, human herpesvirus 6 and 7 and Epstein-Barr virus and Kaposi's sarcoma herpesvirus. The invention further discloses methods for treatment of individuals infected with a virus selected from the group comprising varicella zoster virus, human cytomegalovirus, human herpesvirus 6 and 7 and Epstein-Barr virus and Kaposi's sarcoma herpesvirus.
 IT 179324-59-5
 RL: PAC (Pharmacological activity); THU (Therapeutic use); BIOL (Biological study); USES (Uses)
 (proteasome inhibitors for treatment of herpesviridae infection)
 RN 179324-59-5 CAPLUS
 CN Boronic acid, ((1R)-3-methyl-1-[[[(2S)-1-oxo-4-phenyl-2-[[2-(quinolinylcarbonyl)amino]butyl]amino]butyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

L4 ANSWER 8 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)




REFERENCE COUNT: 5 THERE ARE 5 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 9 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
 ACCESSION NUMBER: 2004:41276 CAPLUS
 DOCUMENT NUMBER: 140:105251
 TITLE: 3,4-Dihydroisoquinolin-1-one derivatives as inducers of apoptosis
 INVENTOR(S): Gangloff, Anthony R.; Litvak, Joanne; Pararajasingham, Keith; Sperandio, David
 PATENT ASSIGNEE(S): Axys Pharmaceuticals, Inc., USA
 SOURCE: PCT Int. Appl., 107 pp.
 CODEN: PIXXD2
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2004004727	A1	20040115	WO 2003-US21102	20030703
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MY, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW, AM, AZ, BY, BG, KZ, MD, RU, TJ, TM				
RW: GH, GM, KE, LS, MW, MZ, SD, SI, SZ, TZ, UG, ZM, ZW, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GN, GQ, GW, ML, MR, NE, SN, TD, TG				

PRIORITY APPLN. INFO.: MARPAT 140:105251
 OTHER SOURCE(S):
 AB The invention discloses 3,4-dihydroisoquinolin-1-one derivs. that are activators of caspases and inducers of apoptosis, as well as pharmaceutical compns. comprising these compds., and methods for treating cancer using these compds. Preparation of selected compds. of the invention is included.
 IT 646028-31-1
 RL: PAC (Pharmacological activity); THU (Therapeutic use); BIOL (Biological study); USES (Uses)
 (dihydroisoquinolinone derivs. as inducers of apoptosis, and use for cancer treatment)
 RN 646028-31-1 CAPLUS
 CN L-Phenylalanine, N-[[[3-(3,5-bis(trifluoromethyl)phenyl)-1,2,3,4-tetrahydro-2-methyl-1-oxo-4-isoquinolinyl]carbonyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




L4 ANSWER 9 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)

REFERENCE COUNT: 7 THERE ARE 7 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 10 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 2004:2899 CAPLUS
 DOCUMENT NUMBER: 140:27711
 TITLE: Preparation of etoposide aglycon analogs as antitumor agents and DNA topoisomerase II inhibitors
 INVENTOR(S): Lee, Kuo-Hsiung; Xiao, Zhiyan; Bastow, Kenneth F.
 PATENT ASSIGNEE(S): University of North Carolina at Chapel Hill, USA
 SOURCE: PCT Int. Appl., 33 pp.
 CODEN: PIXXD2
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 2
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2004000859	A2	20031231	WO 2003-US19629	20030620
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ			
RW:	GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LJ, MC, NL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG			
US 6566393	B1	20030520	US 2002-177147	20020621
US 2004006126	A1	20040108	US 2003-349351	20030122
PRIORITY APPL. INFO.:			US 2002-177147 A	20020621
			US 2003-349351 A	20030122
OTHER SOURCE(S):		MARPAT 140:27711		
GI				



AB Etoposide amino acid analogs I (X = O, S, NH, CO, CH₂NH; R1 = covalent linkage between X and Y, alkyl, alkenyl, (un)substituted Ph; Y = NHCO, CONH; Z = CHR₂(CH₂)nR₃, R2 = CO₂H, NH₂, ester, etc., R3 = alkyl,

L4 ANSWER 10 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)

alkenyl, aryl, n= 0-2; D = CH₂OC(O), CH₂OC(=CH₂), CH₂CH₂C(O), CH₂OCH₂, CH₂OC(S), CH₂O(SO₂)CH₂, etc.) were prepd. as DNA topoisomerase II inhibitors. Thus, 4'-O-demethyl-4β-[(4''-(methyl-L-tyrosine-N-carbonyl)anilino]-4-deoxy-podophyllotoxin (II) and 4'-O-demethyl-4β-[(4''-(methyl-L-tryptophan-N-carbonyl)anilino]-4-deoxypodophyllotoxin were prepd. from podophyllotoxin and their pharmaceutical activity evaluated. The antitumor ED₅₀ of II against A 549 human cell line was 2.4 μM. Cancer is selected from the group consisting of skin cancer, lung cancer, Kaposi's sarcoma, testicular cancer, lymphoma, leukemia, esophageal cancer, stomach cancer, colon cancer, breast cancer, endometrial cancer, ovarian cancer, central nervous system cancer, liver cancer and prostate cancer.

IT 527678-17-7p


RL: PAC (Pharmacological activity); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)

(preparation of etoposide amino acid analogs as DNA topoisomerase II inhibitors)

RN 527678-17-7 CAPLUS

CN L-Tyrosine, N-[[[5S,5aR,8aR,9R)-5,5a,6,8,8a,9-hexahydro-9-(4-hydroxy-3,5-dimethoxyphenyl)-8-oxofuro[3',4':6,7]naphtho[2,3-d]-1,3-dioxol-5-yl]carbonyl]-, methyl ester (9CI) (CA INDEX NAME)

Absolute stereochemistry.



L4 ANSWER 11 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 2004:2730 CAPLUS
 DOCUMENT NUMBER: 140:71043
 TITLE: Combination treatment for depression and anxiety by NK1 and NK3 antagonists
 INVENTOR(S): Sobolov-Jaymes, Susan Beth; Lowe, John Adams, III; McLean, Stafford
 PATENT ASSIGNEE(S): Pfizer Products Inc., USA
 SOURCE: PCT Int. Appl., 124 pp.
 CODEN: PIXXD2
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2004000355	A1	20031231	WO 2003-1B2516	20030610
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM			
RW:	GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LJ, MC, NL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG			
US 2004006135	A1	20040108	US 2003-366582	20030312
PRIORITY APPL. INFO.:			US 2002-389975P P	20020619
OTHER SOURCE(S):		MARPAT 140:71043		


AB The invention discloses a method for treating depression or anxiety in a mammal, including a human, by administering to the mammal a CNS-penetrant NK1 receptor antagonist (e. g., a substance P receptor antagonist) in combination with an NK3 antagonist agent. It also relates to pharmaceutical compns. containing a pharmaceutically acceptable carrier, a CNS-penetrant NK1 receptor antagonist and an NK3 antagonist.

IT 174635-51-9

RL: PAC (Pharmacological activity); THU (Therapeutic use); BIOL (Biological study); USES (Uses)
 (NK1 and NK3 antagonist combination treatment for depression and anxiety)

RN 174635-51-9 CAPLUS

CN Benzeneacetic acid, α-[[[2-phenyl-4-quinolinyl)carbonyl]amino]-, methyl ester (9CI) (CA INDEX NAME)




REFERENCE COUNT:

3

THERE ARE 3 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT


L4 ANSWER 12 OF 261 CAPLUS COPYRIGHT 2004 ACS ON STN
 ACCESSION NUMBER: 2003:951029 CAPLUS
 DOCUMENT NUMBER: 140:16715
 TITLE: Preparation of azepeinoindole and pyridoindole derivatives as modulators of farnesoid X and/or orphan nuclear receptors
 INVENTOR(S): Martin, Richard; Wang, Tie-Lin; Platt, Brenton Todd; Gu, Xiao-Hui; Griffith, Ronald
 PATENT ASSIGNEE(S): X-Cepto Therapeutics, Inc., USA
 SOURCE: PCT Int. Appl., 268 pp.
 CODEN: PIXXD2
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2003099821	A1	20031204	WO 2003-US16767	20030527
<p>W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TN</p> <p>RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG</p> <p>US 2004023947 A1 20040205 US 2003-447302 20030527 PRIORITY APPL. INFO.: US 2002-383574P P 20020524 OTHER SOURCE(S): MARPAT 140:16715 GI</p>				





L4 ANSWER 12 OF 261 CAPLUS COPYRIGHT 2004 ACS ON STN (Continued)
 AB The present invention is directed to azepeinoindole and pyridoindole derivatives (shown as I and II; variables defined below; e.g. R1, 1,2,3,6-tetrahydroazepino[4,5-b]indole-5-carboxylate). These compounds were used in pharmaceutical compns. and methods for modulating the activity of farnesoid X receptor and/or orphan nuclear receptors. A farnesoid X receptor/ECREx7 co-transfection assay and a TR-FRET assay were used to establish the EC50/IC50 values for potency and percent activity or inhibition for efficacy; efficacy defines the activity of a compound relative to a high control (chenodeoxycholic acid, CDCA) or a low control (DMSO/vehicle). Most of the compds. disclosed and tested exhibited activity in at least one of the assays (EC50 or IC50 <10 μM); most showed activity at <1 μM, e.g. Pr 3-(4-fluorobenzoyl)-2-methyl-1,2,3,6-tetrahydroazepino[4,5-b]indole-5-carboxylate exhibited agonist activity <1 μM EC50 and >100 % efficacy and 8-(3-cyclopropyl-1-methylureido)-3-(4-fluorobenzoyl)-1,1-dimethyl-1,2,3,6-tetrahydroazepino[4,5-b]indole-5-carboxylic acid Et ester exhibited antagonist activity with IC50 <100 nM and 100 % inhibition. Although the methods of preparation are not claimed, 74 example preps. of I and II and characterization data for many more I and II are included. For I and II: n = 0-4; A is -N(R9)-, -O- or -S(O)t- (t = 0-2); R1 and R2 = H, alkyl, alkenyl, alkynyl, aryl, heteroaryl, cycloalkyl, heterocyclyl, aralkyl, heteroaralkyl, -OR14, -SR14, -N(R15)R16, -N(R15)S(O)2R43, -N(R17)N(R15)R16, -N(R17)N(R15)S(O)2R43, -C(O)R18, -C(O)OR14, -C(S)OR14, -C(O)SR14, -C(O)N(R15)R16, -C(O)N(R15)S(O)2R43, -C(O)N(R15)N:R16 and -C(O)N(R17)N(R15)R16; or -C(O)N(R17)N(R15)S(O)2R43; or R1 and R2, together with the atom to which they are attached, form a cycloalkyl, heterocyclyl, aryl, or heteroaryl ring. R3 is H, alkyl, alkenyl, alkynyl, cycloalkyl, aryl, aralkyl, heteroaryl, heterocyclyl, heteroaralkyl, -C(O)R10, -C(O)OR10, -S(O)2R10, -C(O)N(R11)R12, -C(O)N(R11)S(O)2R43, -C(O)N(R13)N(R11)R12, -C(O)N(R13)N(R11)S(O)2R43, -N(R13)C(O)R10, -N(R13)C(O)N(R11)R12, -N(R13)C(O)N(R11)S(O)2R43, -N(R10)C(O)N(R13)N(R11)R12, -N(R10)C(O)N(R13)N(R11)S(O)2R43, -N(R13)C(O)OR10, -P(O)OR10, or -P(O)(OR19)OR12. R4, R5, R6 and R7 = H, alkyl, alkenyl, alkynyl, aryl, heteroaryl, cycloalkyl, heterocyclyl, aralkyl, heteroaralkyl, -OR14, -SR14, -S(O)2R14, -N(R15)R16, -N(R15)S(O)2R43, -C(O)R18, -C(O)OR20, -C(O)N(R21)R22, -C(O)N(R21)S(O)2R43, -C(O)N(R42)N(R21)R22; or -C(O)N(R42)N(R21)S(O)2R43; or R4 and R5, or R4 and R6, or R4 and R7, or R5 and R6, or R5 and R7, or R6 and R7, together with the C atom to which they are attached, form a cycloalkyl, heterocyclyl, or cycloalkenyl ring, or together form a double bond and the others of R4, R5, R6 and R7 are as described above; or R6 and R7 together form an oxo, thioxo, imine, oxime or a hydrazone; or R6 and R7, together with the C atom to which they are attached, form an exocyclic double bond, and R4 and R5 are as described above. R8 = alkyl, alkenyl, alkynyl, aryl, heteroaryl, cycloalkyl, heterocyclyl, halo, pseudohalo, cyano, nitro, -C(O)OR23, -C(O)N(R24)R25, -C(O)N(R24)S(O)2R43, -C(O)R26, -OR27, -SR27, -C(S)OR23, -C(O)SR23, -N(R28)R29, and -N(R28)S(O)2R43, or two adjacent R8 groups, together with the carbons to which they are attached, form an aryl, cycloalkyl, heterocyclyl or heteroaryl; addnl. details including provisos are given in the claims
 IT RL: PAC (Pharmacological activity); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)
 (drug candidate; preparation of azepeinoindole and pyridoindole derivs. as modulators of farnesoid X and/or orphan nuclear receptors)

L4 ANSWER 12 OF 261 CAPLUS COPYRIGHT 2004 ACS ON STN (Continued)
 RN 629667-63-6 CAPLUS
 CN Azepeino[4,5-b]indole-5-carboxylic acid, 1,2,3,6-tetrahydro-3-[[[2-methoxy-2-oxo-1-(phenylmethyl)ethyl]amino]carbonyl]-, ethyl ester (9CI) (CA INDEX NAME)



REFERENCE COUNT: 10 THERE ARE 10 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT


L4 ANSWER 13 OF 261 CAPLUS COPYRIGHT 2004 ACS ON STN
 ACCESSION NUMBER: 2003:946309 CAPLUS
 DOCUMENT NUMBER: 140:128675
 TITLE: Electrochemical and Surface Study of Ferrocenyl Oligopeptides
 AUTHOR(S): Bediako-Amoa, Irene; Sutherland, Todd C.; Li, Chen-Zhong; Silerova, Roberta; Kraatz, Heinz-Bernhard
 CORPORATE SOURCE: Department of Chemistry, University of Saskatchewan, Saskatoon, SK, S7N 5C9, Can.
 SOURCE: Journal of Physical Chemistry B (2004), 108 (2), 704-714
 CODEN: JPCBFF; ISSN: 1520-6106
 PUBLISHER: American Chemical Society
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 GI



AB The syntheses and characterizations of several sym. ferrocenyl (Fc)-peptide cystamines (CSA), [Fc-Gly-CSA]2, [Fc-Ala-CSA]2, [Fc-Ala-Ala-CSA]2, [Fc-Phe-Ala-CSA]2, I (X = Gly, Ala, Ala-Ala, Phe-Ala, resp.), together with an unym. Fc-Ala-CSA-Ala-Fc (II), are reported. All systems show intermol. hydrogen bonding in solution. In the solid-state, [Fc-Gly-CSA]2 and [Fc-Ala-CSA]2 exhibit strong intermol. hydrogen bonding, as expected from solution studies, forming a network of β-helical supramol. structures. Monolayers of the Fc-peptide cystamines produced structures that show a uniform thickness of 7(2) Å but are not well ordered, leaving about 10-15% of the Au surface exposed as determined by Cu underpotential deposition. E0' values of all the monolayers are in the range of 460-510 mV. Monolayer dilution with hexanethiol caused an anodic


L4 ANSWER 13 OF 261 CAPLUS COPYRIGHT 2004 ACS on STM (Continued)
redox shift of approx. 20 mV and a slight increase in the
electron-transfer kinetics.
IT 651019-87-3P
RL: PEP (Physical, engineering or chemical process); PRP (Properties); PYP
(Physical process); SPN (Synthetic preparation); PREP (Preparation); PROC
(Process)
(preparation of disulfide-bonded ferrocenyl peptides, studies of their
solid-state structures, their monolayer assembly on Au surface and
electrochem. properties)
RN 651019-87-3 CAPLUS
CN L-Alaninamide, 2,2'-(dithiodi-2,1-ethanediyl)bis(N-(ferrocenylcarbonyl)-L-
phenylalanyl- (9CI) (CA INDEX NAME)

PAGE 1-A




L4 ANSWER 13 OF 261 CAPLUS COPYRIGHT 2004 ACS on STM (Continued)

PAGE 1-B




PAGE 2-A




REFERENCE COUNT: 94 THERE ARE 94 CITED REFERENCES AVAILABLE FOR THIS
RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 14 OF 261 CAPLUS COPYRIGHT 2004 ACS on STM
ACCESSION NUMBER: 2003:940762 CAPLUS
DOCUMENT NUMBER: 140:216323
TITLE: Quantification of ochratoxin A in foods by a stable
isotope dilution assay using high-performance liquid
chromatography-tandem mass spectrometry
Lindenmeier, Michael; Schieberle, Peter; Rychlik,
Michael
CORPORATE SOURCE: Institut fuer Lebensmittelchemie der Technischen
Universitaet Muenchen, Garching, D-85748, Germany
SOURCE: Journal of Chromatography, A (2004), 1023(1), 57-66
CODEN: JCRABY, ISSN: 0021-9673
PUBLISHER: Elsevier Science B.V.
DOCUMENT TYPE: Journal
LANGUAGE: English
AB A stable isotope dilution assay (SIDA) was developed for quantification of
the mycotoxin ochratoxin A (OTA) by using [2H5]-OTA as internal standard. The
synthesis of labeled OTA was accomplished by acid hydrolysis of unlabeled
OTA and subsequent coupling one of the products, ochratoxin A, to
[2H5]-L-phenylalanine. The mycotoxin was quantified in foods by LC-tandem
MS after extraction with buffers containing [2H5]-OTA and clean-up by immuno
affinity chromatog. or by solid phase extraction on silica. The method showed
a sufficient sensitivity with a low detection and quantification limit of
0.5 and 1.4 µg/kg, resp., and good precision in inter-assay studies
showing a CV (n = 3) of 3.6%. The anal. of certified reference materials
resulted in a low bias of 2.1% from the certified values and revealed
excellent accuracy of the new method. To prove the suitability of SIDA,
OTA was quantified in a number of food samples in which OTA was mostly
undetectable. However, three samples of raisins exceeded the legal limit
of 10 µg/kg and highlighted the need for further controlling the
contamination by the mycotoxin.
IT 666236-26-6P
RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT
(Reactant or reagent)
(determination of ochratoxin A in food by stable isotope dilution assay
using HPLC-tandem mass spectrometry)
RN 666236-26-6 CAPLUS
CN L-Phenyl-d5-alanine, N-([[(3R)-5-chloro-3,4-dihydro-8-hydroxy-3-methyl-1-
oxo-1H-2-benzopyran-7-yl]carbonyl]-, methyl ester (9CI) (CA INDEX NAME)
Absolute stereochemistry.



REFERENCE COUNT: 26 THERE ARE 26 CITED REFERENCES AVAILABLE FOR THIS
RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 15 OF 261 CAPLUS COPYRIGHT 2004 ACS on STM
ACCESSION NUMBER: 2003:913002 CAPLUS
DOCUMENT NUMBER: 139:395952
TITLE: Substituted piperazine derivatives as melanocortin
receptor ligands, and their preparation,
pharmaceutical compositions, and use
Pontillo, Joseph; Marinkovic, Dragan; Lanier, Marion
C.; Tran Joe Ahn; Arellano, Melissa; Parker, Jessica;
Nelson, Jodie; Chen, Chen; Chen, Caroline; Jiang,
Wanglong; White, Nicole; Tucci, Fabio C.
PATENT ASSIGNEE(S): Neurocrine Biosciences, Inc., USA
SOURCE: PCT Int. Appl., 153 pp.
CODEN: PIXXD2
DOCUMENT TYPE: Patent
LANGUAGE: English
FAMILY ACC. NUM. COUNT: 1
PATENT INFORMATION:
PATENT NO. KIND DATE APPLICATION NO. DATE
WO 2003094918 A1 20031120 WO 2003-US14628 20030509
W: AR, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN,
CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GE, GD, GE, GH,
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR,
LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM,
PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT,
TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW, AM, AZ, BY, KG, KZ,
MD, RU, TJ, TM
RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AT, BE, BG,
CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC,
NL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ,
GW, ML, MR, NE, SN, TD, TG
US 2004053933 A1 20040318 US 2003-434803 20030509
PRIORITY APPLN. INFO.: US 2002-379517P P 20020510
US 2002-422272P P 20021029
OTHER SOURCE(S): MARPAT 139:395952
GI




AB Comps. are disclosed, which function as melanocortin receptor ligands (no data), and which have utility in the treatment of melanocortin receptor-based disorders. The comps. have structure I [q = 1 or 2; p = 1-3; W, Q, Y, Z = CH or N, provided that ≤ 2 are N, and that when 2 are N, then the N atoms are not adjacent; Ar = (un)substituted Ph or naphthyl; X = bond, O, S, N(R6a), N(R6a)(C(O)), N(R6a)(S(O)2), N(R6a)(C(O)N(R6b)), C(O)(O), OC(O), N(R6a)(C(O)N(R6b)O), N(R6a)(C(O)N(R6b)N(R6c)), or N(R6a)(C(O)N(R6b)N(R6c)N(R6d)), (un)substituted alkyl, arylalkyl, heterocyclyl, or heterocyclylalkyl; R4a and R4b optional ring substituents selected from OH, (un)substituted alkyl, cyano, halo, alkoxy, or alkylamino; R5 = H, (un)substituted alkyl, aryl, or heterocyclyl; R6a, R6b, R6c = H, (un)substituted alkyl; R7a, R7b = optional ring substituents selected from H and (un)substituted alkyl; provided that when p = 1 then R1, R2, R3a, and R3b cannot all be H; including stereoisomers, prodrgs, and pharmaceutically acceptable salts. Pharmaceutical comps. containing I, as well as methods relating to their use, are also disclosed. Approx. 450 examples of comps. and salts were prepared, as well as various intermediates. For instance, 1-Cbz-piperazine was N-arylated with 2-fluorobenzaldehyde (53%), followed by reductive amination of the aldehyde with 2-thiophenethanamine, N-protection of the chain amino as the BOC derivative (Fmoc, 2 steps), hydrogenolysis of CBZ (35%), peptide coupling with D-N-monomer-4-chlorophenylalanine using EDC, removal of Fmoc (87%, 2 steps), another peptide coupling with N-BOC-β-alanine, and removal of BOC, to give invention compound II, isolated as the trifluoroacetate salt.

632119-09-8P

IT RL: RCT (Reactant), SPN (Synthetic preparation), PREP (Preparation), RACT (Reactant or reagent)

(intermediate), preparation of substituted piperazine derivs. as melanocortin


L4 ANSWER 15 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)
receptor ligands)
RN 626129-09-8 CAPLUS
CN 2-(1H)-isoquinolinecarboxylic acid, 3-[[[1-({4-chlorophenyl)methyl]-2-
ethoxy-2-oxoethyl]amino]carbonyl]-3,4-dihydro-, 1,1-dimethylethyl ester
(9CI) (CA INDEX NAME)



REFERENCE COUNT: 7 THERE ARE 7 CITED REFERENCES AVAILABLE FOR THIS
RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

LA ANSWER 16 OF 261 CAPLUS COPYRIGHT 2004 ACS ON STN
ACCESSION NUMBER: 2003:818297 CAPLUS
DOCUMENT NUMBER: 139:302018
TITLE: Proteasome inhibitors for treating Flaviviridae
infections
INVENTOR(S): Schubert, Ulrich; Will, Hans; Sirma, Huseyin
PATENT ASSIGNEE(S): Viromics G.m.b.H., Germany
SOURCE: PCT Int. Appl., 64 pp.
CODEN: PIXXD2
DOCUMENT TYPE: Patent
LANGUAGE: German
FAMILY ACC. NUM. COUNT: 1
PATENT INFORMATION:

L4 ANSWER 16 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)



REFERENCE COUNT: 3 THERE ARE 3 CITED REFERENCES AVAILABLE FOR THIS
RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2003084551	A1	20031016	WO 2003-DE1213	20030404
W:	AE, AG, AI, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, FB, GD, GE, GH, GM, HR, HU, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MX, MY, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SK, SL, TJ, TM, TR, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM			
RW:	GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, ZW, AM, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IS, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR, BF, BJ, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG			
DE 10316735	A1	20031120	DE 2003-10316735	20030404
PRIORITY CLAIM INFO:			DE 2002-102162Z A	20020405
AB	The invention discloses agents used in the treatment and inhibition of Flaviviridae virus infections containing, as active constituents, proteasome inhibitors. The agents, which are used for inhibiting the release, maturation and replication of Flaviviridae, contain, as active components in pharmaceutical prepn.s, compds. which inhibit the 26S proteasome. The compds. include proteasome inhibitors that influence the activities of the ubiquitin/proteasome pathway, particularly the enzymic activities of the 26S and 20S proteasome complexes. The invention is useful for antiviral therapy of Flaviviridae infections, especially			
IT	in preventing the establishment and the sustainment of a chronic hepatitis C virus infection and of hepatopathogenesis associated therewith.			
	197324-59-5			
	RL: AGR (Agricultural use); PAC (Pharmacological activity); THU (Therapeutic use); BIOL (Biological study); USES (Uses) (proteasome inhibitors for treating Flaviviridae infections)			
RN	197324-59-5 CAPLUS			
CN	Boronic acid, [(1R)-3-methyl-1-[(12S)-1-oxo-4-phenyl-2-[(2-quinolinylcarbonyl)amino]butyl]amino]butyl]- (9CI) (CA INDEX NAME)			

Absolute stereochemistry

L4 ANSWER 17 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 2003:788828 CAPLUS
 DOCUMENT NUMBER: 140:246179
 TITLE: In vitro and in vivo antipseudomonal activity, acute toxicity, and mode of action of a newly synthesized fluoroquinolonyl ampicillin derivative
 AUTHOR(S): Lin, Wen-po; Ji, Dar-der; Shiau, Chia-yang; Yang, Tee-chun; Yang, Yung-wen; Tsou, Tai-ii; Tang, Shang-tao; Chen, Chi-hsing; Liu, Yu-tien
 CORPORATE SOURCE: Institutes of Microbiology and Immunology, Preventive Medicine and Medical Science, Section of Bacteriology, Division of Clinical Pathology, National Defense Medical Center, Tri-Service General Hospital, Taipei, Taiwan
 SOURCE: Journal of Laboratory and Clinical Medicine (2003), 142(3), 158-165
 CODEN: JLCMAK; ISSN: 0022-2143
 PUBLISHER: Mosby, Inc.
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 AB Compds. N-(6,7-difluoroquinolonyl)-ampicillin (AU-1) and N-(6-fluoroquinolonyl)-ampicillin (FQ-1), synthesized by coupling of the carboxyl group of 6,7-difluoroquinolone (FP-3) and 6-fluoroquinolone (FP4), resp., with the α -amino-group of ampicillin side chain, exhibit antipseudomonal activity similar to and lower acute toxicity than that of norfloxacin, whereas neither ampicillin nor the fluoroquinolone moieties, compound FP-3 or FP4, alone have such activity. Also, AU-1 and FQ-1 are active against tested clin. isolates of *Pseudomonas aeruginosa* that are highly resistant to norfloxacin, gentamicin, or both. The therapeutic efficacies of FQ-1 and norfloxacin were assessed and compared in neutropenic mice infected with a 90% LD of *P. aeruginosa*. Mice i.p. administered FQ-1 (10 mg/kg) 4, 8, 24, and 48 h after infection had survival rates as high as 80%, comparable to those of mice treated with norfloxacin at the same dosage and dosing schedule. The study of protoplast formation revealed that FQ-1 did not inhibit cell-wall biosynthesis but did induce cell filamentation of *Bacillus subtilis* at a level close to its minimal inhibition concentration. Both AU-1 and FQ-1 were able to intercalate into the double-stranded DNA. However, that FQ-1 lost such activity after it was treated with penicillinase suggests that the lactam-ring structure in ampicillin moiety of FQ-1 was hydrolyzed by penicillinase and that the hydrolyzed structure of FQ-1 does not own DNA-intercalation activity.
 IT 190902-80-8
 RL: ADV (Adverse effect, including toxicity); DMA (Drug mechanism of action); PAC (Pharmacological activity); THU (Therapeutic use); BIOL (Biological study); USES (Uses)
 (fluoroquinolonyl ampicillin derivative in vitro and in vivo antipseudomonal activity and acute toxicity and mode of action)
 RN 190902-80-8 CAPLUS
 CN 4-Thia-1-azabicyclo[3.2.0]heptane-2-carboxylic acid, 6-[[[(2R)-[[[(6,7-difluoro-1,4-dihydro-4-oxo-3-quinolonyl)carbonyl]amino]phenylacetyl]amino]-3,3-dimethyl-7-oxo-, (2S,5R,6R)- (9CI) (CA INDEX NAME)
 Absolute stereochemistry.


L4 ANSWER 17 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)



REFERENCE COUNT: 26 THERE ARE 26 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 18 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 2003:746685 CAPLUS
 TITLE: Stoichiometric preference in copper-promoted oxidative DNA damage by ochratoxin A. [Erratum to document cited in CAL39:392373]
 AUTHOR(S): Manderville, Richard A.; Calcutt, M. Wade; Dai, Jian; Park, Gyungse; Gillman, Ivan G.; Nofle, Ronald E.; Mohammed, Abdul K.; Birincioglu, Mustafa; Dizdaroglu, Miral; Rodriguez, Henry; Akman, Steven A.
 CORPORATE SOURCE: Department of Chemistry, Wake Forest University, Winston-Salem, NC, 27109-7486, USA
 SOURCE: Journal of Inorganic Biochemistry (2003), 97(2), 249
 CODEN: JIBIDJ; ISSN: 0162-0134
 PUBLISHER: Elsevier Science Inc.
 DOCUMENT TYPE: Journal; Errata
 LANGUAGE: English
 AB An erratum.
 IT INDEXING IN PROGRESS
 IT 560134-09-0P
 RL: BSU (Biological study, unclassified); FMU (Formation, unclassified); PRP (Properties); RCT (Reactant); SPN (Synthetic preparation); BIOL (Biological study); FORM (Formation, nonpreparative); PREP (Preparation); RACT (Reactant or reagent)
 (stoichiometric preference in copper-promoted oxidative DNA damage by ochratoxin A)
 RN 560134-09-0 CAPLUS
 CN 1-Phenylalanine, N-[[[(3R)-6-[[[(2R)-2-amino-2-carboxyethyl]thio]-3,4-dihydro-5,8-dihydroxy-3-methyl-1-oxo-1H-2-benzopyran-7-yl]carbonyl]- (9CI) (CA INDEX NAME)
 Absolute stereochemistry.




L4 ANSWER 19 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 2003:737774 CAPLUS
 DOCUMENT NUMBER: 139:246221
 TITLE: Preparation of acylaminopiperidine-1-carboxamides as inhibitors of plasma kallikrein
 INVENTOR(S): Evans, David Michael
 PATENT ASSIGNEE(S): Ferring BV, Neth.
 SOURCE: PCT Int. Appl., 35 pp.
 CODEN: PIXXD2
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2003076458	A2	20030918	WO 2003-GH908	20030304
WO 2003076458	A3	20031106		


W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MY, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW, AM, AZ, BY, BG, KZ, MD, RU, TJ, TM
 RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG
 PRIORITY APPLN. INFO.: GB 2002-5527 A 20020308
 OTHER SOURCE(S): MARPAT 139:246221
 GI



AB Peptides I [R1 is H, alkyl, R4-CO, R4-O2CCH2, R5-OCO, R5-SO2 (R4 is H, alkyl, Ph and R5 is alkyl, Ph, benzyl), R2 is alkyl, cycloalkyl or cycloalkylalkyl optionally substituted with an alkyl or alkyloxy group, aralkyl optionally substituted with up to three groups chosen from F, Cl, Br, I, OH, alkyl, O-alkyl, O-benzyl, NH2, NO2, NH-acyl, CN, or CF3, or aralkyloxymethyl optionally substituted with up to three groups chosen from F, Cl, Br, OH, alkyl, or O-alkyl; or R1 and R2 together are an α -xylylene group optionally substituted on the aromatic ring by F, Cl, Br, OH, alkyl and O-alkyl; R3 is H, OH, O-alkyl or pharmaceutically-acceptable salts were prepared as inhibitors of plasma kallikrein. Thus,


L4 ANSWER 19 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)
 (2'S,2'R)-4-[2'-[2''-amino-3''-(4'''-ethoxyphenyl)propanoylamino]-3'-phenylpropanoylamino]piperidine-1-carboxamide (I; R1, R3 = H, R2 = p-ETOC6H4CH2, stereo not shown) trifluoroacetate was prepd. via peptide coupling in soln. and showed Ki = 4.5 nM for inhibition of plasma kallikrein.
 IT 599201-08-8P
 RL: PAC (Pharmacological activity); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)
 (preparation of peptidyl acylaminopiperidinecarboxamides as inhibitors of plasma kallikrein)
 RN 599201-08-8 CAPLUS
 CN 3-Isoquinolinecarboxamide, N-[(1S)-2-[[1-(aminoiminomethyl)-4-piperidinylamino]-2-oxo-1-(phenylmethyl)ethyl]-1,2,3,4-tetrahydro-, (3R)-(9CI) (CA INDEX NAME)

Absolute stereochemistry.



L4 ANSWER 20 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
 ACCESSION NUMBER: 2003:721766 CAPLUS
 DOCUMENT NUMBER: 139:323784
 TITLE: Synthesis of 1-(m-hydroxybenzyl)-substituted 1,2,3,4-tetrahydroisoquinoline-3-carboxylic acid derivatives as opioid peptide mimetics - unexpected amide bond cleavages under mild conditions
 AUTHOR(S): Mannekens, Els; Crisma, Marco; Van Cauwenberghe, Sylvia; Tourwe, Dirk
 CORPORATE SOURCE: Laboratorium voor Organische Chemie, Vrije Universiteit Brussel, Brussels, 1050, Belg.
 SOURCE: European Journal of Organic Chemistry (2003), (17), 3300-3307
 CODEN: EJOCHF; ISSN: 1434-193X
 PUBLISHER: Wiley-VCH Verlag GmbH & Co. KGaA
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 OTHER SOURCE(S): CASREACT 139:323784
 AB N-Glycyl-(1R,3S)-1-(m-hydroxybenzyl)-1,2,3,4-tetrahydroisoquinoline-3-carboxylic acid (Tic) was prepared as a Tyr-Tic dipeptide mimetic for exploration of its potential as a delta opioid receptor selective ligand. The compound was successfully obtained by a stereoselective synthesis starting from L-Tic. In the course of the reactions, unusual peptide bond cleavages were observed under mild conditions, and reaction mechanisms have been proposed.
 IT 613232-52-3P
 RL: BSU (Biological study, unclassified); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation)
 (asym. synthesis of hydroxybenzyl-substituted tetrahydroisoquinoline carboxylic acids as opioid peptidomimetics starting from Tic via stereoselective alkylation)
 RN 613232-52-3 CAPLUS
 CN L-Phenylalanine, glycyl-(1R,3S)-1,2,3,4-tetrahydro-1-[(3-hydroxyphenyl)methyl]-3-isoquinolinecarbonyl-, monoacetate (salt) (9CI) (CA INDEX NAME)
 CM 1
 CRN 613232-51-2
 CMF C28 H29 N3 O5

Absolute stereochemistry.



L4 ANSWER 20 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)

CM 2
 CRN 64-19-7
 CMF C2 H4 O2




REFERENCE COUNT: 18 THERE ARE 18 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 21 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
 ACCESSION NUMBER: 2003:678812 CAPLUS
 DOCUMENT NUMBER: 139:214483
 TITLE: Preparation of pyrrolopyrimidine derivatives as GSK-3 inhibitors
 INVENTOR(S): Kataoka, Kenichiro; Kosugi, Tomomi; Ishii, Toshihiro; Takeuchi, Takahiro; Tsutsumi, Takaharu; Nakano, Akira; Yamamoto, Yoji; Yoshioka, Noboru
 PATENT ASSIGNEE(S): Teijin Limited, Japan
 SOURCE: PCT Int. Appl., 210 pp.
 CODEN: PIXXD2
 DOCUMENT TYPE: Patent
 LANGUAGE: Japanese
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2003070730	A1	20030828	WO 2003-JP1978	20030224
W: AB, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, GR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MY, NZ, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM				
RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, NG, SN, TD, TG				

PRIORITY APPLN. INFO.: JP 2002-46129 A 20020222
 OTHER SOURCE(S): MARPAT 139:214483
 GI




AB The title pyrrolopyrimidine derivs. with general formula of I [wherein X = O or S; n = 0-2; A = N or CH; G0 = (un)substituted CH2, 2 valence group of (un)substituted benzene, furan, thiophene, pyrrole, isoxazole, cyclopentane, or cyclohexane; G1 = a single bond, CO2, CO, SO2, (un)substituted CONH, CSNH, or CONHSO2; R3 = a single bond,

L4 ANSWER 21 OF 261 CAPLUS COPYRIGHT 2004 ACS ON STN (Continued)
 (un)substituted aliph. (cyclo)hydrocarbyl, arom. hydrocarbyl, or heterocyclyl; R4 = a single bond, aliph. cyclohydrocarbyl, arom. hydrocarbyl, or heterocyclyl; G2 = H, CO₂H, CONH₂, SO₃H, or tetrazol-5-yl) and pharmaceutically acceptable salts thereof are prep'd. as glycogen synthetase kinase 3 (GSK-3) inhibitors. For example, the comp'd. II was prep'd. in a multi-step synthesis in good yield. Some of comp'ds. I showed IC₅₀ of <50 nM against GSK-3. I are useful as remedies or preventives for diseases in which GSK-3 participates such as diabetes, diabetic complications, Alzheimer's disease, neurodegenerative disease, depression, traumatic brain injury, hair loss, inflammatory diseases, cancer, immunodeficiency (no data). Formulations contg. I as an active ingredient were also described.

IT 590388-69-5P
 RL: PAC (Pharmacological activity); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)
 (drug candidate; preparation of pyrrolopyrimidine derivs. as GSK-3 inhibitors)

RN 590388-69-5 CAPLUS
 CN L-Phenylalanine, N-[(11-cyano-1,4,6,7,9,10-hexahydro-4-thioxo-8H-pyrimido[4',5':4,5]pyrrolo[1,2-d][1,4]diazepin-8-yl)carbonyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




REFERENCE COUNT: 14 THERE ARE 14 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 22 OF 261 CAPLUS COPYRIGHT 2004 ACS ON STN
 ACCESSION NUMBER: 2003:665404 CAPLUS
 TITLE: Potent and selective cathepsin L inhibitors do not inhibit human osteoclast resorption in vitro. [Erratum to document cited in CA135:120165]
 AUTHOR(S): James, Ian E.; Marquis, Robert W.; Blake, Simon M.; Hwang, Shing Mei; Gress, Catherine J.; Ru, Yu; Zembyki, Denise; Yamashita, Dennis S.; McQueney, Michael S.; Tomaszek, Thaddeus A.; Oh, Hye-Ja; Gowen, Maxine; Veber, Daniel F.; Lark, Michael W.
 CORPORATE SOURCE: Departments of Bone and Cartilage Biology, SmithKline Beecham Pharmaceuticals, King of Prussia, PA, 19406, USA
 SOURCE: Journal of Biological Chemistry (2003), 278(34), 32484
 CODEN: JBCHA3; ISSN: 0021-9258
 PUBLISHER: American Society for Biochemistry and Molecular Biology
 DOCUMENT TYPE: Journal; Errata
 LANGUAGE: English
 AB An erratum.
 IT INDEXING IN PROGRESS
 IT 350796-41-7
 RL: BSU (Biological study, unclassified); BUU (Biological use, unclassified); BIOL (Biological study); USES (Uses) (potent and selective cathepsin L inhibitors do not inhibit human osteoclast resorption in vitro (Erratum))

RN 350796-41-7 CAPLUS
 CN 8-Quinolincarboxamide, N-[(1S)-2-[(4S)-hexahydro-3-oxo-1-(2-pyridinylsulfonyl)-1H-azepin-4-yl]amino]-2-oxo-1-(phenylmethyl)ethyl]- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



L4 ANSWER 23 OF 261 CAPLUS COPYRIGHT 2004 ACS ON STN
 ACCESSION NUMBER: 2003:656752 CAPLUS
 DOCUMENT NUMBER: 139:197392
 TITLE: Preparation of N-carbamoyl nitrogen-containing fused ring compounds as mitochondrial benzodiazepine receptor (MBR) antagonists
 INVENTOR(S): Seko, Takuya; Katsumata, Seishi; Kato, Masaaki; Manako, Jun-ichi; Ohmoto, Kazuyuki
 PATENT ASSIGNEE(S): Omo Pharmaceutical Co., Ltd., Japan
 SOURCE: PCT Int. Appl., 222 pp.
 CODEN: PIXXD2
 DOCUMENT TYPE: Patent
 LANGUAGE: Japanese
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2003068753	A1	20030821	WO 2003-JP1481	20030213
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM				
RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GR, HU, IE, IT, LU, MC, NL, PT, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				

PRIORITY APPLN. INFO.: JP 2002-36340 A 20020214
 OTHER SOURCE(S): MARPAT 139:197392
 GI




AB The title comp'ds. (I) [wherein the ring A = C5-8 monocyclic carbocyclic ring or 5- to 8-membered monocyclic heterocyclic ring containing 1 or 2 N, 1 or 2 O and/or one S atom; X = CH₂, O, S, SO, SO₂; L1, L2 = a single bond, C1-4 alkylene, C2-4 alkenylene, provided that a sum total of C atoms in L1 and L2 is 3 or 4; R1, R2 = each (un)substituted C1-8 alkyl, C2-8 alkenyl, or C2-8 alkynyl; ring B, ORS, NR6R7, COR8, O2CR8, O2CNR6R7, CO2R8, CONR6R7, SR9, SOR8, SO2R8, SO2NR6R7, halo, CO₂H, cyano, NO₂, oxo, etc.; the ring B = (un)substituted C3-10 monocyclic or bicyclic carbocyclic ring or monocyclic or bicyclic heterocyclic ring containing 1 or 2 N, 1 or 2 O and/or one S atom; R5 = each (un)substituted C1-8 alkyl, C2-8 alkenyl, or C2-8 alkynyl, etc.; R6, R7 = H, -D1-D2 (wherein D1 = a single bond, CO, CO₂, or SO₂; D2 = each (un)substituted C1-8 alkyl, C2-8 alkenyl, or C2-8 alkynyl, etc.); R8 = groups listed in D2; R9 = each (un)substituted C1-8 alkyl, C2-8 alkenyl, or C2-8 alkynyl; ring B; R3 = H, ring B, each (un)substituted C1-8 alkyl, C2-8 alkenyl, or C2-8 alkynyl; R4 = H, C1-8 alkyl, C2-8 alkenyl, or C2-8 alkynyl; or NR3R4 = (un)substituted 5- to 10-membered monocyclic or bicyclic heterocyclyl containing one N atom and

L4 ANSWER 23 OF 261 CAPLUS COPYRIGHT 2004 ACS ON STN (Continued)
 optionally 1-3 N, one O, and/or one S atom; m, n = an integer of 0-4; W = O, S) were prep'd. Also disclosed are mitochondrial benzodiazepine receptor (MBR) antagonists comprising these comp'ds. I and preventives and/or remedies for diseases caused by stress which comprise the above comp'ds. I as the active ingredient. Because of having an MBR antagonistic activity and inhibiting the prodn. of neurosteroids, the comp'ds. I are useful as preventives and/or remedies for diseases caused by stress. The diseases induced, worsened, or exacerbated by stress include digestive organ disease, circulatory system disease, endocrine-metabolic disease, respiratory disease, nerve-muscular disease, skin disease, surgical disease, orthopedic disease, urinary organ-reproductive disease, gynecol. disease (gynopathy), eye disease, otolaryngol. (ear, nose and throat) disease, dental-oral surgical disease, and cancer. The digestive organ disease include functional indigestion, stomach-duodenal ulcer, ulcerative colitis, irritable bowel syndrome, biliary tract dyskinesia, esophagium, gastrotonia (atony of stomach), chronic hepatitis, and chronic pancreatitis. Thus, 250 mg Ph isocyanate was added to a soln. of 560 mg 5-(tert-butylidimethylsilyloxy)-2,3,4,5-tetrahydro-1H-1-benzazepine (prepn. given) in toluene and refluxed overnight to give, after silica gel chromatog., 655 mg 5-(tert-butylidimethylsilyloxy)-1-phenylcarbamoyl-2,3,4,5-tetrahydro-1H-1-benzazepine. The latter comp'd. (676 mg) was dissolved in 3 mL THF, treated with 2 mL 1 M Bu₄NF, and stirred for 5 h to give, after silica gel chromatog., 459 mg 5-hydroxy-1-phenylcarbamoyl-2,3,4,5-tetrahydro-1H-1-benzazepine. 1-(2,6-Dichlorophenylcarbamoyl)-2,3,4,5-tetrahydro-1H-1-benzazepine in vitro inhibited the binding of [3H]PK11195 (MBR-selective ligand) to rat brain membrane sample with K_i of 0.09 nM. A tablet formulation contg. 8-fluoro-5-(4-fluorophenyl)-1-(4-hydroxyphenylcarbamoyl)-2,3,4,5-tetrahydro-1H-1-benzazepine was described.

IT 585576-70-1P
 RL: PAC (Pharmacological activity); RCT (Reactant); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); RACT (Reactant or reagent); USES (Uses) (preparation of carbamoyl nitrogen-containing fused ring comp'ds. as mitochondrial benzodiazepine receptor antagonists for treating or preventing diseases caused by stress)

RN 585576-70-1 CAPLUS
 CN L-Phenylalanine, N-[(2,3,4,5-tetrahydro-1H-1-benzazepin-1-yl)carbonyl]-, methyl ester (9CI) (CA INDEX NAME)

Absolute stereochemistry.



REFERENCE COUNT: 40 THERE ARE 40 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 24 OF 261 CAPLUS COPYRIGHT 2004 ACS ON STM

ACCESSION NUMBER: 2003:633668 CAPLUS
DOCUMENT NUMBER: 139:197505TITLE: Preparation of aryl- or heteroaryl-containing guanidines as melanocortin-4-receptor agonists useful against disorders such as obesity or type II diabetes
INVENTOR(S): Boyce, Rustum; Chu, Daniel
PATENT ASSIGNEE(S): Chiron Corporation, USASOURCE: PCT Int. Appl., 102 pp.
CODEN: PIXXD2


DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2003066597	A2	20030814	WO 2003-US1078	20030203
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MY, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW, AM, AZ, BY, BG, KZ, MD, RU, TJ, TM			
RW:	GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG			
US 2003195187	A1	20031016	US 2003-351574	20030127
PRIORITY APPL. INFO.:			US 2002-353188P	P 20020204
			US 2003-351574	A 20030127
OTHER SOURCE(S):			MARPAT 139:197505	
GI				



AB A variety of small, guanidino group-containing mols. (I; A1-A2-A3-A4;


L4 ANSWER 24 OF 261 CAPLUS COPYRIGHT 2004 ACS ON STM (Continued)

variables defined below, e.g. (3S)-N'-[4-(3,4-dihydroquinolin-1(2H)-ylcarbonyl)phenyl]-3-methyl-N-[(1S,2S,3S,5R)-2,6,6-trimethylbicyclo[3.1.1]hept-3-yl]piperazine-1-carboxinidamide (shown as I) capable of acting as MC4-R agonists are provided. The compds. are useful in treating MC4-R mediated diseases and may be formulated into pharmaceutical formulations and compns. Although the methods of prepns. are not claimed, several example prepns. of I and a no. of example prepns. of intermediates are included; 131 addnl. examples of I are tabulated with mass spectral characterization data. Some of the I have -log EC50 values above -apprx.3. Compds. I showed beneficial effects in in vivo studies on energy intake, body wt., hyperinsulinemia, and glucose levels in male 9-10 wk old ob/ob mice that display early onset of obesity, insulin resistance and diabetes due to leptin deficiency. For I: A1 = R1'R2'NC(=NR3')NR4'-, R1'R2'NC(=NR3')R4')N-; R1' = H, and (un)substituted alkyl, alkenyl, alkynyl, cycloalkyl, aryl, heteroaryl, heterocyclyl, arylalkyl, heteroarylalkyl, and cycloalkylalkyl; R2' = (un)substituted alkyl, alkenyl, alkynyl, cycloalkyl, aryl, heteroaryl, heterocyclyl, arylalkyl, heteroarylalkyl, and cycloalkylalkyl; or R1' and R2', together with the N to which they are bound, form a (un)substituted heterocyclyl or heteroaryl; R3' = (un)substituted aryl, alkyl, alkenyl, alkynyl, cycloalkyl, heteroaryl, heterocyclyl, heterocyclylalkyl, arylalkyl, heteroarylalkyl, and cycloalkylalkyl; R4' = H, and (un)substituted alkyl, alkenyl, alkynyl, cycloalkyl, heterocyclylalkyl, cycloalkylalkyl, aryl, heteroaryl, heterocyclyl, arylalkyl, and heteroarylalkyl. A2 = (un)substituted aryl and heteroaryl; A3 is a covalent bond such that A2 is directly bonded to A4, or A3 is a linking group O, S, -NRa-, -C(O)-, -C(O)O-, -NRaC(O)-, -SO2NRa-, -C(S)-, -C(O)S-, -P(O)Rb-, -SO2-, and -S(O)-, wherein if A3 is a linking group, then it is bonded to A2 and A4 in a configuration A2-O-A4, A2-S-A4, A2-NRa-A4, A2-C(O)-A4, A2-C(O)O-A4, A4-C(O)O-A2, A2-NRaC(O)-A4, A4-NRaC(O)-A2, A2-SO2NRa-A4, A4-SO2NRa-A2, A2-C(S)-A4, A2-C(O)S-A4, A4-(C(O)S)-A2, A2-(P(O)Rb)-A4, A2-SO2-A4, and A2-S(O)-A4 provided that if A3 is a linking group with the configuration A4-NRaC(O)-A2, then A2 is not a (un)substituted Ph and is not a (un)substituted 6-membered N-contg. heteroaryl. A4 = (un)substituted arylalkyl, heteroarylalkyl, aryl, heteroaryl, heterocyclyl, cycloalkyl, heterocyclylalkyl, cycloalkylalkyl, alkenyl, alkynyl, and alkyl; Ra = H, and (un)substituted arylalkyl, heteroarylalkyl, aryl, heteroaryl, heterocyclyl, cycloalkyl, heterocyclylalkyl, cycloalkylalkyl, alkenyl, alkynyl, and alkyl; Rb = (un)substituted arylalkyl, heteroarylalkyl, aryl, heteroaryl, heterocyclyl, cycloalkyl, heterocyclylalkyl, cycloalkylalkyl, alkenyl, alkynyl, and alkyl.

IT 582297-17-AP: (3R)-N-[(1R)-1-[[4-(chlorophenyl)methyl]-2-[[4-[[[(Z)-[(3R,5S)-3,5-dimethylpiperazin-1-yl]](1S,2S,3S,5R)-2,6,6-trimethylbicyclo[3.1.1]hept-3-yl]imino]methyl]amino]phenyl]amino]-2-oxoethyl]-1,2,3,4-tetrahydroisoquinoline-3-carboxamide
RL: PAC (Pharmacological activity); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)
(drug candidate; preparation of aryl- or heteroaryl-containing guanidines as melanocortin-4-receptor agonists useful against disorders such as obesity or type II diabetes)

RN 582297-17-4 CAPLUS
CN 3-isoquinolinecarboxamide, N-[(1R)-1-[[4-(chlorophenyl)methyl]-2-[[4-[[[(Z)-[(3R,5S)-3,5-dimethyl-1-piperazinyl]](1S,2S,3S,5R)-2,6,6-trimethylbicyclo[3.1.1]hept-3-yl]imino]methyl]amino]phenyl]amino]-2-oxoethyl]-1,2,3,4-tetrahydro-, (3R)- (9CI) (CA INDEX NAME)

L4 ANSWER 24 OF 261 CAPLUS COPYRIGHT 2004 ACS ON STM (Continued)



L4 ANSWER 25 OF 261 CAPLUS COPYRIGHT 2004 ACS ON STM

ACCESSION NUMBER: 2003:610477 CAPLUS
DOCUMENT NUMBER: 139:169275TITLE: Trojan inhibitors for use in treatment of viral infections
INVENTOR(S): Schubert, Ulrich; Schubert, Evelyn; Tessmer, Uwe;

PATENT ASSIGNEE(S): Viromics G.m.b.H., Germany

SOURCE: PCT Int. Appl., 78 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: German

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2003064453	A2	20030807	WO 2003-DE265	20030127
WO 2003064453	A3	20040212		
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MY, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW, AM, AZ, BY, BG, KZ, MD, RU, TJ, TM			
RW:	GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG			
DE 10304203	A1	20031204	DE 2003-10304202	20030127
PRIORITY APPL. INFO.:			DE 2002-10203862 A	20020127
			DE 2002-10204210 A	20020127
			DE 2002-10209064 A	20020228

AB The invention relates to active inhibitors - Trojan inhibitors (TI) - and the use thereof in the form of specifically shaped Trojan proteasome-inhibitors (TPI) or Trojan assembling-inhibitors (TAI), such as proteasome-and-assembling-inhibitors which are, initially, inactive and are only activated in the target cell by means of a protease specific for the target cell. According to the invention, said inhibitor can be used in the treatment of viral infections, whereby a virus-specific protease is expressed, particularly in HIV-infections and AIDS-therapy, and optionally in the inhibition of the release, maturing and replication of filo viruses, and in the treatment and prevention of viral haemorrhagic fever (activated by Ebola or Marburg-viruses) and in the therapy of tumoral diseases, whereby the tumor cells are characterised by a specific protease.

IT 179324-59-5, eS 325


RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)
(Trojan inhibitors for use in treatment of viral infections)

RN 179324-59-5 CAPLUS

CN Boronic acid, [(1R)-3-methyl-1-[(2S)-1-oxo-4-phenyl-2-[(2-quinolinylcarbonyl)amino]butyl]amino]butyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

L4 ANSWER 25 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)




L4 ANSWER 26 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
 ACCESSION NUMBER: 2003:591206 CAPLUS
 DOCUMENT NUMBER: 139:145837
 TITLE: Substrates for monitoring Staphylococcus aureus cysteine peptidase activity and use for screening antibacterial agents
 INVENTOR(S): Ramjee, Manoj Kumar
 PATENT ASSIGNEE(S): Amura Therapeutics Limited, UK
 SOURCE: PCT Int. Appl., 96 pp.
 CODEN: PIXXD2
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2003062267	A2	20030731	WO 2003-GB120	20030116
WO 2003062267	A3	20030904		
W:	AE, AG, AI, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MY, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM			
RW:	GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GM, GQ, GW, ML, MR, NE, SN, TD, TG			
PRIORITY APPLN. INFO.:			GB 2002-1040 A 20020117	
			GB 2002-16508 A 20020716	

OTHER SOURCE(S): MARPAT 139:145837
 AB The invention provides peptide substrates for monitoring Staphylococcus aureus cysteine peptidase activity and use for screening antibacterial agents. Comps. of general formula (I): (A) A compound of general formula (I) Y-AA1-AA2-AA3-AA4-R-Glu-NH₂, Y-A1-A2-A3-A4-R-Glu-NH₂ where Y and R are reporter groups; A1 is either the D-isomer or the L-isomer form of isoleucine; each of A2 and A3 is independently Ala or 2-aminobutyric acid (Abu); A4 is Pro, L-3-hydroxyproline (Pro(3-OH)), L-4-hydroxyproline (Pro(4-OH)), L-tetrahydroisoquinoline-1-carboxylic acid (THIQ), L-tetrahydroisoquinoline-3-carboxylic acid (Tic), L-pipecolic acid (Pip), L-amino-1-cyclopentane carboxylic acid (1-ACP) or D-Pro; and any one of A1, A2, A3 and A4 can be replaced by any other amino acid group (Xaa); and the other protein based substrates are cleavable by Staphylococcus sp. extracellular cysteine protease and are therefore likely to be of use in monitoring enzyme activity for purposes such as investigating candidate enzyme modulators and screening for therapeutic agents.
 IT 572919-07-4
 RL: ARG (Analytical reagent use); PRP (Properties); THU (Therapeutic use); ANST (Analytical study); BIOL (Biological study); USES (Uses) (substrate sequence; substrates for monitoring Staphylococcus aureus cysteine peptidase activity and use for screening antibacterial agents)
 RN 572919-07-4 CAPLUS
 CN L-α-Glutamine, N-(2-aminobenzoyl)-L-leucyl-L-alanyl-L-alanyl-(3S)-1,2,3,4-tetrahydro-3-isoquinolinecarboxyl-3-nitro-L-tyrosyl- (9CI) (CA INDEX NAME)

L4 ANSWER 26 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)


Absolute stereochemistry.



L4 ANSWER 27 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
 ACCESSION NUMBER: 2003:591007 CAPLUS
 DOCUMENT NUMBER: 139:149922
 TITLE: Preparation of piperazinyl amino acid derivatives as melanocortin receptor agonists
 INVENTOR(S): Backer, Ryan Thomas; Collado Cano, Ivan; De Frutos Garcia, Oscar; Doecke, Christopher William; Fisher, Matthew Joseph; Kuklish, Steven Lee; Mancuso, Vincent; Martinelli, Michael John; Mullaney, Jeffrey Thomas; Ornstein, Paul Leslie; Xie, Chaoyu
 PATENT ASSIGNEE(S): Eli Lilly and Company, USA
 SOURCE: PCT Int. Appl., 222 pp.
 CODEN: PIXXD2
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2003061660	A1	20030731	WO 2003-US33	20030121
W:	AE, AG, AI, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW, AM, AZ, BY			
RW:	GH, GM, KE, LS, MW, MZ, SD, SI, SL, SZ, TZ, UG, ZM, ZW, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG			
PRIORITY APPLN. INFO.:			US 2002-351200P P 20020123	
OTHER SOURCE(S):			CASREACT 139:149922; MARPAT 139:149922	

GI




AB The invention relates to melanocortin receptor (MC-R) agonists I [LL1 = H₂ or oxo; E = O, S, NR1b, SO, SO₂, CR₉, CR₉₂, where R1b = H, alkyl, alkylsulfonyl, etc. and R₉ = H, alk(en)(yn)yl, alkanoyl, Ph, (hetero)aryl, or R₉ may combine with adjacent R₁ to form a carbocycle; X = CH₂ or CH₂CH₂; Y = (CH₂)₀₋₂; the ring containing E may have a double bond; T = substituted (tetrahydro)isoquinolinyl, dihydroisoindolinyl, or piperazinyl; n = 0-8; R₁ = H, alkyl, (D)cycloalkyl, aryl, carbalkoxy, etc.; R_{1a} = H, (cyclo)alkyl, (D)(hetero)aryl, aminoalkyl, etc.; R₂ = H, alkyl, alkylcarbonyl, (D)phenyl, (D)cycloalkyl, or oxo adjacent to N attached to the ring containing E; p = 0-4; R₃ = (un)substituted Ph, aryl, or thienyl; R₄ = H, alkyl, alkoxyalkyl, alkanoyl, or carbalkoxy] or their

L4 ANSWER 27 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)
pharmaceutically-acceptable salts or stereoisomers, which are useful in the treatment of obesity, diabetes, and male and/or female sexual dysfunction. Compds. I comprise three domains, i.e., a piperazinyl fragment, an amino acid, and a radical $CL_1(CH_2)_n-T$. Thus, N-[1-(4-chlorobenzyl)-2-(4-(4-isobutyl-1-isopropylpiperidin-4-yl)piperazin-1-yl)-2-oxoethyl]-2-(2,3-dihydro-1H-indol-1-yl)acetamide TFA salt was prep'd. via acylation of the piperazine moiety and assayed for treatment of sexual dysfunction in rat models (MC4 Ki = 9 nM, MC4 EC50 = 4.2 nM).

IT 252008-71-2P
RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)
(preparation of piperazinyl amino acid derivs. as melanocortin receptor agonists)

RN 252008-71-2 CAPLUS
CN 2(1H)-Isoquinolinecarboxylic acid, 3-[[[1R]-1-carboxy-2-(4-chlorophenyl)ethyl]amino]carbonyl]-3,4-dihydro-, 2-(1,1-dimethylethyl) ester, (3R) - (9CI) (CA INDEX NAME)

Absolute stereochemistry.




REFERENCE COUNT: 5 THERE ARE 5 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 28 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
ACCESSION NUMBER: 2003:598812 CAPLUS
DOCUMENT NUMBER: 139:133836
TITLE: Preparation of 4-aminoazepan-3-ones as protease inhibitors
INVENTOR(S): Marquis, Robert Wells; Ru, Yu; Veber, Daniel Frank; Cummings, Maxwell David; Thompson, Scott Kevin; Yamashita, Dennis Shinji
PATENT ASSIGNEE(S): Smithkline Beecham Corporation, USA
SOURCE: U.S. Pat. Appl. Publ. 126 pp., Cont.-in-part of U.S. Ser. No. 593,845, abandoned.
CODEN: USXXCO
DOCUMENT TYPE: Patent
LANGUAGE: English
FAMILY ACC. NUM. COUNT: 4
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 2003144175	A1	20030731	US 2001-881334	20010614
WO 2000038687	A1	20000706	WO 1999-US30730	19991221
W: AE, AL, AU, BA, BB, BG, BR, CA, CN, CZ, EE, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KP, KR, LC, LK, LR, LT, LV, MA, MG, MK, MN, MX, NO, NZ, PL, RO, SG, SI, SK, SL, TR, TT, UA, US, UZ, VN, YU, ZA, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM				
RW: GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG				
EP 1384713	A1	20040128	EP 2003-76211	19991221
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, FI, RO, CY				
ZA 2001004208	A	20020523	ZA 2001-4208	20010523
WO 2002017924	A1	20020307	WO 2001-US27178	20010831
W: AB, AG, AI, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM				
RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
AU 2001085983	A5	20020313	AU 2001-86983	20010831
EP 1320370	A1	20030625	EP 2001-966474	20010831
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR				
JP 2004509083	T2	20040325	JP 2002-522897	20010831
US 2004002487	A1	20040101	US 2003-404716	20030401
PRIORITY APPLN. INFO.:				
US 1998-113636P P 19981223				
US 1999-164581P P 19991110				
WO 1999-US30730 A2 19991221				
US 2000-593845 B2 20000614				
EP 1999-963112 A3 19991221				
US 2000-653815 A2 20000901				
US 2001-881334 A2 20010614				
WO 2001-US27178 W 20010831				

GI

L4 ANSWER 28 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)




AB Aminoazepanones I [R1 = alkanoyl, amino-, alkoxy-, or alkylthioalkanoyl, etc.; R2 = H, alkyl, cycloalkyl, cycloalkylalkyl, (thio)acyl, alkylsulfonyl, etc.; R3 = H, alkyl, cycloalkyl, cycloalkylalkyl, aryl, etc.; R4 = H, alkyl, arylalkyl, etc.] or their pharmaceutically-acceptable salts were prepared as protease inhibitors, including cathepsin K, for treating diseases of excessive bone loss or cartilage or matrix degradation, gingival disease, arthritis, Paget's disease, hypercalcemia of malignancy, and metabolic bone disease. Thus, compound II (Cbz = benzyloxycarbonyl) was prepared by a multistep procedure.

IT 281217-12-7P
RL: PAC (Pharmacological activity); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)
(preparation of (acylamino)azepanones as protease inhibitors)

RN 281217-12-7 CAPLUS
CN 8-Quinolinesulfonyl-1H-azepin-4-yl]amino]-2-oxo-1-(phenylmethyl)ethyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




L4 ANSWER 29 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
ACCESSION NUMBER: 2003:598842 CAPLUS
DOCUMENT NUMBER: 140:218058
TITLE: Solid supported parallel synthesis of dimer libraries
AUTHOR(S): Subra, Gilles; Ambard, Muriel; Durand, Philippe; Komesil, Sylvianne; Renaut, Patrice; Martinez, Jean
CORPORATE SOURCE: Laboratoire des Aminoacides, Peptides et Proteines, Faculte de Pharmacie, UMR 5810, Montpellier, 34060, Fr.
SOURCE: Peptides 2000, Proceedings of the European Peptide Symposium, 26th, Montpellier, France, Sept. 10-15, 2000 (2001), Meeting Date 2000, 973-974. Editor(s): Martinez, Jean; Fehrentz, Jean-Alain. Editions EDK: Paris, Fr.
CODEN: 69EDWK; ISBN: 2-84254-048-4
DOCUMENT TYPE: Conference
LANGUAGE: English

AB A symposium report. Dimer libraries, particularly the JMV 1783 dimer library, were synthesized using lysine as a central template via the Multipin technol. The core of the compds. in the dimer library synthesis is a diamino acid template which is linked to the Synphase crown by a Rink amide type linker. Eleven libraries generated a family of 650 members, of which 10 showed a growth hormone binding inhibition of > 80% at 10-5 M.

IT 664335-91-5P, JMV 1946
RL: CPN (Combinatorial preparation); CMBI (Combinatorial study); PREP (Preparation)
(solid supported parallel synthesis of peptide dimer libraries and their growth factor hormone agonist activity)

RN 664335-91-5 CAPLUS
CN 6-Quinolinesulfonyl-1H-azepin-4-yl]amino]-2-oxo-1-(phenylmethyl)ethyl]- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



PAGE 1-A

L4 ANSWER 29 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)


PAGE 1-B



REFERENCE COUNT: 5 THERE ARE 5 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 30 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
 ACCESSION NUMBER: 2003:509769 CAPLUS
 DOCUMENT NUMBER: 140:164185
 TITLE: Amide bond cleavages under unusual mild conditions in N-pivaloyl-1,2,3,4-tetrahydroisoquinoline-3-carboxylic acid (N-pivaloyl-Tic) compounds: A mechanistic study
 AUTHOR(S): Mannekens, Els; Tourwe, Dirk; Crisma, Marco
 CORPORATE SOURCE: Laboratorium voor Organische Chemie, Vrije Universiteit Brussel, Brussels, B-1050, Belg.
 SOURCE: Peptides 2000, Proceedings of the European Peptide Symposium, 26th, Montpellier, France, Sept. 10-15, 2000 (2001), Meeting Date 2000, 827-828. Editor(s): Martinez, Jean; Fehrentz, Jean-Alain. Editions EDK: Paris, Fr.
 CODEN: 69EDWK; ISBN: 2-84254-048-4
 DOCUMENT TYPE: Conference
 LANGUAGE: English
 AB A symposium report. A mechanistic study has been conducted for two cases of amide cleavages during the deprotection of the N-pivaloyl protecting group of 1-[3-(benzyloxy)benzyl]-1,2,3,4-tetrahydroisoquinoline-3-carboxylic acid derivs. Mechanisms are proposed for cleavage of the pivaloyl-Tic amide bond and the Tic-Phe amide bond.
 IT 613232-44-3
 RL: RCT (Reactant); RACT (Reactant or reagent) (mechanism of amide bond cleavage of pivaloyltetrahydroisoquinolinecarbonylic acid compds.)
 RN 613232-44-3 CAPLUS
 CN L-Phenylalanine, N-[[[(1R,3S)-2-(2,2-dimethyl-1-oxopropyl)-1,2,3,4-tetrahydro-1-[[3-(phenylmethoxy)phenyl]methyl]-3-isoquinolinyl]carbonyl]-, phenylmethyl ester (9CI) (CA INDEX NAME)


Absolute stereochemistry.



REFERENCE COUNT: 2 THERE ARE 2 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 31 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
 ACCESSION NUMBER: 2003:509722 CAPLUS
 DOCUMENT NUMBER: 140:193456
 TITLE: Opioid activity profiles of TIPP-related peptides containing 2'-hydroxy,6'-methyltyrosine (Hmt) in place of Tyr
 AUTHOR(S): Schiller, Peter W.; Berezowska, Irena; Weltrowska, Grazyna; Lemieux, Carole; Chung, Nga N.
 CORPORATE SOURCE: Laboratory of Chemical Biology and Peptide Research, Clinical Research Institute of Montreal, Montreal, QC, H2W 1R7, Can.
 SOURCE: Peptides 2000, Proceedings of the European Peptide Symposium, 26th, Montpellier, France, Sept. 10-15, 2000 (2001), Meeting Date 2000, 731-732. Editor(s): Martinez, Jean; Fehrentz, Jean-Alain. Editions EDK: Paris, Fr.
 CODEN: 69EDWK; ISBN: 2-84254-048-4
 DOCUMENT TYPE: Conference
 LANGUAGE: English
 AB Syntheses and in vitro opioid activity profiles of the Hmt1-analogs of TIPP-related peptides were presented. Results indicate that Hmt1-analogs of TIPP-related peptides may have unexpected opioid activity profiles, possibly due to the hydrogen bonding capability of the 2'-OH group of Hmt.
 IT 660850-07-7
 RL: BSU (Biological study, unclassified); PKT (Pharmacokinetics); PRP (Properties); BIOL (Biological study) (opioid activity profiles of TIPP-related peptides containing 2'-hydroxy,6'-methyltyrosine (Hmt) in place of Tyr)
 RN 660850-07-7 CAPLUS
 CN L-Phenylalaninamide, 2-hydroxy-6-methyltyrosyl-(3S)-1,2,3,4-tetrahydro-3-isoquinolinecarboxyl-L-phenylalanyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




REFERENCE COUNT: 4 THERE ARE 4 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT


L4 ANSWER 32 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
 ACCESSION NUMBER: 2003:509658 CAPLUS
 DOCUMENT NUMBER: 140:157610
 TITLE: A structure-activity study of dynorphin(1-13)-peptide amides. Synthesis of analogs with unusual amino acids in positions 0, 2, 3, 4
 AUTHOR(S): Bobrova, Irina; Goodman, Murray; DeHaven, Robert N.; Daubert, Jeffrey D.; Johansson, Lars Erik; Neumuller, Magnus; Terenius, Lars
 CORPORATE SOURCE: Institute of Organic Synthesis, Riga, LV-1006, Latvia
 SOURCE: Peptides 2000, Proceedings of the European Peptide Symposium, 26th, Montpellier, France, Sept. 10-15, 2000 (2001), Meeting Date 2000, 603-604. Editor(s): Martinez, Jean; Fehrentz, Jean-Alain. Editions EDK: Paris, Fr.
 CODEN: 69EDWK; ISBN: 2-84254-048-4
 DOCUMENT TYPE: Conference
 LANGUAGE: English
 AB A series of dynorphin analogs was synthesized by solid-phase methodol. using Fmoc-amino acid pentafluorophenyl esters. The analogs were synthesized with the following modifications: the amino acid residue at position 2 was replaced by amino acids Phe, cPrAla (8-cyclopropyl-Ala), Tic(tetrahydroisoquinoline), Oic (octahydroindole-2-carboxylic acid), NaI2 (2-naphthylalanine); N-terminal extension with lys0; 3 Gly3 was substituted with cPrAla,Tic; exchange of Phe4 for Tic; and Gly2-Gly3 was replaced by cPrAla-Phe, Tic-Phe, Oic-Phe and NaI2-Phe in analogs with double replacement. The new dynorphin analogs were κ-selective. All of them retained modest μ activity and were essentially inactive at the δ opioid receptor. The presence of cPrAla in position 3 resulted in a compound with the highest κ-activity. Double substitution of Gly-Gly by Tic-Phe afforded an analog which exhibited modest δ-activity.
 IT 656256-53-0
 RL: BSU (Biological study, unclassified); PRP (Properties); BIOL (Biological study) (structure-activity study of dynorphin(1-13)-peptide analogs with unusual amino acids in positions 0, 2, 3, 4)
 RN 656256-53-0 CAPLUS
 CN 1-13-Dynorphin A (swine), 3-[(3S)-1,2,3,4-tetrahydro-3-isoquinolinecarboxylic acid]-13-L-lysineamide- (9CI) (CA INDEX NAME)

Absolute stereochemistry.


PAGE 1-A



PAGE 1-B



PAGE 2-A



REFERENCE COUNT: 5 THERE ARE 5 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

ACCESSION NUMBER: 2003:491183 CAPLUS
DOCUMENT NUMBER: 139:69525

TITLE: Synthesis of tetrahydrocarbazole derivatives for use as ligands for G-protein coupled receptors and antagonists of gonadotropin-releasing hormone for treatment of disease

INVENTOR(S): Kopitz, Marcus; Mulin, Hans Peter; Shaw, Ken; Hess-Stump, Holger; Paulini, Klaus

PATENT ASSIGNEE(S): Zentaris Ag, Germany
SOURCE: PCT Int. Appl., 114 pp.



DOCUMENT TYPE: CODEN: PIXXD2

LANGUAGE: Patent

FAMILY ACC. NUM. COUNT: 1 German

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2003051837	A2	20030626	WO 2002-EP14344	20021216
WO 2003051837	A3	20040226		
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MY, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW, AM, AZ, BY, BG, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
DE 10164564	A1	20030626	DE 2001-10164564	20011214
US 2003232873	A1	20031218	US 2002-319833	20021216
PRIORITY APPLN. INFO.: DE 2001-10164564 A 20011214 US 2001-341876P P 20011221				
OTHER SOURCE(S): MARPAT 139:69525				
GI				



AB The invention relates to novel tetrahydrocarbazole derivs. [e.g., (I)] which act as ligands for G-protein coupled receptors (GPCR), especially as antagonists of gonadotropin-releasing hormone (GnRH), and pharmaceutical composition containing them. Furthermore, the invention relates to the administration of tetrahydrocarbazole derivs. for the treatment of pathol. conditions mediated by GPCR, especially for the inhibition of GnRH, to mammals, especially humans, requiring such treatment, and to the use of tetrahydrocarbazole derivs. for producing a pharmaceutical agent for treating pathol. conditions mediated by GPCR, especially for the inhibition of GnRH. Limited synthesis of intermediate materials is given, with many tables of products exemplified by general synthesis steps. Thus, beginning from 4,4-ethylenedioxcyclohexanone and phenylhydrazine, I was prepared in seven generalized steps. In in vitro tests with alpha T3-1 cells, I had IC50 for human GnRH of 1.5 x 10-8 M, with Ca2+ release of 4.5 x 10-8 M.

IT 548754-89-89

RL: BSU (Biological study, unclassified); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)

(preparation of tetrahydrocarbazole derivs. for use as ligands for G-protein coupled receptors and antagonists of gonadotropin-releasing hormone for treatment of disease)

RN 548754-89-8 CAPLUS

CN Carbanic acid, [(1S,2S)-1-[[[(3R)-3-[[[(1S)-1-(aminocarbonyl)-3-phenylpropyl]amino]carbonyl]-6,8-dichloro-2,3,4,9-tetrahydro-1H-carbazol-3-yl]amino]carbonyl]-2-methylbutyl]-, phenylmethyl ester (9CI) (CA INDEX NAME)

Absolute stereochemistry.


L4 ANSWER 34 OF 261 CAPIUS COPYRIGHT 2004 ACS on STN
 ACCESSION NUMBER: 2003:459828 CAPIUS
 DOCUMENT NUMBER: 139:175029
 TITLE: Binding of Ochratoxin A Derivatives to Human Serum Albumin
 AUTHOR(S): Perry, Jennifer L.; Il'ichev, Yuri V.; Kempf, Valerie R.; McClendon, Jamal; Park, Gyungse; Manderville, Richard A.; Rueker, Florian; Dockal, Michael; Simon, John D.
 CORPORATE SOURCE: Department of Chemistry, Duke University, Durham, NC, 27708, USA
 SOURCE: Journal of Physical Chemistry B (2003), 107(27), 6644-6647
 CODEN: JPCBPK; ISSN: 1520-6106
 PUBLISHER: American Chemical Society
 DOCUMENT TYPE: Journal
 LANGUAGE: English

AB Ochratoxins are fungal metabolites known to contaminate human and animal feed. Ochratoxin A (OTA) is the most widespread form of the toxins and is believed to be responsible for human renal diseases. For the majority of its lifetime within the body, OTA remains bound to the plasma protein human serum albumin (HSA). In this paper, the binding of three OTA derivs. (ochratoxin B (OTB), ochratoxin hydroquinone (OHQ), and O-methylated OTA (MOA)) to HSA is examined using optical spectroscopy. The binding consts. decrease as follows: OTA2- (5.2+105 M-1) > OTB2- (1.8+106 M-1) > OHQ2- approx. OHQ- (2.2+105 M-1) > MOA- (3+104 M-1). Studies of the binding of OTB, OHQ, and MOA to recombinant proteins corresponding to the domains of HSA reveal binding to all domains but with different affinities. Similar to OTA, all derivs. exhibit the largest binding constant for domain 2. These ligands are displaced by 2,3,5-triiodobenzoate (TIB), indicating they share a common binding site and bind to Sudlow Site I within domain 2 of HSA. Derivs. with ionizable phenolic protons exhibit a decreased pKa by as much as two units upon interaction with HSA. The magnitude of the change in pKa observed upon binding decreases in the order OTA > OTB > OHQ. These data suggest a model in which the monoanions of OTA, OTB, and OHQ undergo deprotonation by an arginine within domain 2 upon binding to HSA. The difference in binding constant for the three dianions studied results from the stabilization of the dianion by the surrounding protein matrix.

IT 205034-32-0D, serum albumin complexes
 RL: BSU (Biological study, unclassified); BIOL (Biological study) (binding of ochratoxin A derivs. to human serum albumin)
 RN 205034-32-8 CAPIUS
 CN 1-Phenylalanine, N-[[[(3R)-3,4-dihydro-5,8-dihydroxy-3-methyl-1-oxo-1H-2-benzopyran-7-yl]carbonyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

L4 ANSWER 34 OF 261 CAPIUS COPYRIGHT 2004 ACS on STN (Continued)



REFERENCE COUNT: 54 THERE ARE 54 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 35 OF 261 CAPIUS COPYRIGHT 2004 ACS on STN
 ACCESSION NUMBER: 2003:390845 CAPIUS
 DOCUMENT NUMBER: 138:385216
 TITLE: Preparation of etoposide amino acid analogs as DNA topoisomerase II inhibitors
 INVENTOR(S): Lee, Kuo-Hsiung; Xiao, Zhiyan; Bastow, Kenneth F.
 PATENT ASSIGNEE(S): The University of North Carolina At Chapel Hill, USA
 SOURCE: U.S., 17 pp.
 CODEN: USXXAM
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 2
 PATENT INFORMATION:


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 6566393	B1	20030520	US 2002-177147	20020621
US 2004006126	A1	20040108	US 2003-349351	20030122
WO 2004008559	A2	20031231	WO 2003-US19629	20030620

W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ

RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG

PRIORITY APPLN. INFO.: US 2002-177147 A1 20020621
 US 2003-349351 A 20030122

OTHER SOURCE(S): MARPAT 138:385216
 GI




AB Etoposide amino acid analogs I (X = O, S, NH, CO, CH:N, CH2NH; R1 = covalent linkage between X and Y, alkyl, alkenyl, (un)substituted Ph; Y = NHCO, CONH; Z = CHR2(CH2)nR3, R2 = CO2H, NH2, ester, etc., R3 = alkyl, alkenyl, aryl, n = 0-2; D = CH2OC(O), CH2OC(=CH2), CH2CH2C(O), CH2OCH2, CH2OC(S), CH2O(SO2)OCH2, etc.) were prepared as DNA topoisomerase II inhibitors. Thus, 4'-O-demethyl-4β-(4'''-(methyl-L-tyrosine-N-carbonyl)anilino)-4-desoxy-podophyllotoxin (II) and 4'-O-demethyl-4β-

L4 ANSWER 35 OF 261 CAPIUS COPYRIGHT 2004 ACS on STN (Continued)
 [4'''-(methyl-L-tryptophan-N-carbonyl)anilino]-4-desoxypodophyllotoxin were prepd. from podophyllotoxin and their pharmaceutical activity evaluated. The antitumor ED50 of II against A 549 human cell line was 2.4 μM.
 IT 527678-17-7P
 RL: PAC (Pharmacological activity); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)
 (preparation of etoposide amino acid analogs as DNA topoisomerase II inhibitors)

RN 527678-17-7 CAPIUS
 CN L-Tyrosine, N-[[[(5S,5aR,8aR,9R)-5,5a,6,8,8a,9-hexahydro-9-(4-hydroxy-3,5-dimethoxyphenyl)-8-oxofuro[3',4':6,7]naphtho[2,3-d]-1,3-dioxol-5-yl]carbonyl]-, methyl ester (9CI) (CA INDEX NAME)

Absolute stereochemistry.



REFERENCE COUNT: 8 THERE ARE 8 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 36 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
 ACCESSION NUMBER: 2003:38943 CAPLUS
 DOCUMENT NUMBER: 139:392373
 TITLE: Stoichiometric preference in copper-promoted oxidative DNA damage by ochratoxin A
 AUTHOR(S): Manderville, Richard A.; Wade Calcutt, M.; Dai, Jian; Park, Gyungse; Gillman, Ivan G.; Nofle, Ronald E.; Mohammed, Abdul K.; Dizdaroğlu, Miral; Rodriguez, Henry; Akman, Steven A.
 CORPORATE SOURCE: Department of Chemistry, Wake Forest University, Winston-Salem, NC, 27109-7486, USA
 SOURCE: Journal of Inorganic Biochemistry (2003), 95(2-3), 87-96
 CODEN: JIBIDJ; ISSN: 0162-0134
 PUBLISHER: Elsevier Science Inc.
 DOCUMENT TYPE: Journal
 LANGUAGE: English

AB The ability of the fungal carcinogen, ochratoxin A (OTA, 1), to facilitate copper-promoted oxidative DNA damage has been assessed using supercoiled plasmid DNA (Form I)-agarose gel electrophoresis and gas chromatog.-mass spectrometry with selected-ion monitoring (GC-MS-SIM). OTA is shown to promote oxidative cleavage of Form I DNA with optimal cleavage efficiency occurring under excess Cu(II) conditions. As the concentration of OTA was increased and present in excess of Cu(II) the cleavage was less effective. Parallel findings were found for the ability of the OTA-Cu mixture to facilitate oxidative base damage. Yields (lesions per 106 DNA bases) of modified bases upon exposure of calf-thymus DNA (CT-DNA) to OTA-H2O2-Cu(II) were diminished when the OTA:Cu ratio was increased to 5:1. Electrochem. studies carried out in methanol implicate a ligand-centered 2e oxidation of OTA in the presence of excess Cu(II), while product analyses utilizing electrospray mass spectrometry support the intermediacy of the quinone, OTQ (3), in Cu-promoted oxidation of OTA. The implications of these findings with regard to the mutagenicity of OTA are discussed.


IT 560134-09-0P

RL: BSU (Biological study, unclassified); FMU (Formation, unclassified); PRP (Properties); RCT (Reactant); SPN (Synthetic preparation); BIOL (Biological study); FORM (Formation, nonpreparative); PREP (Preparation); RACT (Reactant or reagent)

(stoichiometric preference in copper-promoted oxidative DNA damage by ochratoxin A)
 RN 560134-09-0 CAPLUS
 CN L-Phenylalanine, N-[[[(3R)-6-[[[(2R)-2-carboxyethylthio]-3,4-dihydro-5,8-dihydroxy-3-methyl-1-oxo-1H-2-benzopyran-7-yl]carbonyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

L4 ANSWER 36 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)



REFERENCE COUNT: 48 THERE ARE 48 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 37 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
 ACCESSION NUMBER: 2003:341041 CAPLUS
 DOCUMENT NUMBER: 139:363489
 TITLE: Structure-biological response relationship of fMLP analogs in human neutrophils
 AUTHOR(S): Spisani, S.; Turchetti, M.; Cavicchioni, G.
 CORPORATE SOURCE: Dipartimento di Biochimica e Biologia Molecolare, Università degli Studi di Ferrara, Ferrara, 44100, Italy
 SOURCE: Chemotaxis and Migration (2002), 3(4), 68-83
 CODEN: CMHICK; ISSN: 1608-2265
 PUBLISHER: Vicer Publishing
 DOCUMENT TYPE: Journal
 LANGUAGE: English


AB Neutrophils constitute the first line of defense against bacterial invasion. They migrate to infected tissues along a concentration gradient of chemoattractant mols., the most important of which is for-Met-Leu-Phe-OH. Different responses arise from binding of formylpeptide with different isoforms of the specific receptor. The goal of studies reported herein was to clarify: (i) the role of for-Met-Leu-Phe amide bonds in receptor-ligand crosslinking, (ii) the features peculiar to the Met, Leu, and Phe receptor pockets. The data show: (1) the amide bond at position 2 must be protic, while the amide bond at position 3 participates and links the receptor, but its role is not mandatory. Furthermore, the isoform that elicits chemotaxis is structurally more exacting than the isoform that elicits superoxide anion production; (2) the Met receptor pocket shows a pos. charged area narrow in dimension, located at a well defined distance from the backbone, and oriented in a specific position, not completely surrounding the internally located side chain; (3) the Leu receptor pocket highlights that the hydrophobicity of the second residue is the mandatory key to stimulate a good chemotaxis, while hydrophilicity associated to good steric hindrance elicits a potent superoxide anion production; (4) the features peculiar to the Phe receptor pocket, unexpectedly, prove to be less mandatory in the isoform that elicits chemotaxis, which is strongly stimulated, than in the isoform that triggers superoxide anion production, which instead is less triggered.

IT 177656-62-1

RL: BSU (Biological study, unclassified); PRP (Properties); BIOL (Biological study)
 (structure-biol. response relationship of fMLP analogs in human neutrophils)

RN 177656-62-1 CAPLUS
 CN L-Phenylalanine, N-[[[(3S)-2-[(2S)-2-(formylamino)-4-(methylthio)-1-oxobutyl]-1,2,3,4-tetrahydro-3-isoquinolinyl]carbonyl]-, methyl ester (9CI) (CA INDEX NAME)

Absolute stereochemistry.




L4 ANSWER 37 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)

REFERENCE COUNT: 42 THERE ARE 42 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 38 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
 ACCESSION NUMBER: 2003:319899 CAPLUS
 DOCUMENT NUMBER: 138:338490
 TITLE: Preparation of β -carboline derivatives as protein tyrosine phosphatase (PTP)-inhibitors
 INVENTOR(S): Mjalli, Adnan M. M.; Andrews, Robert C.; Xie, Rongyuan; Yarragunta, Ravindra R.; Ren, Tan
 PATENT ASSIGNEE(S): Transtech Pharma, Inc., USA
 SOURCE: PCT Int. Appl., 110 pp.
 CODEN: PIXXD2
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2003033496	A1	20030424	WO 2002-US33520	20021018
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM			
RW:	GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG			
US 2004014778	A1	20040122	US 2002-274546	20021018
PRIORITY APPLN. INFO.:			US 2001-346125P	P 20011019
			US 2001-346176P	P 20011019
OTHER SOURCE(S):		MARPAT 138:338490		
GI				



AB The invention provides compds. I [RCH:CHR is (un)substituted (hetero)aryl; X is O, S, imino; Y is CH2, CH2CH2; R1 is alk(en)(yn)yl, (hetero)aryl, heterocyclyl, cycloalkyl, (hetero)aryl, etc.; R2 is H, alk(en)(yn)yl, (hetero)aryl, heterocyclyl, cycloalkyl, arylalk(en)(yn)yl, carboxy, etc.; R3 is H, alk(en)(yn)yl, (hetero)arylalk(en)(yn)yl; R4 is OH, (cyclo)alkoxy, (un)substituted amino, etc.] which are useful as inhibitors of protein tyrosine phosphatases (PTases). Thus, N-benzyl-1-(1,1'-biphenyl-4-yl)-1,2,3,4-tetrahydro- β -carboline-3-carboxamide was prepared from DL-tryptophan Me ester, 4-biphenylcarboxaldehyde, and

L4 ANSWER 38 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)
 benzylamine.
 IT 515157-61-6P
 RL: PAC (Pharmacological activity); RCT (Reactant); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); RACT (Reactant or reagent); USES (Uses)
 (preparation of β -carboline derivs. as protein tyrosine phosphatase (PTP)-inhibitors)
 RN 515157-61-6 CAPLUS
 CN [1,1'-Biphenyl]-4-acetic acid, α -[[[(1S,3R)-1-cyclohexyl-2,3,4,9-tetrahydro-1H-pyrido[3,4-b]indol-3-yl]carbonyl]amino]- (9CI) (CA INDEX NAME)



Absolute stereochemistry.



REFERENCE COUNT: 4 THERE ARE 4 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT


L4 ANSWER 39 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
 ACCESSION NUMBER: 2003:301053 CAPLUS
 DOCUMENT NUMBER: 138:321578
 TITLE: Preparation of peptides as ligands of melanocortin receptors
 INVENTOR(S): Dyck, Brian P.; Goodfellow, Val; Phillips, Teresa; Parker, Jessica; Zhang, Xiaohu; Chen, Chen; Tran, Joe Anh; Pontillo, Joseph; Tucci, Fabio C.
 PATENT ASSIGNEE(S): Neurocrine Biosciences, Inc., USA
 SOURCE: PCT Int. Appl., 112 pp.
 CODEN: PIXXD2
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2003031410	A1	20030417	WO 2002-US32282	20021009
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM			
RW:	GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG			
US 2003158209	A1	20030821	US 2002-268923	20021009
PRIORITY APPLN. INFO.:			US 2001-328295P	P 20011009
			US 2002-366745P	P 20020322
OTHER SOURCE(S):		MARPAT 138:321578		
GI				



L4 ANSWER 39 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)
 AB The invention relates to peptides I [m = 1-4; n = 0-4; A is (un)substituted alkanediyl; R1, R2, R3a, R3b = H, (un)substituted alkyl, aryl, arylalkyl, heterocyclyl, or heterocyclylalkyl or may combine to form rings; R1 or R2 may also be acyl; R4 = (un)substituted (hetero)aryl; R5 = H, OH, (un)substituted alkyl, aryl, or heterocyclyl; R6 = cyano, nitro, (un)substituted heterocyclyl, amino, carbamoyl, etc.; R7 = H or 1-4 substituents], or stereoisomers, prodrugs or pharmaceutically-acceptable salts, which function as melanocortin receptor ligands and may be used to treat disorders or illnesses including cachexia, obesity, diabetes, inflammation, and sexual dysfunction. Thus, treatment of cyclohexanone with sodium metabisulfite in H2O, followed by addition of Boc-protected piperazine and then NaCN, afforded 1-Boc-4-(1-cyanocyclohexyl)piperazine. The latter was converted into peptide II via coupling reaction.
 IT 252008-71-2
 RL: RCT (Reactant); RACT (Reactant or reagent)
 (preparation of peptides as ligands of melanocortin receptors)
 RN 252008-71-2 CAPLUS
 CN 2-(1H)-Isoquinolinecarboxylic acid, 3-[[[(1R)-1-carboxy-2-(4-chlorophenyl)ethyl]amino]carbonyl]-3,4-dihydro-, 2-(1,1-dimethylethyl) ester, (3R)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



REFERENCE COUNT: 7 THERE ARE 7 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 40 OF 261 CAPLUS COPYRIGHT 2004 ACS ON STN

ACCESSION NUMBER: 2003:300170 CAPLUS
 DOCUMENT NUMBER: 139:143349
 TITLE: Pharmacological Profiles of Peptide Drug Candidates for the Treatment of Alzheimer's Disease
 AUTHOR(S): Adeesi, Celine; Prossard, Marie-Jose; Boissard, Christophe; Fraga, Santiago; Bieler, Sylvain; Ruckle, Thomas; Vilbois, Francis; Robinson, Sandra M.; Mutter, Manfred; Banks, William A.; Soto, Claudio
 CORPORATE SOURCE: Serono Pharmaceutical Research Institute, Geneva, 1228, Switz.
 SOURCE: Journal of Biological Chemistry (2003), 278(16), 13905-13911
 CODEN: JBCHA3; ISSN: 0021-9258
 PUBLISHER: American Society for Biochemistry and Molecular Biology
 DOCUMENT TYPE: Journal
 LANGUAGE: English

AB Amyloid plaques in brain, composed of aggregates of amyloid- β peptide, play a central role in the pathogenesis of Alzheimer's disease and represent a good target for treatment. We have shown previously that a 5-amino acid β -sheet breaker peptide (iA β Sp), end-protected, has the ability to induce a dramatic reduction in amyloid deposition in two different transgenic Alzheimer's models. The aim of this study was to evaluate the effect of chemical modifications of the peptide bonds at the metabolite cleavage sites on the pharmacol. properties of iA β Sp deriva. Using a rational approach, peptide analogs were designed and tested for in vitro activity and enzymic stability. One peptide analog containing a Me group introduced at the nitrogen atom of one amide bond showed increased stability in vitro, a 10-fold higher in vivo half-life, and good brain uptake compared with iA β Sp while maintaining a similar activity in vitro. Our results suggest that the pharmacol. profile of β -sheet breaker peptides can be improved to produce compds. with drug-like properties that might offer a new promise in the treatment of Alzheimer's disease.

IT 572914-18-2


RL: PAC (Pharmacological activity); PKT (Pharmacokinetics); THU (Therapeutic use); BIOL (Biological study); USES (Uses)
 (pharmacol. profiles of peptide drug candidates for treatment of Alzheimer's disease)

RN 572914-18-2 CAPLUS

CN L-Aspartic acid, L-leucyl-(3S)-1,2,3,4-tetrahydro-3-isoquinolinecarbonyl-L-phenylalanyl-L-phenylalanyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

L4 ANSWER 40 OF 261 CAPLUS COPYRIGHT 2004 ACS ON STN (Continued)



REFERENCE COUNT: 38

THERE ARE 38 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 41 OF 261 CAPLUS COPYRIGHT 2004 ACS ON STN

ACCESSION NUMBER: 2003:178421 CAPLUS
 DOCUMENT NUMBER: 138:349900
 TITLE: Ochratoxin A Forms a Carbon-Bonded C8-Deoxyguanosine Nucleoside Adduct: Implications for C8 Reactivity by a Phenolic Radical
 AUTHOR(S): Dai, Jian; Wright, Marcus W.; Manderville, Richard A.
 CORPORATE SOURCE: Department of Chemistry, Wake Forest University, Winston-Salem, NC, 27109, USA
 SOURCE: Journal of the American Chemical Society (2003), 125(13), 3716-3717
 CODEN: JACSAT; ISSN: 0002-7863
 PUBLISHER: American Chemical Society
 DOCUMENT TYPE: Journal
 LANGUAGE: English

AB The ability of the carcinogenic fungal toxin Ochratoxin A (OTA) to react with deoxyguanosine (dG) has been assessed using electrospray mass spectrometry and NMR. Photoexcitation of OTA (100 μ M) in the presence of 50 mol equiv of dG led to the isolation and identification of the C8-deoxyguanosine nucleoside adduct. Importantly, the same adduct was formed upon oxidative activation of OTA using horseradish peroxidase (HRP)/H₂O₂ or the transition metals Fe(II) and Cu(II), as evidenced by mass spectrometry. Because the mutagenicity and subsequent carcinogenicity of OTA are believed to stem from oxidative DNA damage (strand scission and oxidative base products) and formation of guanine-specific DNA adducts, the adduct confirms the ability of OTA to react covalently with dG and has important implications for the mechanism of action of OTA and other chlorophenolic toxins that undergo oxidation to yield phenoxyl radicals. The C8 position of dG is susceptible to radical attack, as was amply proven through formation of the hydroxyl radical-derived DNA lesion, 8-oxodeoxyguanosine. The adduct is the first structurally characterized nucleoside adduct of a chlorophenolic toxin, and its formation has important implications for the mutagenicity of phenolic xenobiotics.

IT 519005-04-0


RL: BSU (Biological study, unclassified); FMU (Formation, unclassified); PRP (Properties); BIOL (Biological study); FORM (Formation, nonpreparative)
 (ochratoxin A forms carbon-bonded C8-deoxyguanosine nucleoside adduct with implications for C8 reactivity by phenolic radical as mechanism for oxidative DNA damage)

RN 519005-04-0 CAPLUS

CN L-Phenylalanine, N-[[5-[2-amino-9-(2-deoxy- β -D-erythro-pentofuranosyl)-6,9-dihydro-6-oxo-1H-purin-8-yl]-3,4-dihydro-8-hydroxy-1-oxo-1H-2-benzopyran-7-yl]carbonyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

L4 ANSWER 41 OF 261 CAPLUS COPYRIGHT 2004 ACS ON STN (Continued)



REFERENCE COUNT: 28

THERE ARE 28 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 42 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 2003:133239 CAPLUS

DOCUMENT NUMBER: 138:170086

TITLE: Preparation of spiro[isquinoline-piperidine], spiro[indoline-piperidine], and spirocyclohexane compounds as antagonists of neuropeptide Y receptor

INVENTOR(S): Fukami, Takehiro; Nonoshita, Katsumasa; Sagara, Takeshi; Kishino, Hiroyuki

PATENT ASSIGNEE(S): Banyu Pharmaceutical Co., Ltd., Japan

SOURCE: PCT Int. Appl., 220 pp.

CODEN: PIXX2


DOCUMENT TYPE: Patent

LANGUAGE: Japanese

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2003014083	A1	20030220	WO 2002-37922	20020802
<p>W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM</p> <p>RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZH, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG</p>				
PRIORITY APPL. INFO:		JP 2001-239567 A 20010807		
OTHER SOURCE(S):		MARPAT 138:170086		
GI				



AB The invention relates to compds. such as spiro[cyclohexane-1,1'-(3'H)-isobenzofuran], spiro[4-, 5-, 6-, or 7-azaisobenzofuran-1(3H),1'-cyclohexane], spiro[indoline-3,1'-cyclohexane], spiro[indoline-3,4'-piperidine], spiro[isobenzofuran-1(3H),4'-piperidine], and spiro[isquinoline-1(2H),4'-piperidine] represented by the general formula (I) or salts or esters thereof [A = linear C1-6 hydrocarbon group which

L4 ANSWER 42 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)

may be substituted or interrupted by oxygen or nitrogen; Ar1 = (un)substituted aryl or heteroaryl; n = 0,1; R = H, lower alkylene; T, U, V, W = (un)substituted CH or N and at least 2 of T, U, V, and W is (un)substituted CH; X = -N(SO2R1)-, -N(COR2)-, or CO; Y = -C(R3)(R4)-, O, or -N(R5)-; and Z = CH or nitrogen; wherein R1, R2, R5 = H, lower alkyl, aralkyl, aryl; R3, R4 = H, HO, lower alkyl, aralkyl, aryl. These compds. exhibit neuropeptide Y (NPY) receptor antagonism and are therefore useful as treating agents for various diseases in which NPY participates such as circulatory diseases, central nervous system diseases, and metabolic diseases, in particular over eating (hyperphagia), obesity, and diabetes. Thus, 64 mg 4-phenylcyclohexylamine hydrochloride and 115 mg 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride were added to a soln. of 74 mg trans-3'-oxospiro[cyclohexane-1,1'-(3'H)-isobenzofuran]-4-carboxylic acid in 2 mL pyridine and stirred at room temp. for 24 h to give trans-3'-oxo-N-(trans-4-phenylcyclohexyl)spiro[cyclohexane-1,1'-(3'H)-isobenzofuran]-4-carboxamide (II). I and trans-N-[(S)-1-benzyl-2-(benzylamino)ethyl]-1-(methanesulfonyl)spiro[indoline-3,1'-cyclohexane]-4-carboxamide showed IC50 of 2.5 and 0.69 nM for inhibiting the binding of [125I]peptide YY to human NPY Y5 receptor.

IT 497238-78-5P


RL: PAC (Pharmacological activity); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)

(preparation of spiro[isquinoline-piperidine], spiro[indoline-piperidine], and spiro[azaisobenzofuran-cyclohexane], and spirocyclohexane compds. as antagonists of neuropeptide Y receptor for treating overeating, obesity, and diabetes)

RN 497238-78-5 CAPLUS

CN Spiro[cyclohexane-1,3'-(3H)indole]-4-carboxamide, N-1-[(4-fluorophenyl)methyl]-2-oxo-2-[(4-pyridinylmethyl)amino]ethyl]-1',2'-dihydro-1'-(methylsulfonyl)-, trans- (9CI) (CA INDEX NAME)

Relative stereochemistry.



REFERENCE COUNT: 25 THERE ARE 25 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 43 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 2003:126123 CAPLUS

DOCUMENT NUMBER: 139:69500

TITLE: Organometallic β -turn mimetics. A structural and spectroscopic study of inter-strand hydrogen bonding in ferrocene and cobaltocenium conjugates of amino acids and dipeptides

AUTHOR(S): van Staveren, Dave R.; Weyhermueller, Thomas;

CORPORATE SOURCE: Metzler-Nolte, Nils

Muelheim/Ruhr, D-45470, Germany

Dalton Transactions (2003), (2), 210-220

CODEN: DTRAPF, ISSN: 1477-9226

Royal Society of Chemistry

PUBLISHER: Journal

DOCUMENT TYPE: English

LANGUAGE: English

OTHER SOURCE(S): CASREACT 139:69500

AB By using organometallic turn-mimetics, the influence of a pos. charge on the structure and stability of peptide turn structures which are stabilized by H bonds. Starting from metallocene mono- (1) or dicarboxylic acid (2), 11 amide derivs. were prepared: CpM(C5H4-CO-Phe-Ome) (3), CpM(C5H4-CO-Ala-Phe-Ome) (4), CpM(C5H4-CO-NH-CHMe-Ph) (5), M(C5H4-CO-Phe-Ome)2 (6), M(C5H4-CO-Ala-Phe-Ome)2 (7), and Fe(C5H4-CO-NH-CHMe-Ph)2 (8a) with Cp = η -C5H5 and M = Fe (ferrocene, a) or M = Co+ (cobaltocenium, b). All compds. were characterized by elemental anal., MS, IR, electrochem., Mossbauer spectroscopy (a only) and NMR spectroscopy. Solid state structures of 4a, 6a, 7a, 3b, and 5b were determined by single crystal x-ray diffraction. 1H NMR data (δ (NH) and δ S(NH) with T) as well as solution IR spectra were evaluated to determine intramol. H bond interactions in solution. No intramol. H bonds form


in the monosubstituted derivs. 3-5 and in 8a. For 7, a strong intramol. H bond is observed between the NHAla and COAla, of the other ring, forming an 11-membered ring in solution as well as in the solid state. The situation is most complex for 6, which forms an intramol. 8-membered ring by H bonds NHPh...COCP in the solid state (6a), but a sym. 11-membered ring structure with NHPh...COPh...bonds in solution. A comparison of the uncharged ferrocene derivs. with the iso-structural but pos. charged cobaltocenium derivs. reveals only minor differences. Apparently, the presence of a pos. charge does not significantly influence H bonds in peptide turn structures. The results are related to geometries and amino acid sequences in protein turn structures and a nomenclature for turn mimetics with a parallel orientation of the two peptide strands is proposed.

IT 181589-78-6P
 RL: CPS (Chemical process); PEP (Physical, engineering or chemical process); PRP (Properties); RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); PROC (Process); RACT (Reactant or reagent)
 (crystal structure, electrochem. redox, Mossbauer spectra; condensation reaction of ferrocene carboxylic acid derivs. with phenylethylamine, alanine ester and dipeptide in presence of coupling reagent to give amides)

RN 181589-78-6 CAPLUS


CN Ferrocene, 1,1'-bis[[[(1S)-2-methoxy-2-oxo-1-(phenylmethyl)ethyl]amino]carbonyl]- (9CI) (CA INDEX NAME)

L4 ANSWER 43 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)



REFERENCE COUNT: 43 THERE ARE 43 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 44 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
 ACCESSION NUMBER: 2003:89939 CAPLUS
 DOCUMENT NUMBER: 138:271967
 TITLE: A Solid-Phase Synthetic Strategy for Labeled Peptides:
 Synthesis of a Biotinylated Derivative of the δ
 Opioid Receptor Antagonist TIPP (Tyr-Tic-Phe-Phe-OH)
 Kumar, Vivek; Aldrich, Jane V.
 CORPORATE SOURCE: Department of Pharmaceutical Sciences School of
 Pharmacy, University of Maryland, Baltimore, MD,
 21201, USA
 SOURCE: Organic Letters (2003), 5(5), 613-616
 CODEN: ORLEF7; ISSN: 1523-7060
 PUBLISHER: American Chemical Society
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 OTHER SOURCE(S): CASREACT 138.271967
 GI




AB A general solid-phase synthetic strategy for labeled peptides was developed and used to prepare a biotinylated peptide I (Tic = 1,2,3,4-tetrahydroisoquinolinyl-3-carbonyl) analog of the δ opioid receptor antagonist TIPP (H-Tyr-Tic-Phe-Phe-OH). A monoprotected hydrophilic diamine linker was attached to an aldehyde-containing solid-phase resin by reductive amination, followed by introduction of biotin and peptide synthesis to yield biotinyl-peptide I. The high δ receptor affinity and selectivity of I demonstrate the applicability of this design approach for labeled peptide derivs.

IT 319906-09-7
 RL: BSU (Biological study, unclassified); BIOL (Biological study) (biol. activity comparisons at δ - and μ -opioid receptors)
 RN 319906-09-7 CAPLUS
 CN L- α -Asparagine, L-tyrosyl-(3S)-1,2,3,4-tetrahydro-3-isoquinolinecarboxyl-L-phenylalanyl-L-phenylalanyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

L4 ANSWER 44 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)



REFERENCE COUNT: 21 THERE ARE 21 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 45 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
 ACCESSION NUMBER: 2003:76750 CAPLUS
 DOCUMENT NUMBER: 138:137595
 TITLE: Preparation of N-[(aryl or thienyl)sulfonyl]dipeptide derivatives and analogs as α 2 β 1 integrin inhibitors
 INVENTOR(S): Takayanagi, Masaru; Fukuchi, Naoyuki; Sugiki, Masayuki; Futaki, Fumie; Takehana, Shunji; Kajigaya, Yuki; Takamatsu, Yayoi; Tokumaeu, Munetaka; Yoshida, Kaoru
 PATENT ASSIGNEE(S): Ajinomoto Co., Inc., Japan
 SOURCE: PCT Int. Appl., 187 pp.
 CODEN: PIXXD2
 DOCUMENT TYPE: Patent
 LANGUAGE: Japanese
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2003008380	A1	20030130	WO 2002-JP7250	20020717
W: AE, AC, AL, AM, AT, AU, AZ, BA, BB, BS, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM				
RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				

PRIORITY APPLN. INFO.: JP 2001-218507 A 20010718
 JP 2001-351077 A 20011116

OTHER SOURCE(S): MARPAT 138:137595


AB The compds. represented by the following formula D-Y-N(L)-CH(B)-X-CH₂(CH₂n)A)CO₂H (wherein n = an integer of 0-2; X = CH₂O, CO₂, each N-(un)substituted CH₂NH, CONH, C(S)NH, or SO₂NH, CH₂S, CH₂SO₂, Y = CH₂, CO, C(S), SO₂, each N-(un)substituted NHCO or NHC(S); A, D = (un)substituted Ph, aryl, heteroaryl; B = (un)substituted amino-lower alkyl or carbamoyl-lower alkyl; L = H, lower alkyl, lower alkenyl, lower alkynyl, cycloalkyl or cycloalkyl-lower alkyl optionally containing a heteroatom in the ring, aryl, heteroaryl, aryl-lower alkyl, heteroaryl-lower alkyl, halo-lower alkyl, hydroxy-lower alkyl, lower alkoxy-lower alkyl, mercapto-lower alkyl, lower alkylthio-lower alkyl, (un)substituted amino-lower alkyl, carboxy-lower alkyl, lower alkoxy-carbonyl-lower alkyl, (un)substituted carbamoyl-lower alkyl; E = HO, lower alkoxy, (un)substituted amino, hydroxylamino or pharmacol. acceptable salts thereof are prepared. These compds. have α 2 β 1 integrin inhibitory activity (inhibition of binding between platelet GPIIb/IIIa receptor and collagen) and are useful as antiplatelet agents for the prevention or treatment of various diseases such as myocardial infarction, angina pectoris, acute coronary syndrome, peripheral artery obstruction, deep venous thrombosis, cerebral infarction, stroke, and pulmonary embolism. Thus, N-(4-nitrobenzenesulfonyl)-L-prolyl-L-4-aminophenylalanine (prepared by the Fmoc solid phase method) showed IC₅₀ of 2.0 μ g/mL for inhibiting the binding of collagen to platelet GPIIb/IIIa receptor.

IT 493040-65-69
 RL: PAC (Pharmacological activity); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES

L4 ANSWER 45 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)

(Uses)
 (prepn. of N-[(aryl or thienyl)sulfonyl]dipeptide derivs. and analogs as α 2 β 1 integrin inhibitors and antiplatelet agents for the prevention or treatment of various diseases)
 RN 493040-65-6 CAPLUS
 CN D-Phenylalanine, N-[(3S)-1,2,3,4-tetrahydro-2-(phenylsulfonyl)-3-isoquinolinyl]carbonyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



REFERENCE COUNT: 29 THERE ARE 29 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 46 OF 261 CAPLUS COPYRIGHT 2004 ACS ON STM
 ACCESSION NUMBER: 2003:24837 CAPLUS
 DOCUMENT NUMBER: 138:221561
 TITLE: Parallel Solid-Phase Synthesis of 2-Imino-4-oxo-1,3,5-triazino[1,2-a]benzimidazoles via Tandem Aza-Wittig/Heterocumulene-Mediated Annulation Reaction
 AUTHOR(S): Hoesl, Cornelia E.; Nefzi, Adel; Houghten, Richard A.
 CORPORATE SOURCE: Torrey Pines Institute for Molecular Studies, San Diego, CA, 92121, USA
 SOURCE: Journal of Combinatorial Chemistry (2003), 5(2), 155-160
 CODEN: JCCHFF; ISSN: 1520-4766
 PUBLISHER: American Chemical Society
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 OTHER SOURCE(S): CASREACT 138:221561

AB The parallel synthesis of a large number of 2-imino-4-oxo-1,3,5-triazino[1,2-a]benzimidazole deriva. via a solid-phase 1,3,5-triazine annulation reaction is described. The solid-phase approach involves the in situ generation of iminophosphorane deriva. derived from resin-bound 2-aminobenzimidazoles employing Mitsunobu conditions. The subsequent Aza-Wittig reaction of the iminophosphoranes with isocyanates leads to highly reactive carbodiimides, which undergo an intramol. heterocyclization reaction to form tetrasubstituted 2-imino-4-oxo-1,3,5-triazino[1,2-a]benzimidazoles in high yields (74-94%) and good purity (>80%).

IT 500732-13-8P

RL: SPN (Synthetic preparation); PREP (Preparation)


(parallel solid-phase synthesis of 2-imino-4-oxo-1,3,5-triazino[1,2-a]benzimidazoles via tandem aza-Wittig/heterocumulene-mediated annulation reaction)

RN 500732-13-8 CAPLUS

CN 1,3,5-Triazino[1,2-a]benzimidazole-7-carboxamide, N-[(1S)-2-amino-2-oxo-1-phenylethyl]-10-butyl-3-ethyl-2-(ethylimino)-2,3,4,10-tetrahydro-4-oxo- (9CI) (CA INDEX NAME)

Absolute stereochemistry.


Double bond geometry unknown.



REFERENCE COUNT: 33 THERE ARE 33 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 47 OF 261 CAPLUS COPYRIGHT 2004 ACS ON STM
 ACCESSION NUMBER: 2003:23531 CAPLUS
 DOCUMENT NUMBER: 138:90079
 TITLE: Preparation of N-arylsulfonyl aza-bicyclic derivatives as potent cell adhesion inhibitors
 INVENTOR(S): Lin, Linus S.; Doherty, George; Shah, Shrenik K.; Chang, Linda L.; Hagmann, William K.; Mumford, Richard A.
 PATENT ASSIGNEE(S): USA
 SOURCE: U.S. Pat. Appl. Publ., 31 pp.
 CODEN: USXXCO
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 2003008861	A1	20030109	US 2002-96607	20020313
PRIORITY APPLN. INFO:			US 2001-277233P	P 20010320
OTHER SOURCE(S):			MARPAT 138:90079	



L4 ANSWER 49 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
 ACCESSION NUMBER: 2003:146 CAPLUS
 DOCUMENT NUMBER: 138:205334
 TITLE: Novel Antibiotics: Macrocylic Peptides Designed to Trap Holliday Junctions
 AUTHOR(S): Bolla, Megan L.; Azevedo, Enrique V.; Smith, Jason M.; Taylor, Rachel E.; Ranjit, Dev K.; Segall, Anca M.; McAlpine, Shelli R.
 CORPORATE SOURCE: Department of Chemistry, Molecular Biology Institute and Center for Applied and Experimental Genomics, San Diego State University, San Diego, CA 92182-1030, USA
 SOURCE: Organic Letters (2003), 5(2), 109-112
 CODEN: ORLEF7; ISSN: 1523-7060
 PUBLISHER: American Chemical Society
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 OTHER SOURCE(S): CASREACT 138:205334

AB This work describes the synthesis of eight macrocyclic peptides designed to trap Holliday junctions in bacteria, thereby inhibiting bacterial growth. These macrocycles were designed from linear dimerized hexapeptides that bind to the C-2 sym. Holliday junction. They were synthesized from three monomers using a combinatorial-like strategy that permits elucidation of the monomer role in accumulation of Holliday junctions and antibiotic activity. These macrocycles are an important step in designing and synthesizing a new class of antibiotics.
 IT 617688-75-2P
 RL: CPN (Combinatorial preparation); CRT (Combinatorial reactant); RCT (Reactant); SPN (Synthetic preparation); CMBI (Combinatorial study); PREP (Preparation); RACT (Reactant or reagent)
 (combinatorial-style preparation and activity of macrocyclic antibacterial peptides designed to trap Holliday junctions in bacteria)
 RN 617688-75-2 CAPLUS
 CN L-Phenylalanine, N-[(1,1-dimethylethoxy)carbonyl]-L-phenylalanyl-(3S)-1,2,3,4-tetrahydro-3-isoquinolinecarbonyl-L-phenylalanyl-L-phenylalanyl-(3S)-1,2,3,4-tetrahydro-3-isoquinolinecarbonyl-, methyl ester (9CI) (CA INDEX NAME)

Absolute stereochemistry.




REFERENCE COUNT: 22 THERE ARE 22 CITED REFERENCES AVAILABLE FOR THIS

L4 ANSWER 49 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)
 RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 50 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
 ACCESSION NUMBER: 2002:965131 CAPLUS
 DOCUMENT NUMBER: 138:24961
 TITLE: Preparation of N-arylsulfonyl aryl aza-bicyclic derivatives as potent cell adhesion inhibitors
 INVENTOR(S): Lin, Linus S.; Shah, Shrenik K.; Chang, Linda L.; Hagmann, William K.; Mumford, Richard A.
 PATENT ASSIGNEE(S): USA
 SOURCE: U.S. Pat. Appl. Publ., 19 pp.
 CODEN: USXXCO
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 2002193399	A1	20021219	US 2002-97028	20020313
US 6559174	B2	20030506		

PRIORITY APPLM. INFO.: US 2001-277235P P 20010320
 OTHER SOURCE(S): MARPAT 138:24961
 GI



AB Comps. I [R2 is an (un)substituted (hetero)aryl ring; R1 = H, alkyl, arylalkyl; R2, R4 = halo, alkyl, alkoxy; R3 = H, OH, MeO, NH2; Z = N or N:O; Ar1 = (un)substituted Ph, pyridyl, pyridazinyl, pyrimidinyl, pyrazinyl, or triazinyl; Ar2 = 1,4-phenylene or 2,5-pyridylene; X, Y = (CH2)0-2] or their pharmaceutically-acceptable salts were prepared as antagonists of VLA-4 and/or α4β7 and as such are useful in the inhibition or prevention of cell adhesion and cell-adhesion mediated pathologies. Thus, N-[N-(4-methylbenzenesulfonyl)-1,3-dihydro-2H-isoindole-1-carbonyl]-4-[(3',5'-dichloroisocotinyloxy)amino]-L-phenylalanine was prepared by coupling of N-(4-methylbenzenesulfonyl)-1,3-dihydro-2H-isoindole-1-carboxylic acid with 4-[(3',5'-dichloroisocotinyloxy)amino]-L-phenylalanine tert-Bu ester (syntheses given), followed by ester cleavage using TFA.
 IT 478170-92-2P
 RL: PAC (Pharmacological activity); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)
 (preparation of N-arylsulfonyl heteroaryl amino acid derivs. as cell adhesion inhibitors)
 RN 478170-92-2 CAPLUS
 CN L-Phenylalanine, N-[[2-[(3,5-dichlorophenyl)sulfonyl]-1,2,3,4-tetrahydro-3-isoquinolinyl]carbonyl]-4-[[[(3,5-dichloro-4-pyridinyl)carbonyl]amino]-(9CI) (CA INDEX NAME)

L4 ANSWER 50 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)
 Absolute stereochemistry.




L4 ANSWER 51 OF 261 CAPLUS COPYRIGHT 2004 ACS ON STN
 ACCESSION NUMBER: 2002:943502 CAPLUS
 DOCUMENT NUMBER: 138:385709
 TITLE: Identification of TNF- α inhibitors from a split-pool library based on a tyrosine-proline peptidomimetic scaffold
 AUTHOR(S): Jackson, Randy W.; Tabone, John C.; Howbert, J. Jeffrey
 CORPORATE SOURCE: Department of Chemistry, Celltech R&D, Inc., Bothell, WA, 98021, USA
 SOURCE: Bioorganic & Medicinal Chemistry Letters (2003), 13(2), 205-208
 CODEN: BMCLE8; ISSN: 0960-894X
 PUBLISHER: Elsevier Science Ltd.
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 OTHER SOURCE(S): CASREACT 138:385709

AB The design and synthesis of a combinatorial library based on a 4-aryloxyproline scaffold with tyrosine as the aryl portion is described. The 1728 member library was prepared using the split-pool method to generate pools of compds. Screening of the library components as mixts. followed by deconvolution led to the discovery of novel inhibitors of TNF- α induced apoptosis.

IT 526223-58-5P
 RL: BSU (Biological study, unclassified); CPN (Combinatorial preparation); BIOL (Biological study); CMBI (Combinatorial study); PREP (Preparation) (preparation of peptide library based on tyrosine-proline peptidomimetic scaffold and identification of TNF inhibitors)

RN 526223-58-5 CAPLUS
 CN 3-Quinolincarboxamide, N-[(1S)-2-amino-1-[[4-[[[(3S,5S)-5-[(dipentylamino)carbonyl]-1-(3-quinolinylcarbonyl)-3-pyrrolidinyl]oxy]phenyl]methyl]-2-oxoethyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



REFERENCE COUNT: 22 THERE ARE 22 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 52 OF 261 CAPLUS COPYRIGHT 2004 ACS ON STN
 ACCESSION NUMBER: 2002:928230 CAPLUS
 DOCUMENT NUMBER: 138:19472
 TITLE: Method of identifying inhibitors of Cdc25 using three dimensional crystal structure of the catalytic domain of Cdc25
 INVENTOR(S): Taylor, Neil R.; Borhani, David; Epstein, David; Rudolph, Johannes; Ritter, Kurt; Fujimori, Taro; Robinson, Simon; Eckstein, Jens; Haupt, Andreas; Walker, Nigel; Dixon, Richard W.; Choquette, Deborah; Blanchard, Jill; Kluge, Arthur; Pal, Kolloi; Bockovich, Nicholas; Come, Jon; Hediger, Mark
 PATENT ASSIGNEE(S): Australia
 SOURCE: U.S. Pat. Appl., 246 pp., Cont.-in-part of U.S. Ser. No. 645,750.
 CODEN: USXXCO
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 2
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 2002183249	A1	20021205	US 2001-797500	20010301
PRIORITY APPLN. INFO.:			US 1999-172215P	P 19990831
			US 2000-645750	A2 20000824


OTHER SOURCE(S): MARPAT 138:19472
 AB The present invention relates to the x-ray crystallog. study of proteins comprising the catalytic domains of Cdc25. The atomic coordinates which result from this study are of use in identifying compds. which fit in the catalytic domain and are, therefore, potential inhibitors of Cdc25. The present invention further provides proteins which comprise the ligand binding domain of Cdc25, crystalline forms of these proteins and the use of these crystalline forms to determine the three dimensional structure of the catalytic domain of Cdc25. The invention also relates to the use of the three dimensional structure of the Cdc25 catalytic domain in methods of designing and/or identifying potential inhibitors of Cdc25 activity, for example, compds. which inhibit the binding of a native substrate to the Cdc25 catalytic domain. These Cdc25 inhibitors are of use in methods of treating a patient having a condition which is modulated by Cdc25 activity, for example, a condition characterized by excessive, inappropriate or undesirable cellular proliferation such as cancer.

IT 477908-84-2P
 RL: BSU (Biological study, unclassified); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USRS (Uses)
 (method of identifying inhibitors of Cdc25 using three dimensional crystal structure of catalytic domain of Cdc25)
 RN 477908-84-2 CAPLUS
 CN L-Norvalinamide, N-(4-dibenzofuranylcarbonyl)-4-(sulfomethyl)-L-phenylalanyl-L-norvalyl-2-methyl-L-prolyl-3-benzo[b]thien-3-yl-L-alanyl-5-carboxy-N-(1,1-dimethylethyl)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.

L4 ANSWER 52 OF 261 CAPLUS COPYRIGHT 2004 ACS ON STN (Continued)

PAGE 1-A



PAGE 1-B




L4 ANSWER 53 OF 261 CAPLUS COPYRIGHT 2004 ACS ON STN
 ACCESSION NUMBER: 2002:850472 CAPLUS
 DOCUMENT NUMBER: 138:68223
 TITLE: Detection and Characterization of a Glutathione Conjugate of Ochratoxin A
 AUTHOR(S): Dai, Jian; Park, Gyungae; Wright, Marcus W.; Adams, Marissa; Akman, Steven A.; Manderville, Richard A.
 CORPORATE SOURCE: Department of Chemistry, Wake Forest University, Winston-Salem, NC, 27109-7486, USA
 SOURCE: Chemical Research in Toxicology (2002), 15(12), 1581-1588
 CODEN: CRTOC; ISSN: 0893-228X
 PUBLISHER: American Chemical Society
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 AB The ability of the carcinogenic mycotoxin ochratoxin A (OTA) to react with reduced glutathione (GSH) has been assessed using electrospray ionization (ESI)-MS techniques. On the basis of the assumption that OTA undergoes biotransformation into the reactive quinone species, ochratoxin quinone (OTQ), a synthetic sample of the reduced form of OTQ, ochratoxin hydroquinone (OTHQ), was prepared and photoreacted with 6 M equiv of GSH to yield an authentic sample of the conjugate that was definitively identified by mass spectrometry, UV-vis spectroscopy and NMR. With the authentic sample of the conjugate in hand, it was demonstrated that the same conjugate is produced from reaction of 100 μ M OTA (1) in the presence of 5 mM GSH following incubation for 1 h with either horseradish peroxidase (HRP)/H₂O₂, rat liver microsomes (RLM)/NADPH or free Fe(II). In each of these oxidative systems the conjugate was generated in less than 1% yield and the parent OTA mol. is poorly metabolized. Comparison of the peak area ratio of the conjugate to that for the hydroxyOTA metabolite from the RLM/NADPH system implied that the conjugate was produced at a rate of approx. 1-3 pmol min⁻¹ (mg of protein)⁻¹. These studies are the first to demonstrate that OTA undergoes biotransformation to a reactive intermediate [OTQ] that covalently reacts with GSH to yield the conjugate. The biol. implications of the reactivity of OTA toward GSH are discussed.

IT 481053-26-3
 RL: ANT (Analyte); BSU (Biological study, unclassified); FMU (Formation, unclassified); PRP (Properties); ANST (Analytical study); BIOL (Biological study); FORM (Formation, nonpreparative)
 (detection, characterization and biol. implications of glutathione conjugate of ochratoxin A as biotransformation product)
 RN 481053-26-3 CAPLUS
 CN Glycine, L- γ -glutamyl-S-[(3R)-7-[[[(1S)-1-carboxy-2-phenylethyl]amino]carbonyl]-3,4-dihydro-5,8-dihydroxy-3-methyl-1-oxo-1H-2-benzopyran-6-yl]-L-cysteinyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

L4 ANSWER 53 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)




REFERENCE COUNT: 48 THERE ARE 48 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 54 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 2002:832787 CAPLUS
 DOCUMENT NUMBER: 137:337786
 TITLE: Preparation of chiral alkylaminochroman derivatives as β_3 -adrenoreceptor agonists
 INVENTOR(S): O'Connor, Stephen J.; Ladouceur, Gaetan H.; Bullock, William H.; Campbell, Ann-Marie; Dai, Miao; Dally, Robert; Dumas, Jacques; Hatoum-Mokdad, Holia N.; Khire, Uday; Lee, Wendy; Liu, Qingjie; Lowe, Derek B.; Magnuson, Steven R.; Qi, Ming; Shelekhn, Tatiana E.; Shen, Quanrong; Smith, Roger A.; Wang, Ming
 PATENT ASSIGNEE(S): Bayer Corporation, USA
 SOURCE: PCT Int. Appl., 193 pp.
 CODEN: PIXXD2
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2002085891	A1	20021031	WO 2002-US12940	20020422
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, ME, MK, MN, MW, MX, MY, NZ, NO, NZ, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM			
RW:	GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG			
US 2003078260	A1	20030424	US 2002-131448	20020422
US 6660752	B2	20031209		
EP 1389202	A1	20040218	EP 2002-723958	20020422
R:	AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR			
US 2004072828	A1	20040415	US 2003-666903	20030917
PRIORITY APPLN. INFO.:			US 2001-285719P	P 20010423
			US 2001-324518P	P 20010926
			US 2002-131448	A1 20020422
			WO 2002-US12940	W 20020422
OTHER SOURCE(S):		MARPAT 137:337786		
GI				

L4 ANSWER 54 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)



II

L4 ANSWER 54 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)



REFERENCE COUNT: 2 THERE ARE 2 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

AB This invention relates to novel 2,6-substituted chroman deriva. which are useful in the treatment of β_3 -adrenoreceptor mediated conditions. Title compds. I [wherein R = independently OH, -O, halo, CN, NO₂, (halo)alkyl, CF₃, NR1R1, SR1, OR1, SO₂R2, OCOR2, NR1COR2, COR2, NR1SO₂R2, or (un)substituted Ph or heterocyclyl; R1 = independently H, (CH₂)_mOC(CH₂)_mR5, or (un)substituted (cyclo)alkyl, Ph, or naphthyl; or NR1R1 = heterocyclyl; R2 = independently R1, OR1, NR1R1, or (un)substituted NHSO₂-2-Ph, NHSO₂-2-naphthyl, NHSO₂-2-alkyl, or heterocyclyl; R3 = H, alkyl, or COR3, R4 = H, alkyl(phenyl), or alkylpyridyl; R5 = H or CO₂H; R6 = H or (un)substituted alkyl or alkyl-SO₂-2-alkyl; Ar = Ph or (fused) hetero(aryl); Y = halo, NO₂, R6, SR1, SO₂-2C6H4CO₂R1, (CONR4CR4R4)PCO₂R1, or (un)substituted Ph or heterocyclyl; m = 1-3; n = 0-5; p = 1 or 2; and pharmaceutically acceptable salts and esters thereof] were prepared as β_3 -adrenoreceptor agonists. For example, coupling of (2R)-6-iodo-3,4-dihydro-2H-chromene-2-carboxylic acid and (1R)-2-amino-1-(3-pyridinyl)ethanol•2HCl with 1-hydroxybenzotriazole, 1-[3-dimethylaminopropyl]-3-ethylcarbodiimide•HCl, and TEA in CH₂Cl₂ gave the amide (744). Reduction using borane-dimethylsulfide complex in THF afforded the chromanmethanamine II (844). Over one hundred compds. of the invention demonstrated β_3 -adrenergic receptor agonist activity with EC₅₀ values \leq 1 μ M. I are useful in the treatment of β_3 -adrenergic receptor mediated conditions, including obesity, diabetes, gastrointestinal disorders, cardiovascular disorders, and urinary disorders. (no data).

IT 474114-60-BP, N-[[[(2R)-2-[[[(2R)-2-(6-amino-3-pyridinyl)-2-hydroxyethyl]amino]methyl]-3,4-dihydro-2H-chromen-6-yl]carbonyl]-N-methyl-L-phenylalanine

RL: PAC (Pharmacological activity); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)

(β_3 -adrenoreceptor agonist; preparation of chiral alkylaminochroman deriva. as β_3 -adrenoreceptor agonists)

RN 474114-60-B CAPLUS

CN L-Phenylalanine, N-[[[(2R)-2-[[[(2R)-2-(6-amino-3-pyridinyl)-2-hydroxyethyl]amino]methyl]-3,4-dihydro-2H-1-benzopyran-6-yl]carbonyl]-N-methyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

L4 ANSWER 55 OF 261 CAPLUS COPYRIGHT 2004 ACS ON STN

ACCESSION NUMBER: 2002-814137 CAPLUS

DOCUMENT NUMBER: 137:325439

TITLE: Preparation of hydroxy cyclohexenylphenyl benzopyrrolodiazepine and triazabenzazulene carboxamides as tocolytic oxytocin receptor antagonists

INVENTOR(S): Failli, Amedeo Arturo; Sanders, William Jennings; Trybulski, Eugene John

PATENT ASSIGNEE(S): Wyeth, John, and Brother Ltd., USA

SOURCE: PCT Int. Appl., 87 pp. CODEN: PIXXD2

DOCUMENT TYPE: Patent



LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2002083679	A1	20021024	WO 2002-US11529	20020411
W:	AR, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZM, ZW, AM, AZ, BY, BG, KZ, MD, RU, TJ, TJ, TM			
RW:	GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF, BJ, CF, CG, CI, CM, CA, GN, GO, GW, ML, MR, NE, SN, TD, TG			
US 2002183311	A1	20021205	US 2002-119973	20020410
EP 1385849	A1	20040204	EP 2002-721727	20020411
R:	AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR			
PRIORITY APPL. INFO.:			US 2001-283316P	20010412
			WO 2002-US11529	20020411
OTHER SOURCE(S):		MARPAT 137:325439		
GI				

L4 ANSWER 55 OF 261 CAPLUS COPYRIGHT 2004 ACS ON STN (Continued)



AB The present invention provides substituted 10,11-Dihydro-5H-benzof[1,2-a]pyrrolo[1,2-a][1,4]diazepine and 9,10-Dihydro-4H-3a,5,9-triazabenzof[1,2-a]pyrrolo[1,2-a][1,4]diazepine compds. (shown as I; 10-(5-chloro-4-(cyclohex-1-enyl)-2-methoxybenzoyl)-N-methyl-N-((2S,3R,4R,5R)-2,3,4,5,6-pentamethoxyhexyl)-10,11-dihydro-5H-pyrrolo[2,1-c][1,4]benzodiazepine-3-carboxamide) as well as methods and pharmaceutical compns. using these compds. for the treatment and/or prevention and/or suppression of disorders which may be remedied or alleviated by oxytocin antagonist activity, including prevention and/or suppression of preterm labor, suppression of labor at term prior to cesarean deliver, and for the treatment of dysmenorrhea. These compds. are also useful in enhancing fertility rates, enhancing survival rates and synchronizing estrus in farm animals. In I: the Z ring is R1- and R2-substituted o-phenylene or 2,3-pyridinediyl; R1 and R2 are H, (C1-C6)alkyl, halo, cyano, trifluoromethyl, hydroxy, amino, (C1-C6)alkylamino, (C1-C6)alkoxy, -OCF3, (C1-C6alkoxy)carbonyl, -NHCO[(C1-C6)alkyl], carboxy, -CONH2, -CONH(C1-C6)alkyl, or -CON[(C1-C6)alkyl]2. R3 is H, (C1-C6)alkyl, (C1-C6)alkoxy, hydroxy, amino, (C1-C6)alkylamino, -COalkyl(C1-C6), or halo. R4 is B-C wherein B is II and III and C is (un)substituted cyclohexenyl, 3,4-dihydro-1-naphthyl or 5,6-dihydro-8-quinolynyl where A is CH or N; R5, R6, and R7 are H, (C1-C6)alkyl, (C1-C6)alkoxy, hydroxy(C1-C6)alkyl, loweralkoxy(C1-C6)alkyl, (C2-C7)acyloxy(C1-C6)alkyl, (C3-C8)cycloalkylcarbonyl, (C2-C6)alkenyl, (C2-C6)alkynyl, (C3-C8)cycloalkyl, formyl, (C3-C8)cycloalkylcarbonyl, carboxy, loweralkoxy carbonyl, (C3-C8)cycloalkyloxy carbonyl, (arylloweralkyl)oxy carbonyl, carbamoyl, -OCH2CH2, halogen, haloloweralkyl including trifluoromethyl, -OCF3, -S(lower alkyl), -OC(O)N(lower alkyl)2, -CONH(lower alkyl), -CON(lower alkyl)2, lower alkylamino, diloweralkylamino, diloweralkylaminoloweralkyl, hydroxy, cyano, trifluoromethylthio, nitro, amino, lower alkylsulfonyl, aminosulfonyl, lower alkylaminosulfonyl, (un)substituted 2-dioxolanyl, Ph or naphthyl. R is acyclic or cyclic amino as defined in the claims. Although the methods of preparation are not claimed, 17 example preps. are included. Binding to membranes of Chinese Hamster Ovary (CHO) cell line

L4 ANSWER 55 OF 261 CAPLUS COPYRIGHT 2004 ACS ON STN (Continued)

stably transfected with human vasopressin V1a receptor subtype, human vasopressin V2 receptor subtype and human oxytocin receptor by approx. 17 I are reported.

IT 473665-66-6P, (2S)-2-[[10-(4-(Cyclohex-1-enyl)-3-methylbenzoyl)-10,11-dihydro-5H-pyrrolo[2,1-c][1,4]benzodiazepine-3-carbonyl]amino]-3-(4-hydroxyphenyl)propionic acid ethyl ester


RL: PAC (Pharmacological activity); RCT (Reactant); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); RACT (Reactant or reagent); USES (Uses)

(drug candidate; preparation of hydroxy cyclohexenylphenyl benzopyrrolodiazepine and triazabenzazulene carboxamides as tocolytic oxytocin receptor antagonists)

RN 473665-66-6 CAPLUS

CN L-Tyrosine, N-[[10-(4-(1-cyclohexen-1-yl)-3-methylbenzoyl)-10,11-dihydro-5H-pyrrolo[2,1-c][1,4]benzodiazepin-3-yl]carbonyl]-, ethyl ester (9CI) (CA INDEX NAME)

Absolute stereochemistry.



REFERENCE COUNT: 2 THERE ARE 2 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 56 OF 261 CAPLUS COPYRIGHT 2004 ACS ON STN

ACCESSION NUMBER: 2002:696111 CAPLUS

DOCUMENT NUMBER: 137:228607

TITLE: Crystal structure and three-dimensional structure of human Cdc25 catalytic domains and its use in designing peptidomimetic inhibitors

INVENTOR(S): Taylor, Neil R.; Borhani, David; Epstein, David; Rudolph, Johannes; Ritter, Kurt; Fujimori, Taro; Robinson, Simon; Eckstein, Jens; Haupt, Andreas; Walker, Nigel; Dixon, Richard W.; Choquette, Deborah; Blanchard, Jill; Kluge, Arthur; Pal, Kollol; Bockovich, Nicholas; Come, Jon; Hediger, Mark

PATENT ASSIGNEE(S): BASF Aktiengesellschaft, Germany; GPC Biotech Inc.

SOURCE: PCT Int. Appl., 351 pp. CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2002070680	A1	20020912	WO 2001-US6587	20010301

W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW, AM, AZ, BY, BG, KZ, MD, RU, TJ, TM

RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF, BJ, CF, CG, CI, CM, CA, GN, GW, ML, MR, NE, SN, TD, TG

PRIORITY APPL. INFO.: WO 2001-US6587 20010301

OTHER SOURCE(S): MARPAT 137:228607

AB Due to its role in regulating the cell cycle, Cdc25 (a family of dual specificity phosphatases) is a potential target for therapies aimed at controlling proliferative diseases, but rational, structure-based design has not been possible because of the lack of accurate 3-dimensional data. The present invention relates to polypeptides which comprises the ligand binding domain of human Cdc25 proteins, crystalline forms of these polypeptides, and the use of these crystalline forms to determine the 3-dimensional

structure of the catalytic domain of Cdc25. In particular, a high resolution crystal structure was obtained for the polypeptide denoted Cdc25B(ANB), comprising residues Glu-368 through Arg-562 of human Cdc25B, complexed with a pentapeptide inhibitor denoted cdc1249 (2-methoxynaphthyl-1-carboxy-(4-sulfomethyl)-L-Phe-L-Glu-L-Glu-L-naphthylalanine-L-Glu-amide). The invention also relates to the use of the 3-dimensional structure of the Cdc25 catalytic domain in methods of designing and/or identifying potential inhibitors of Cdc25 activity, for example, compds. which inhibit the binding of a native substrate to the Cdc25 catalytic domain. The syntheses and structures of a large number of putative pentapeptide inhibitors are also provided. Such inhibitors have potential in the treatment of diseases associated with excessive cellular proliferation, such as cancer, restenosis, reocclusion of coronary artery, and inflammation.


IT 329274-06-8P

RL: SPN (Synthetic preparation); PREP (Preparation)


(crystal structure and three-dimensional structure of human Cdc25 catalytic domains and its use in designing peptidomimetic inhibitors)

L4 ANSWER 56 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)
 RN 329274-06-8 CAPLUS
 CN L-Norvalinamide, N-(1-dibenzofuran-2-yl)-4-(sulfomethyl)-L-phenylalanyl-L-norvalyl-2-methyl-L-prolyl-3-benzo[b]thien-3-yl-L-alanyl-5-carboxy-N-(1,1-dimethylethyl)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



PAGE 1-A




PAGE 1-B

REFERENCE COUNT: 8 THERE ARE 8 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 57 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
 ACCESSION NUMBER: 2002:692540 CAPLUS
 DOCUMENT NUMBER: 138:338460
 TITLE: Modifications of the Tic residue in TIPP-peptides
 AUTHOR(S): Tourwe, D.; Van Cauwenberghe, S.; Vanommeslaeghe, K.; Mannekens, E.; Geerlings, P.; Toth, G.; Peter, A.; Csombos, J.
 CORPORATE SOURCE: Department of Organic Chemistry, Vrije Universiteit Brussel, Brussels, B-1050, Belg.
 SOURCE: Peptides: The Wave of the Future, Proceedings of the Second International and the Seventeenth American Peptide Symposium, San Diego, CA, United States, June 9-14, 2001 (2001), 683-684. Editor(s): Lebl, Michal; Houghten, Richard A. American Peptide Society: San Diego, Calif.
 CODEN: 69DBAL; ISBN: 0-9715560-0-8
 DOCUMENT TYPE: Conference
 LANGUAGE: English
 AB A symposium report. Four 1,2,3,4-tetrahydroisoquinoline-3-carboxylic acid (Tic) analogs were prepared and used as replacements for Tic in the selective δ -opioid antagonist N-Tyr-Tic-Phe-Phe-OH (TIPP). Mol. modeling of tripeptide TIP analogs containing modified Tics with (3S) configuration indicated that the exact distance between the aromatic rings in TIP and the positioning of the phenolic group are crucial for δ -affinity. The positioning of the Tic carbonyl also determined the orientation of the Phe residue.
 IT 517891-95-1P
 RL: PAC (Pharmacological activity); PRP (Properties); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation) (modifications of Tic residue in TIPP-peptides)
 RN 517891-95-1 CAPLUS
 CN L-Phenylalanine, L-tyrosyl-(1aS,7bS)-1,2,3,7b-tetrahydro-1aH-cycloprop[c]isoquinoline-1a-carbonyl-L-phenylalanyl- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



REFERENCE COUNT: 5 THERE ARE 5 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 58 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
 ACCESSION NUMBER: 2002:692518 CAPLUS
 DOCUMENT NUMBER: 138:268230
 TITLE: Design, synthesis and biological evaluation of pilicides: inhibitors of pilus assembly in pathogenic bacteria
 AUTHOR(S): Larsson, Andreas; Emteneas, Hans; Svensson, Anette; Pinkner, Jerome S.; Hultgren, Scott J.; Almqvist, Fredrik; Kihlberg, Jan
 CORPORATE SOURCE: Department of Organic Chemistry, Umea University, Umea, SE-901 87, Swed.
 SOURCE: Peptides: The Wave of the Future, Proceedings of the Second International and the Seventeenth American Peptide Symposium, San Diego, CA, United States, June 9-14, 2001 (2001), 636-637. Editor(s): Lebl, Michal; Houghten, Richard A. American Peptide Society: San Diego, Calif.
 CODEN: 69DBAL; ISBN: 0-9715560-0-8
 DOCUMENT TYPE: Conference
 LANGUAGE: English
 AB A crystal structure of the complex between the periplasmic chaperone PapD, involved in assembly of P Pili in uropathogenic Escherichia coli, and a 19-mer peptide corresponding to the C-terminus of the adhesin PapG was used to develop two classes of peptidomimetics as potential inhibitors of the chaperone/subunit complex by rational drug design. The amino acid derivs. were synthesized through an N-alkylation of an amino acid followed by acylation of the resulting secondary amine. The 2-pyridinones were obtained via a novel procedure based on the use of acid chlorides and nitriles as starting materials. Within the amino acid derivs. and 2-pyridinones, which bind to periplasmic chaperones and even dissociate chaperone, pilus subunit complexes were detected.
 IT 503305-58-6
 RL: PAC (Pharmacological activity); THU (Therapeutic use); BIOL (Biological study); USES (Uses) (design and synthesis and biol. evaluation of pilicides such as inhibitors of pilus assembly in pathogenic bacteria that dissociate periplasmic chaperone-pilus subunit complexes)
 RN 503305-58-6 CAPLUS
 CN L-Tyrosine, N-[3-(1H-indol-3-yl)propyl]-N-[(2-oxo-2H-1-benzopyran-3-yl)carbonyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



L4 ANSWER 58 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)
 REFERENCE COUNT: 3 THERE ARE 3 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 59 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
 ACCESSION NUMBER: 2002:637480 CAPLUS
 DOCUMENT NUMBER: 137:190724
 TITLE: Melanocortin metallopeptides for treatment of sexual dysfunction
 INVENTOR(S): Sharma, Shubh D.; Shi, Yi-qun; Yang, Wei; Cai, Hui-zhi; Shadiack, Annette
 PATENT ASSIGNEE(S): Palatin Technologies, Inc., USA
 SOURCE: PCT Int. Appl., 58 pp.
 CODEN: PIXXD2
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2002064091	A2	20020822	WO 2002-US4431	20020213
WO 2002064091	A3	20030313		

W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM

RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF, BJ, CF, CG, CI, CM, CA, GN, GQ, GW, ML, MR, NE, SN, TD, TG

US 2004038897 A1 20040226 US 2003-640755 20030813

PRIORITY APPLN. INFO.: US 2001-268591P P 20010213
 WO 2002-US4431 A 20020213


OTHER SOURCE(S): MARPAT 137:190724
 AB Metallopeptides are provided for use in treatment of sexual dysfunction in mammals. The metallopeptides are agonists for at least one of melanocortin-3 or melanocortin-4 receptors. The metallopeptides are conformationally fixed on complexation of a metal ion-binding portion thereof with a metal ion. Also provided are metallopeptides that are antagonists for at least one of melanocortin-3 or melanocortin-4 receptors.

IT 448903-51-3
 RL: PAC (Pharmacological activity); PRP (Properties); THU (Therapeutic use); BIOL (Biological study); USES (Uses)
 (melanocortin metallopeptides for treatment of sexual dysfunction)

RN 448903-51-3 CAPLUS
 CN L-Cysteineamide, (3S)-1,2,3,4-tetrahydro-3-isoquinolinecarbonyl-4-chloro-D-phenylalanyl-L-arginyl-L-tryptophyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

L4 ANSWER 59 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)



L4 ANSWER 61 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
 ACCESSION NUMBER: 2002:575075 CAPLUS
 DOCUMENT NUMBER: 137:140779
 TITLE: Preparation of piperazine- and piperidine-derivatives as melanocortin receptor agonists
 INVENTOR(S): Briner, Karin; Doecke, Christopher William; Mancoso, Vincent; Martinelli, Michael John; Richardson, Timothy Ivo; Rothhaar, Roger Ryan; Shi, Qing; Xie, Chaoyu
 PATENT ASSIGNEE(S): Eli Lilly and Company, USA
 SOURCE: PCT Int. Appl., 272 pp.
 CODEN: PIXXD2
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 2
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2002059117	A1	20020801	WO 2002-US515	20020123
WO 2002059117	C1	20031106		

W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM

RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF, BJ, CF, CG, CI, CM, CA, GN, GQ, GW, ML, MR, NE, SN, TD, TG


EP 1370558 A1 20031117 EP 2002-701922 20020123

R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR

PRIORITY APPLN. INFO.: US 2001-263471P P 20010123
 WO 2002-US515 W 20020123

OTHER SOURCE(S): MARPAT 137:140779

GI



AB The compds. of formula I (G = CR1, or N; A = alkyl, or cycloalkyl; L and L1 = H, or (together) oxo; T = substituted indolyl, or pyrazinyl; X = CH2, or CH2CH2; Z = (CH2)n; R1 = H, alkyl, Ph, alkylaryl, alkylcarboxamide, cycloalkyl, or oxo; R2 = H, halo, alkyl, alkylsulfonyl, cycloalkyl, alkylaryl, or haloalkyl; R3 = (un)substituted aryl, or thienyl; R4 = H, alkyl, cycloalkyl, etc.; R5 = NH2, NPh2, alkylamide, alkylsulfonylamide, NHCOH, NHCONH2, NHSO2NH2, (un)substituted


L4 ANSWER 60 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
 ACCESSION NUMBER: 2002:576051 CAPLUS
 DOCUMENT NUMBER: 137:279456
 TITLE: Solid Phase Synthesis and Evaluation of Tyr-Tic-Phe-Phe(p-NMCOCH2Br) ([Phe(p-bromoacetamide)4]TIPP), a Potent Affinity Label for δ Opioid Receptors
 AUTHOR(S): Kumar, Vivek; Murray, Thomas P.; Aldrich, Jane V.
 CORPORATE SOURCE: Department of Pharmaceutical Sciences School of Pharmacy, University of Maryland, Baltimore, MD, 21201, USA
 SOURCE: Journal of Medicinal Chemistry (2002), 45(18), 3820-3823
 CODEN: JMCMAR; ISSN: 0022-2623
 PUBLISHER: American Chemical Society
 DOCUMENT TYPE: Journal
 LANGUAGE: English

AB Derivs. of the δ -opioid receptor-selective peptide H-Tyr-Tic-Phe-Phe-OH (TIPP) containing a p-bromoacetamide moiety on the Ph ring of Phe3 or Phe4 were prepared by solid-phase synthesis. [Phe(p-NMCOCH2Br)4]TIPP exhibited high affinity for cloned δ receptors (IC50 = 5.4 nM), and incubation with only 2.5 nM resulted in 85% wash resistant inhibition of radioligand binding to δ receptors. This peptide is a potent affinity label for further study of δ opioid receptors.

IT 320782-32-99
 RL: BSU (Biological study, unclassified); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation)
 (solid-phase preparation and δ -opioid receptor-binding affinity measurements of TIPP peptides with bromoacetamide groups)

RN 320782-32-9 CAPLUS
 CN L-Phenylalanine, L-tyrosyl-(3S)-1,2,3,4-tetrahydro-3-isoquinolinecarbonyl-4-amino-L-phenylalanyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




REFERENCE COUNT: 28 THERE ARE 28 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 61 OF 261 CAPLUS COPYRIGHT 2004 ACS on STM (Continued)
heterocyclyl, etc.; n = 0-8, m = 0-1, and p = 0-4), pharmaceutically acceptable salts, or stereoisomers were prepd. as melanocortin receptor agonists for treatment of obesity, diabetes and male and/or female sexual dysfunction. Thus, coupling of 2-[(2-tert-butoxycarbonyl-1,2,3,4-tetrahydroisoquinolin-3-ylmethyl)amino]-3-(4-chlorophenyl)propionate with 3-(2-piperazin-1-yltrifluoromethylphenoxy)-S-pyrrolidine-1-carboxylic acid tert-Bu ester, followed by deprotection and addn. of HCl, gave 3-D-(4-chlorophenyl)-1-[4-[5-(trifluoromethyl-2-S-(pyrrolidin-3-yloxy)phenyl)piperazin-1-yl]-2-D-[(1,2,3,4-tetrahydroisoquinoline-3-ylmethyl)amino]propan-1-one hydrochloride in 84% yield.

IT 252008-71-2P
RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)
(preparation of piperazine- and piperidine-derivs. as melanocortin receptor agonists for treatment of obesity, diabetes and sexual dysfunction)

RN 252008-71-2 CAPLUS
CN 2-[(1H)-isoquinoline-3-carboxylic acid, 3-[[[(1R)-1-carboxy-2-(4-chlorophenyl)ethyl]amino]carbonyl]-3,4-dihydro-, 2-(1,1-dimethylethyl) ester, (3R)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



REFERENCE COUNT: 1 THERE ARE 1 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 62 OF 261 CAPLUS COPYRIGHT 2004 ACS on STM
ACCESSION NUMBER: 2002:575056 CAPLUS
DOCUMENT NUMBER: 137:140777
TITLE: Preparation of piperazinyl and hexahydro-1,4-diazepinyl amino acid derivatives as melanocortin receptor agonists
INVENTOR(S): Biggers, Christopher Kelly; Briner, Karin; Doecke, Christopher William; Fisher, Matthew Joseph; Hertel, Larry Wayne; Mancoso, Vincent; Martinelli, Michael John; Mayer, John Philip; Ornstein, Paul Leslie; Richardson, Timothy Ivo; Shah, Jikesh Arvind; Shi, Qing; Wu, Zhipel; Xie, Chaoyu
PATENT ASSIGNEE(S): Eli Lilly and Company, USA
SOURCE: PCT Int. Appl., 356 pp.
CODEN: PIXXD2
DOCUMENT TYPE: Patent
LANGUAGE: English
FAMILY ACC. NUM. COUNT: 2
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2002059108	A1	20020801	WO 2002-US517	20020123
W:	AR, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MY, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW, AM, AZ, BY, BG, KZ, MD, RU, TJ, TM			
RW:	GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG			
EP 1368340	A1	20031210	EP 2002-714719	20020123
R:	AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR			
PRIORITY APPL. INFO.:			US 2001-263471P	P 20010123
			WO 2002-US517	W 20020123
OTHER SOURCE(S):			MARPAT 137:140777	
GI				




AB The invention relates to melanocortin receptor (MC-R) agonists I [X = CH₂ or CH₂CH₂; L₁ = H₂ or oxo; T = isoquinolinyl or tetrahydro derivative,

L4 ANSWER 62 OF 261 CAPLUS COPYRIGHT 2004 ACS on STM (Continued)
isoindolinyl, or piperazinyl; n = 0-8; R = H, OH, CN, NO₂, halo, alkyl, acyl, etc.; R₁ = H, alkyl, alkylcarbonyl, (D)phenyl, (D)cycloalkyl, or oxo (unless amide is formed); p = 0-5; R₃ = (un)substituted aryl or thienyl; R₄ = H, alkyl, acyl, cycloalkyl, or alkoxyalkyl, or their pharmaceutically acceptable salts or stereoisomers, which are useful in the treatment of obesity, diabetes, and male and/or female sexual dysfunction. Comps. 1 comprise three domains, i.e., a piperazinyl or hexahydro-1,4-diazepinyl fragment, an amino acid, and a radical CLL1(CH₂)_n-T. Thus, 1-(D-Tic-4-Cl-D-Phe)-4-[2-(methanesulfonylamino)phenyl]piperazine (Tic = 1,2,3,4-tetrahydroisoquinoline-3-carbonyl, claimed compd.) was prepd. via acylation of the piperazine moiety.

IT 452330-88-0
RL: RCT (Reactant); RACT (Reactant or reagent)
(preparation of piperazinyl and hexahydrodiazepinyl amino acid deriva. as melanocortin receptor agonists)

RN 452330-88-0 CAPLUS
CN D-Phenylalanine, 4-chloro-N-(3-isoquinolinylcarbonyl)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



REFERENCE COUNT: 2 THERE ARE 2 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 63 OF 261 CAPLUS COPYRIGHT 2004 ACS on STM
ACCESSION NUMBER: 2002:575056 CAPLUS
DOCUMENT NUMBER: 137:140776
TITLE: Preparation of piperidinyl and piperazinyl amino acid derivatives as melanocortin receptor agonists
INVENTOR(S): Backer, Ryan Thomas; Briner, Karin; Doecke, Christopher William; Fisher, Matthew Joseph; Kuklish, Steven Lee; Mancoso, Vincent; Martinelli, Michael John; Mullaney, Jeffrey Thomas; Xie, Chaoyu
PATENT ASSIGNEE(S): Eli Lilly and Company, USA
SOURCE: PCT Int. Appl., 263 pp.
CODEN: PIXXD2
DOCUMENT TYPE: Patent
LANGUAGE: English
FAMILY ACC. NUM. COUNT: 1
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2002059107	A1	20020801	WO 2002-US516	20020123
W:	AR, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MY, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW, AM, AZ, BY, BG, KZ, MD, RU, TJ, TM			
RW:	GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG			
EP 1368339	A1	20031210	EP 2002-701923	20020123
R:	AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR			
US 2004058936	A1	20040325	US 2003-466249	20030711
PRIORITY APPL. INFO.:			US 2001-263595P	P 20010123
			WO 2002-US516	W 20020123
OTHER SOURCE(S):			MARPAT 137:140776	
GI				




AB The invention relates to melanocortin receptor (MC-R) agonists I [G = CR₁ or N; L₁ = H₂ or oxo; T = isoquinolinyl or tetrahydro derivative, isoindolinyl, or piperazinyl; n = 0-8; R = H, OH, CN, NO₂, halo, alkyl, acyl, etc.; R₁ = H, alkyl, alkylcarbonyl, (D)phenyl, (D)cycloalkyl, or oxo (unless amide is formed); p = 0-4; CR₂CR₂ is a 5- or 6-membered

L4 ANSWER 63 OF 261 CAPLUS COPYRIGHT 2004 ACS ON STN (Continued)
 carbocycle optionally substituted by 1-3 groups R; R3 = (un)substituted
 aryl or thienyl; R4 = H, alkyl, acyl, cycloalkyl, or alkoxyalkyl, or
 their pharmaceutically-acceptable salts or stereoisomers, which are useful
 in the treatment of obesity, diabetes, and male and/or female sexual
 dysfunction. Comps. I comprise three domains, i.e., a piperidino or
 piperazinyl fragment, an amino acid, and a radical CLL1(CH2)n-T. Thus,
 1,2,3,4-tetrahydroisoquinoline-3-carboxylic acid [1-(4-chlorobenzyl)-2-[4-
 [2-(methylsulfonyl)-1,2,3,4-tetrahydroisoquinolin-8-yl]piperazin-1-yl]-2-
 oxoethylamide (claimed compd.) was prepd. via acylation of the
 piperazine moiety.

IT 252008-71-2P
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT
 (Reactant or reagent)
 (preparation of piperidinyl and piperazinyl amino acid derivs. as
 melanocortin receptor agonists)

RN 252008-71-2 CAPLUS
 CN 2(1H)-Isoquinolinecarboxylic acid, 3-[[[(1R)-1-carboxy-2-(4-
 chlorophenyl)ethyl]amino]carbonyl]-3,4-dihydro-, 2-(1,1-dimethylethyl)
 ester, (3R)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



REFERENCE COUNT: 6 THERE ARE 6 CITED REFERENCES AVAILABLE FOR THIS
 RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 64 OF 261 CAPLUS COPYRIGHT 2004 ACS ON STN
 ACCESSION NUMBER: 2002:575055 CAPLUS
 DOCUMENT NUMBER: 137:140775
 TITLE: Preparation of piperazinyl and hexahydro-1,4-
 diazepinyl amino acid derivatives as melanocortin
 receptor agonists


INVENTOR(S): Backer, Ryan Thomas; Briner, Karin; Collado Cano,
 Ivan; De Frutos-Garica, Oscar; Doecke, Christopher
 William; Fisher, Matthew Joseph; Garcia-Paredes,
 Cristina; Kuklish, Steven Lee; Mancoso, Vincent;
 Martinelli, Michael John; Matteo Herranz, Ana Isabel;
 Mullaney, Jeffrey Thomas; Ornstein, Paul Leslie; Wu,
 Zhipei; Xie, Chaoyu

PATENT ASSIGNEE(S): Eli Lilly and Company, USA

SOURCE: PCT Int. Appl., 554 pp.
 CODEN: PIXXD2

DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2002059095	A1	20020801	WO 2002-US518	20020123
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, ME, MK, MN, MW, MX, MY, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW, AM, AZ, BY, BG, KZ, MD, RU, TJ, TM			
RW:	GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZM, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG			
EP 1358163	A1	20031105	EP 2002-701924	20020123
R:	AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR			
PRIORITY APPLN. INFO.:			US 2001-263380P P	20010123
			WO 2002-US518 W	20020123
OTHER SOURCE(S):			CASREACT 137:140775; MARPAT 137:140775	
GI				




AB The invention relates to melanocortin receptor (MC-R) agonists I [X = CH2
 or CH2CH2; L1 = H2 or oxo; R2 = H, alkyl, alkylcarbamoyl, (D)phenyl,

L4 ANSWER 64 OF 261 CAPLUS COPYRIGHT 2004 ACS ON STN (Continued)
 (D)cyclohexyl, or oxo if adjacent to N-Q; p = 0-4; R3 = (un)substituted
 Ph, aryl, or thienyl; R4 = H, alkyl, alkenyl, alkanoyl, or (D)phenyl; Q =
 various carbon-attached groups; T = isoquinolinyl or tetrahydro deriv.,
 isoindolinyl, or piperazinyl; n = 0-8] which are useful in the treatment
 of obesity, diabetes, and male and/or female sexual dysfunction. Comps.
 I comprise three domains, i.e., a piperazinyl or hexahydro-1,4-diazepinyl
 fragment, an amino acid, and a radical CLL1(CH2)n-T. Thus,
 N-[1-(4-chlorobenzyl)-2-[4-[1-(cyclohexylmethyl)-2-
 morpholinoethyl]piperazin-1-yl]-2-oxoethyl]-2-(2,3-dihydro-1H-isoindol-1-
 yl)acetamide tris(trifluoroacetate) salt was prepd. via acylation of the
 piperazine moiety and showed EC50 = 69.3 nM in the MC-4 agonist assay.

IT 252008-71-2P
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT
 (Reactant or reagent)
 (preparation of piperazinyl and hexahydrodiazepinyl amino acid deriva. as
 melanocortin receptor agonists)

RN 252008-71-2 CAPLUS
 CN 2(1H)-Isoquinolinecarboxylic acid, 3-[[[(1R)-1-carboxy-2-(4-
 chlorophenyl)ethyl]amino]carbonyl]-3,4-dihydro-, 2-(1,1-dimethylethyl)
 ester, (3R)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



REFERENCE COUNT: 4 THERE ARE 4 CITED REFERENCES AVAILABLE FOR THIS
 RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 65 OF 261 CAPLUS COPYRIGHT 2004 ACS ON STN
 ACCESSION NUMBER: 2002:484863 CAPLUS
 DOCUMENT NUMBER: 137:47448
 TITLE: Preparation of substituted phenylalaninol derivatives
 as protein tyrosine phosphatase inhibitors


INVENTOR(S): Larsen, Scott D.; May, Paul D.; Bleasdale, John E.;
 Liljeborg, Charlotta; Schostarez, Heinrich Josef;
 Barf, Tjeerd; Nilsson, Marianne

PATENT ASSIGNEE(S): USA

SOURCE: U.S., 144 pp., Cont.-in-part of U.S. Ser. No. 138,642.
 CODEN: USXXAM

DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 3
 PATENT INFORMATION:


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 6410585	B1	20020625	US 1999-265410	19990310
US 6353023	B1	20020305	US 1998-138642	19980824
WO 2000053583	A1	20000914	WO 2000-US6022	20000309
W:	AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, AM, AZ, BY, BG, KZ, MD, RU, TJ, TM			
RW:	GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG			
EP 1161421	A1	20011212	EP 2000-917793	20000309
R:	AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO			
JP 2002539115	T2	20021119	JP 2000-604023	20000309
PRIORITY APPLN. INFO.:			US 1997-57730P P	19970828
			US 1998-138642 A2	19980824
			US 1999-265410 A	19990310
			WO 2000-US6022 W	20000309
OTHER SOURCE(S):			MARPAT 137:47448	
GI				



AB The invention comprises phenylalaninol deriva., e.g., I [R1 = OSO3H,
 OCH(CO2R5)2, OCH2CO2R5, OCH(CO2R5)CH2CO2R5, OC(CO2R5)CHCO2R5,
 CH2CH(CO2R5)2, CH2C(CO2R5)2, OCH2CONHCH, N(CH2CO2R5)2, OCHFCO2R5 (R5 = H,
 alkyl, alkylphenyl); R2 = CHR7NHXR6, group Q (R6 = alkyl, alkylCONH2,
 alkyl-NHCO2R5, etc.; R7 = H, any group given for R6); R10 = H, CO2R5,
 CONHCH, 5-tetrazolyl, F, OCH2CO2R5], or their pharmaceutically acceptable
 salts, as small mol. weight, non-peptidic inhibitors of protein tyrosine
 phosphatase 1 (PTP1) which are useful for the treatment and/or prevention
 of non-insulin dependent diabetes mellitus. Thus, 5-[(2S)-2-[(2S)-2-

L4 ANSWER 65 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)
 [(tert-butoxycarbonylamino)-3-phenylpropanoylamino]-3-hydroxypropyl-2-(carboxymethoxy)benzoic acid (claimed compd.) was prepd. and showed 80% inhibition of protein tyrosine phosphatase 1B at a concn. of 10 μ M.
 IT 292834-82-3P
 RL: CPN (Combinatorial preparation); PAC (Pharmacological activity); THU (Therapeutic use); BIOL (Biological study); CMBI (Combinatorial study); PREP (Preparation); USES (Uses)
 (preparation of substituted phenylalanine derivs. as protein tyrosine phosphatase inhibitors)
 RN 292834-82-3 CAPLUS
 CN Benzoic acid, 2-(carboxymethoxy)-5-[(2S)-3-oxo-3-[(4-phenylbutyl)amino]-2-[(3-quinolinylcarbonyl)amino]propyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




REFERENCE COUNT: 19 THERE ARE 19 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 66 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
 ACCESSION NUMBER: 2002:477236 CAPLUS
 DOCUMENT NUMBER: 137:197780
 TITLE: Molecular Imprinting for the Recognition of N-Terminal Histidine Peptides in Aqueous Solution
 AUTHOR(S): Hart, Bradley R.; Shea, Kenneth J.
 CORPORATE SOURCE: Department of Chemistry, University of California, Irvine, CA, 92697-2025, USA
 SOURCE: Macromolecules (2002), 35(16), 6192-6201
 CODEN: MAMORX; ISSN: 0024-9297
 PUBLISHER: American Chemical Society
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 AB A new procedure for creating macromol. receptors for peptides using mol. imprinting has been developed. The polymeric receptor exhibits selective uptake of specific N-terminal histidine containing sequences of simple dipeptides. The polymerization and binding are carried out in water. The approach utilizes a strong Ni(II)-His binding to attract the N-terminus histidine of the dipeptide to the polymer surface and secondary interactions between peptide and polymer to discriminate between the peptide sequence. These developments are enabled by utilizing an aqueous based monomer formulation that includes N,N'-ethylenebis(acrylamide) as a water-soluble crosslinking monomer and a polymerizable NTA ligand, which can be used to incorporate nickel and other metals into these polyacrylamides. The Ni-NTA complex provides a strong histidine binding site that draws the dipeptide to the polymer surface. Mild polymerization conditions that utilize low concns. of water-soluble initiator and low-temperature result in quant. polymer yields. Variation of monomer composition reveals an optimum crosslinking for achieving maximum selectivity for these polymers.
 IT 452928-69-7P
 RL: BSU (Biological study, unclassified); PEP (Physical, engineering or chemical process); PYP (Physical process); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation); PROC (Process)
 (protocol for creating macromol. receptors for peptides using mol. imprinting with nickel-nitrilotriacetic acid complex)
 RN 452928-69-7 CAPLUS
 CN Nickelate(2-), [N2,N2-bis[(carboxy- κ O)methyl]-N6-(2-methyl-1-oxo-2-propenyl)-L-lysinate(3-)- κ N2, κ O1](D-histidyl- κ N, κ N3-L-phenylalaninate)-, (OC-6-52)-, polymer with N,N'-1,2-ethanediyldis[2-propenamide] and 2-propenamide (9CI) (CA INDEX NAME)
 CM 1
 CRN 452928-68-6
 CMF C29 H36 N6 Ni O10
 CCI CCS


L4 ANSWER 66 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)

PAGE 1-A



PAGE 1-B


L4 ANSWER 66 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)
 CRN 79-06-1
 CMF C3 H5 N O



REFERENCE COUNT: 35 THERE ARE 35 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

— Ph

CM 2
 CRN 2956-58-3
 CMF C8 H12 N2 O2



CM 3

L4 ANSWER 67 OF 261 CAPLUS COPYRIGHT 2004 ACS ON STN
 ACCESSION NUMBER: 2002:394477 CAPLUS
 DOCUMENT NUMBER: 137:103998

TITLE: Structure-Activity Relationships of the Melanocortin Tetrapeptide Ac-His-DPhe-Arg-Trp-NH₂ at the Mouse Melanocortin Receptors. 1. Modifications at the His Position

AUTHOR(S): Holder, Jerry Ryan; Bauzo, Rayna M.; Xiang, Zhimin; Haskell-Luevano, Carrie

CORPORATE SOURCE: Department of Medicinal Chemistry, University of Florida, Gainesville, FL, 32610, USA

SOURCE: Journal of Medicinal Chemistry (2002), 45(13), 2801-2810

CODEN: JMCMAR; ISSN: 0022-2623

PUBLISHER: American Chemical Society

DOCUMENT TYPE: Journal

LANGUAGE: English

AB The melanocortin pathway is an important participant in obesity and energy homeostasis. The centrally located melanocortin-3 and melanocortin-4 receptors (MC3R, MC4R) are involved in the metabolic and food intake aspects of energy homeostasis and are stimulated by melanocortin agonists such as α -melanocyte stimulation hormone (α -MSH). The melanocortin agonists contain the putative message sequence "His-Phe-Arg-Trp", and it has been well documented that inversion of chirality of the Phe to DPhe results in a dramatic increase in melanocortin receptor potency. Herein, the authors report a tetrapeptide library based on the template Ac-His-DPhe-Arg-Trp-NH₂, consisting of 17 members that have been modified at the His6 position (α -MSH numbering) and pharmacol. characterized for agonist activity at the mouse melanocortin receptors MC1R, MC3R, MC4R, and MC5R. These studies provide further exptl. evidence that the His6 position can determine MC4R vs. MC3R agonist selectivity and that chemical nonreactive side chains may be substituted for the imidazole ring (generally needs to be side chain protected in syntheses) in the design of MC4R-selective, small-mol., non-peptide agonists. Specifically, the tetrapeptide containing the amino-2-naphthylcarboxylic acid (Anc) amino acid at the His position resulted in a potent agonist at the mMC4R (EC₅₀ = 21 nM), was a weak mMC3R micromolar antagonist (pA₂ = 5.6, K_i = 2.5 μ M), and possessed >4700-fold agonist selectivity for the MC4R vs. the MC3R. Substitution of the His6 amino acid in the tetrapeptide template by the Phe, Anc, 3-(2-thienyl)alanine (2Thi), and 3-(4-pyridinyl)alanine (4-Pal) resulted in equipotency or only up to a 7-fold decrease in potency, compared to the His6-containing tetrapeptide at the mMC4R, demonstrating that these amino acid side chains may be substituted for the imidazole in the design of MC4R-selective non-peptide mols.


IT 443789-84-2P
 RL: RSU (Biological study, unclassified); PRP (Properties); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation)
 (structure-activity relationships of melanocortin tetrapeptide analogs at mouse melanocortin receptors)

RN 443789-84-2 CAPLUS

CN L-Tryptophanamide, (3S)-2-acetyl-1,2,3,4-tetrahydro-3-isoquinolinecarboxyl-D-phenylalanyl-L-arginyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

L4 ANSWER 67 OF 261 CAPLUS COPYRIGHT 2004 ACS ON STN (Continued)



REFERENCE COUNT: 49 THERE ARE 49 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 68 OF 261 CAPLUS COPYRIGHT 2004 ACS ON STN

ACCESSION NUMBER: 2002:392338 CAPLUS

DOCUMENT NUMBER: 137:140503

TITLE: Parallel Solid-Phase Synthesis of Triasubstituted Triazinobenzimidazolidiones

AUTHOR(S): Klein, Gerard; Acharya, Achyuta N.; Ostresh, John M.; Houghten, Richard A.

CORPORATE SOURCE: Torrey Pines Institute for Molecular Studies, San Diego, CA, 92121, USA

SOURCE: Journal of Combinatorial Chemistry (2002), 4(4), 345-351

CODEN: JCHFFP; ISSN: 1520-4766


PUBLISHER: American Chemical Society

DOCUMENT TYPE: Journal

LANGUAGE: English

OTHER SOURCE(S): CASREACT 137:140503

GI



AB An efficient method for the solid-phase synthesis of triasubstituted [1,3,5]triazino[1,2-a]benzimidazole-2,4(3H,10H)-diones I (R₁ = Me, Me₂CHCH₂, HOCH₂, PhCH₂, 4-FC₆H₄CH₂; R₂ = Bu, cyclopentyl, EtCHMe, etc.; R₃ = Me, 4-FC₆H₄CH₂, 3-MeC₆H₄CH₂, 2,4-F₂C₆H₃CH₂, 3-MeOC₆H₄CH₂) from resin-bound amino acids was developed. First, N-acylation of the primary amine of a resin-bound amino acid with 4-fluoro-3-nitrobenzoic acid, followed by displacement of the fluoro group with an amine and reduction of the nitro group, generated a resin-bound o-dianilino derivative. The dianilino compound was treated with cyanogen bromide to generate the corresponding iminobenzimidazole, which, following treatment with N-(chlorocarbonyl)isocyanate, afforded the resin-bound triazinedione derivative. Alkylation of the triazinedione compound with an alkyl halide following cleavage of the solid-support gave I.


IT 444813-51-8P
 RL: SPN (Synthetic preparation); PREP (Preparation)
 (parallel solid-phase synthesis of triasubstituted triazinobenzimidazolidiones)

RN 444813-51-8 CAPLUS

CN 1,3,5-Triazino[1,2-a]benzimidazole-7-carboxamide, N-[(1S)-2-amino-1-[(4-fluorophenyl)methyl]-2-oxoethyl]-10-butyl-2,3,4,10-tetrahydro-3-methyl-2,4-dioxo- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

L4 ANSWER 68 OF 261 CAPLUS COPYRIGHT 2004 ACS ON STN (Continued)



REFERENCE COUNT: 33 THERE ARE 33 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 69 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
 ACCESSION NUMBER: 2002:378541 CAPLUS
 DOCUMENT NUMBER: 136:386402
 TITLE: Preparation of alkenylamino acids as proteasome inhibitors
 INVENTOR(S): Kono, Yasushi; Ando, Naoki; Sawada, Takayuki; Kudo, Shinji; Kuriyama, Kazuhiko; Iwanami, Akira
 PATENT ASSIGNEE(S): Kyorin Pharmaceutical Co., Ltd., Japan
 SOURCE: Jpn. Kokai Tokkyo Koho, 23 pp.
 CODEN: JRXAAF
 DOCUMENT TYPE: Patent
 LANGUAGE: Japanese
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 2002145848	A2	20020522	JP 2000-343930	20001110
PRIORITY APPLN. INFO.:			JP 2000-343930	20001110

OTHER SOURCE(S): MARPAT 136:386402
 AB A[NR1CHR2CO]mNHCHR3CONHCHR4CH:CR5R6 [A = Z, Boc, RCO, R(CO)2, RS02; R = (un)substituted Ph, (un)substituted PhCH2, (un)substituted styryl, etc.; R1 = H; R1R2 may be linked to form pyrrolidine ring; R2-R4 = H, (un)substituted C1-4 alkyl, cyclohexylmethyl, (un)substituted PhCH2, etc.; R5 = H, F, C1-4 alkoxy, carbonyl; R6 = C1-4 alkoxy, carbonyl, CO2H, cyano, phenylsulfonyl, etc.; m = 0, 1], their pharmaco. acceptable salts, and their hydrates, useful as immunosuppressants, anti-inflammatory agents, antiallergy agents, anticancer agents, and nerve disorder-treating agents, are prepared by condensation of A[NR1CHR2CO]mNHCHR3CONHCHR4COH (A, R1-R4, m = same as above) with R7CHR8PO(OEt)2 (R7 = H, F; R8 = C1-4 alkoxy, carbonyl, CO2H, C1-4 alkoxyphosphoryl, cyano, etc.) or R9CH2CO2Ra (R9 = C1-4 alkoxy, carbonyl; Ra = C1-4 alkyl), followed by optional hydrolysis and further chemical modification. Thus, 150 mg MeSO2CH2PO(OEt)2 was treated with NaH in THF at room temperature for 1 h and condensed with 300 mg Z-L-Leu-L-Phe-L-Phe-H to give 89 mg Z-L-Leu-L-Phe-NH-L-CH(CH2Ph)CH:CHSO2Me, which inhibited proteasome with IC50 value of 0.14 µg/mL.

IT 428511-71-1P
 RL: PAC (Pharmacological activity); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USRS (Uses)
 (preparation of alkenylamino acids as proteasome inhibitors)
 RN 428511-71-1 CAPLUS
 CN 2-Quinolincarboxamide, N-[(1S)-2-[[[(1S)-3-(methylsulfonyl)-1-(phenylmethyl)-2-propenyl]amino]-2-oxo-1-(phenylmethyl)ethyl]- (9CI) (CA INDEX NAME)


Absolute stereochemistry.
 Double bond geometry unknown.

L4 ANSWER 69 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)



L4 ANSWER 70 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
 ACCESSION NUMBER: 2002:312355 CAPLUS
 DOCUMENT NUMBER: 137:63365
 TITLE: Relative and Absolute Stereochemistry of the Didemnaketals, Metabolites of a Palauan Ascidian, Didemnum sp.
 AUTHOR(S): Salomon, Christine E.; Williams, David H.; Lobkovsky, Emil; Clardy, Jon C.; Faulkner, D. John
 CORPORATE SOURCE: Scripps Institution of Oceanography, UCSD, La Jolla, CA, 92093-0212, USA
 SOURCE: Organic Letters (2002), 4(10), 1699-1702
 CODEN: ORLEF7; ISSN: 1523-7060
 PUBLISHER: American Chemical Society
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 OTHER SOURCE(S): CASREACT 137:63365
 AB The absolute stereochem. of the heptaprenoids didemnaketals B and C, isolated from a Palauan ascidian, was determined using a combination of degradation and derivatization expts., chiral shift methods, and comparison of fragments to known compds.
 IT 438462-64-79
 RL: SPN (Synthetic preparation); PREP (Preparation)
 (isolation of didemnaketal C from Didemnum sp. and determination of relative and absolute configuration of didemnaketals B and C)
 RN 438462-64-7 CAPLUS
 CN Benzeneacetic acid, α-[[[(2S,4R,6S,8S,10S)-8-[(1S,2S)-4-hydroxy-2-methyl-1-(3-methyl-1-oxobutoxy)butyl]-4,10-dimethyl-1,7-dioxaspiro[5.5]undec-2-yl]carbonylamino]-, methyl ester, (αS) (9CI) (CA INDEX NAME)

Absolute stereochemistry.



REFERENCE COUNT: 6 THERE ARE 6 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 71 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
 ACCESSION NUMBER: 2002:312037 CAPLUS
 DOCUMENT NUMBER: 136:325436
 TITLE: Preparation of quinolylindoles as antimicrobial agents
 INVENTOR(S): Cuny, Gregory D.; Hauske, James R.; Hoemann, Michael Z.; Chopra, Ian
 PATENT ASSIGNEE(S): Sepracor Inc., USA
 SOURCE: U.S., 167 pp., Cont. of U.S. Ser. No. 639,622.
 CODEN: USXXAM
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 7
 PATENT INFORMATION:


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 6376670	B1	20020423	US 2000-658690	20000908
US 6207679	B1	20010327	US 1998-45051	19980319
US 6172084	B1	20010109	US 1998-99640	19980618
US 6103905	A	20000815	US 1998-213385	19981211
PRIORITY APPLN. INFO.:			US 1997-878781	B2 19970619
			US 1998-45051	A2 19980319
			US 1998-99640	A2 19980618
			US 1998-213385	A1 19981211
			US 2000-639622	A2 20000815

OTHER SOURCE(S): MARPAT 136:325436
 GI

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

AB The title compds. [I, Z = CO, CR2; R = H, alkyl; R5-R8, R14-R17 = H, halo, alkyl, etc.; R9, R10 = H, alkyl, cycloalkyl, etc.; R3 = H, alkyl; R11 = H, alkyl; R12 = H, alkyl] which are bactericidal to a Gram-pos. bacterium via a non-lytic mechanism at its MIC (data given), were prepared E.g., a multi-step synthesis of II, was given.
 IT 275357-08-9P
 RL: PAC (Pharmacological activity); RCT (Reactant); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation); RACT (Reactant or reagent)
 (preparation of quinolylindole derivs. as antimicrobial agents)
 RN 275357-08-9 CAPLUS
 CN 4-Quinolincarboxamide, N-[2-amino-1-[(4-nitrophenyl)methyl]-2-oxoethyl]-2-(5-chloro-1H-indol-3-yl)- (9CI) (CA INDEX NAME)

L4 ANSWER 71 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)



REFERENCE COUNT: 42 THERE ARE 42 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 72 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN


ACCESSION NUMBER: 2002:293470 CAPLUS
 DOCUMENT NUMBER: 136:319354
 TITLE: Agents for the treatment of viral infections
 INVENTOR(S): Schubert, Ulrich; Will, Hans; Teasmer, Uwe; Sirna, Huesseyin; Prassolow, Alexij; Schubert, Evelyn; Hohenberg, Heinz
 PATENT ASSIGNER(S): Germany
 SOURCE: PCT Int. Appl., 117 pp.
 CODEN: PIXXD2
 DOCUMENT TYPE: Patent
 LANGUAGE: German
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2002030455	A2	20020418	WO 2001-DE3908	20011011
WO 2002030455	A3	20020808		
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM				
RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF, BJ, CP, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
DE 10051716	A1	20020425	DE 2000-10051716	20001012
DE 10149398	A1	20030424	DE 2001-10149398	20011003
AU 2002018133	A5	20020422	AU 2002-18133	20011011
EP 1326632	A2	20030716	EP 2001-986607	20011011
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR				
JP 2004510826	T2	20040408	JP 2002-533894	20011011
PRIORITY APPLN. INFO.: DE 2000-10051716 A 20001012 DE 2001-10149398 A 20011003 WO 2001-DE3908 W 20011011				
AB The invention relates to agents for the treatment of viral infections, in particular, infections with hepatitis and retroviruses. Said agents inhibit the release, maturation and replication of both retroviruses and also hepatitis viruses. In the example of human immune deficiency virus (HIV) and hepatitis-B viruses it has been shown that proteasome inhibitors block the release of virus particles and the infectiousness of the released viral particles and thus the reproduction of the viruses. The proteasome inhibitors affect the activities in the ubiquitin/proteasome pathway, in particular the enzymic activities of the 26S and the 20S proteasome complexes. The application for the above invention lies in antiretroviral therapy, particularly the treatment of HIV infections and AIDS and in the antiviral therapy of hepatitis infections, in particular the treatment of acute and chronic HBV and HCV infections and the associated liver carcinomas.				
IT 179324-59-5, PS 325 RI: PAC (Pharmacological activity); THU (Therapeutic use); BIOL (Biological study); USES (Uses) (proteasome-inhibitory agents for treatment of viral infections)				
RN 179324-59-5 CAPLUS				

L4 ANSWER 72 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)

CN Boronic acid, {[(1R)-3-methyl-1-[(2S)-1-oxo-4-phenyl-2-[(2-quinolinylcarbonyl)amino]butyl]amino]butyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




L4 ANSWER 73 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 2002:256223 CAPLUS
 DOCUMENT NUMBER: 136:295089
 TITLE: Preparation of amino acid aromatic derivatives with HIV integrase inhibitory properties
 INVENTOR(S): N'zamba, Blaise Magloire; Sauve, Gilles; Sevigny, Guy; Velle, Jocelyn
 PATENT ASSIGNER(S): Pharmacor, Inc., Can.
 SOURCE: PCT Int. Appl., 173 pp.
 CODEN: PIXXD2
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2002026697	A2	20020404	WO 2001-CA1367	20010925
WO 2002026697	A3	20020516		
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM				
RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF, BJ, CP, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
AU 2001095310	A5	20020408	AU 2001-95310	20010925
US 6528655	B1	20030304	US 2001-963329	20010926
PRIORITY APPLN. INFO.: CA 2000-2321348 A 20000927 WO 2001-CA1367 W 20010925				
OTHER SOURCE(S): MARPAT 136:295089				
AB Amino acid derivs. R1CO-A-CONHR2 [A = NR3CR4R5, where R3, R4 = H or Me; R5 = H, alkyl, carboxyalkyl, benzyl, MeSCH2CH2, 1-indolylmethyl, 3,4-(HO)2C6H2CH2, etc.; R3R4 may be trimethylene, which may be substituted; R1, R2 are certain rings (Ph, 3-pyridyl, 2-quinolyl, 2-thienyl, etc.), which may be substituted and attached to alkyl; R2 may also be acylamino] were prepared as inhibitors of HIV integrase. Thus, N-[N-(3,4-dihydroxybenzoyl)-N-trityl-L-histidinyl]dopamine was prepared by coupling of Na-(9-fluorenylmethoxycarbonyl)-N-trityl-L-histidine with dopamine hydrochloride, deprotection, and acylation with 3,4-dihydroxybenzoic acid and showed anti-integrase activity IC50 = 65 nM.				
IT 406728-01-6P RI: PAC (Pharmacological activity); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses) (preparation of amino acid aromatic derivs. with HIV integrase inhibitory properties)				
RN 406728-01-6 CAPLUS				
CN 2-Quinolincarboxamide, N-[(1S)-2-[[2-(3,4-dihydroxyphenyl)ethyl]amino]-1-[(4-hydroxyphenyl)methyl]-2-oxoethyl]-4,8-dihydroxy- (9CI) (CA INDEX NAME)				

Absolute stereochemistry.


L4 ANSWER 73 OF 261 CAPLUS COPYRIGHT 2004 ACS ON STN (Continued)



L4 ANSWER 74 OF 261 CAPLUS COPYRIGHT 2004 ACS ON STN

ACCESSION NUMBER: 2002:252956 CAPLUS
 DOCUMENT NUMBER: 136:273215
 TITLE: Combination of an NK-3 receptor antagonist and a CNS-penetrant NK-1 receptor antagonist for treating depression and anxiety
 INVENTOR(S): Lowe, John Adams, III; McLean, Stafford; Sobolov-Jaynes, Susan Beth
 PATENT ASSIGNEE(S): Pfizer Products Inc., USA
 SOURCE: Eur. Pat. Appl., 65 pp.
 CODEN: EPXXDW
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:



PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
EP 1192952	A2	20020403	EP 2001-307657	20010910
EP 1192952	A3	20030326		
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO				
BR 2001004345	A	20020521	BR 2001-4345	20010928
JP 2002338497	A2	20021127	JP 2001-300136	20010928
PRIORITY APPLN. INFO.: US 2000-236375P P 20000928				
OTHER SOURCE(S): MARPAT 136:273215				
AB A composition for the treatment of anxiety or depression in a mammal, including a human, comprises (a) an NK-3 receptor antagonist or its salt, (b) a CNS-penetrant NK-1 receptor antagonist or its salt, and (c) a pharmaceutically acceptable carrier. When administered in combination, either as a single or as sep. pharmaceutical composition(s), the CNS-penetrant NK-1 receptor antagonist and an NK-3 antagonist, are presented in a ratio which is consistent with the manifestation of the desired effect. In particular, the ratio by weight of the CNS-penetrant NK-1 receptor antagonist and the NK-3 antagonist will suitably be between 0.001:1 to 1000:1, and especially between 0.01:1 and 100:1.				
IT 174635-51-9				
RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses) (combination of NK3 receptor antagonist and CNS-penetrant NK1 receptor antagonist for treating depression and anxiety)				
RN 174635-51-9 CAPLUS				
CN Benzeneacetic acid, α -[[(2-phenyl-4-quinolinyl)carbonyl]amino]-, methyl ester (9CI) (CA INDEX NAME)				



L4 ANSWER 75 OF 261 CAPLUS COPYRIGHT 2004 ACS ON STN

ACCESSION NUMBER: 2002:240759 CAPLUS
 DOCUMENT NUMBER: 136:279469
 TITLE: Preparation of quinoline and quinoxaline derivatives as farnesyl transferase inhibitors for treatment of tumors and proliferative diseases
 INVENTOR(S): Angibaud, Patrick Rene; Venet, Marc Gaston; Pilatte, Isabelle Noelle Constance
 PATENT ASSIGNEE(S): Janssen Pharmaceutica N.V., Belg.
 SOURCE: PCT Int. Appl., 66 pp.
 CODEN: PIXXD2
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2002024682	A1	20020328	WO 2001-EP10867	20010918
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF, RJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
EP 1322635	A1	20030702	EP 2001-974271	20010918
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR				
JP 2004509883	T2	20040402	JP 2002-529092	20010918
AU 2001093826	A5	20020402	AU 2001-93826	20020402
US 2003203904	A1	20031030	US 2003-381363	20030324
PRIORITY APPLN. INFO.: EP 2000-203365 A 20000925				
WO 2001-EP10867 W 20010918				
OTHER SOURCE(S): MARPAT 136:279469				
GI				




L4 ANSWER 75 OF 261 CAPLUS COPYRIGHT 2004 ACS ON STN (Continued)

heterocycloxy, alkylthio, or (un)substituted (cyclo)alkyl, alkenyl, alkynyl, carbamoyl, amino, sulfamoyl, etc.; or R1R2 = OCH2O, OCH2CH2O, OCH:CH, OCH2CH2, OCH2CH2CH2, CH:CHCH:CH; R3 = H, halo, CN, alkenyl, alkynyl, hydroxycarbonyl, alkoxy, carbonyl, aryl, heterocyclyl, alkoxy, alkylthio, (un)substituted (cyclo)alkyl or amino, etc.; R4 = (un)substituted imidazolyl, triazolyl, or pyridyl; R5 = CN, OH, halo, alkenyl, alkynyl, hydroxycarbonyl, alkoxy, carbonyl, or (un)substituted (cyclo)alkyl, alkoxy, amino, or carbamoyl, etc.; R7 = halo or (un)substituted (cyclo)alkyl, alkenyl, alkynyl, alkylthio, carboxy, carbamoyl, acyl(amino), etc.; or pharmaceutically acceptable salts, N-oxides, or stereochem. isomeric forms thereof were prepd. For example, N-[2-(3-chlorobenzoyl)-4-(4-chlorobenzoyl)phenyl]acetamide was cyclized with NH₃ in i-PrOH to give (4-chlorophenyl)-4-(3-chlorophenyl)-2-methyl-6-quinazolinylmethanone (36f). Addn. of 1-methyl-1H-imidazole in the presence of BuLi and SiEt₃Cl in THF afforded II (40f). I have potent farnesyl transferase inhibitory effect and are useful for inhibiting proliferative diseases and growth of tumors expressing an activated ras oncogene (no data).

IT 405549-73-7P
 RL: PAC (Pharmacological activity); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)
 (farnesyl transferase inhibitor; preparation of quinoline and quinoxaline derivs. as farnesyl transferase inhibitors for treatment of tumors and proliferative diseases)
 RN 405549-73-7 CAPLUS
 CN Benzeneacetic acid, α -[[(4-(3-chlorophenyl)-6-[(4-chlorophenyl)(1-methyl-1H-imidazol-5-yl)methyl]-2-quinolinyl)carbonyl]amino]-, methyl ester, (4S) (9CI) (CA INDEX NAME)

Absolute stereochemistry.




REFERENCE COUNT: 5 THERE ARE 5 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

AB Title compds. I [wherein m and n = independently 0-5; q = 0-3; Y1Y2 = C:N or C:CR9; C9 = H, halo, CN, (cyclo)alkyl, hydroxyalkyl, alkoxy(alkyl), aminoalkyl, (amino)alkenyl, (amino)alkynyl, halocarbonyl, hydroxycarbonyl, alkoxy, carbonyl, aryl, (un)substituted amino or carbamoyl, etc.; R1 and R2= independently azido, OH, halo, CN, NO₂, trihalomethyl, alkoxy, aryloxy,

L4 ANSWER 76 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
 ACCESSION NUMBER: 2002:209379 CAPLUS
 DOCUMENT NUMBER: 137:33156
 TITLE: Asymmetric construction of the azaspiro[5.6]dodec-9-ene system in marine natural toxins
 AUTHOR(S): Ishihara, Jun; Horie, Mariko; Shimada, Yoshikatsu; Tojo, Shingo; Murai, Akio
 CORPORATE SOURCE: Division of Chemistry, Graduate School of Science, Hokkaido University, Sapporo, 060-0810, Japan
 SOURCE: Synlett (2002), (3), 403-406
 CODEN: SYNLSS; ISSN: 0936-5214
 PUBLISHER: Georg Thieme Verlag
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 OTHER SOURCE(S): CASREACT 137:33156
 AB The asym. formation of azaspiro[5.6]dodec-9-ene system is described. The Diels-Alder reactions of an α -methylene caprolactam and diene in the presence of a Cu(II) and (S,S)-tert-Bu-BOX complex afford the desired spirocyclic compds. with good exo-selectivity as well as excellent enantioselectivity. These exo-adducts would be applicable to the synthesis of marine natural toxins including the corresponding cyclic imine moiety.
 IT 436153-34-3P
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)
 (asym. synthesis of azaspiro[5.6]dodec-9-ene system in marine natural toxins via copper-catalyzed stereoselective Diels-Alder reaction)
 RN 436153-34-3 CAPLUS
 CN 8-Azaspiro[5.6]dodec-2-ene-8-carboxylic acid, 1-[[[(1R)-2-methoxy-2-oxo-1-phenylethyl]amino]carbonyl]-3-methyl-7-oxo-, phenylmethyl ester, (1S,6S)-(9CI) (CA INDEX NAME)

Absolute stereochemistry.




REFERENCE COUNT: 31 THERE ARE 31 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 77 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
 ACCESSION NUMBER: 2002:209378 CAPLUS
 DOCUMENT NUMBER: 137:78803
 TITLE: Asymmetric construction of the azaspiro[5.5]undec-8-ene system towards gymnodimine synthesis
 AUTHOR(S): Tsujimoto, Takashi; Ishihara, Jun; Horie, Mariko; Murai, Akio
 CORPORATE SOURCE: Division of Chemistry, Graduate School of Science, Hokkaido University, Sapporo, 060-0810, Japan
 SOURCE: Synlett (2002), (3), 395-402
 CODEN: SYNLSS; ISSN: 0936-5214
 PUBLISHER: Georg Thieme Verlag
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 OTHER SOURCE(S): CASREACT 137:78803
 GI

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

AB In the course of the synthetic studies on gymnodimine (I), a potent shellfish toxin, the asym. construction of the azaspirocyclic part II was achieved by the (-)-siam-Cu(SbF₆)₂ complex catalyzed intermol. Diels-Alder reaction in high exo- and enantioselectivities.
 IT 440677-28-1P
 RL: SPN (Synthetic preparation); PREP (Preparation)
 (asym. construction of the azaspiro[5.5]undec-8-ene system towards gymnodimine synthesis)
 RN 440677-28-1 CAPLUS
 CN 2-Azaspiro[5.5]undec-8-ene-2-carboxylic acid, 7-[[[(1S)-2-methoxy-2-oxo-1-phenylethyl]amino]carbonyl]-9-[[[(4-methoxyphenyl)methoxy]methyl]-8-methyl-1-oxo-, phenylmethyl ester, (6R,7S)-(9CI) (CA INDEX NAME)

Absolute stereochemistry.




REFERENCE COUNT: 25 THERE ARE 25 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 78 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
 ACCESSION NUMBER: 2002:171694 CAPLUS
 DOCUMENT NUMBER: 136:232208
 TITLE: Preparation of 4-aminoazepan-3-one parasitic cysteine protease inhibitors effective against malaria and other diseases
 INVENTOR(S): Tew, David G.; Thompson, Scott K.; Veber, Daniel F.
 PATENT ASSIGNEE(S): Smithkline Beecham Corporation, UK
 SOURCE: PCT Int. Appl., 220 pp.
 CODEN: PIXXD2
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 4
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2002017924	A1	20020307	WO 2001-US27178	20010831
W:	AS, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM			
RW:	GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GO, GW, ML, MR, NE, SN, TD, TG			
US 2003144175	A1	20030731	US 2001-881334	20010614
AU 2001086983	A5	20020313	AU 2001-86593	20010831
EP 1320370	A1	20030625	EP 2001-966474	20010831
R:	AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR			
JP 2004509083	T2	20040325	JP 2002-522897	20010831
PRIORITY APPLN. INFO.:			US 2000-653815	A2 20000901
			US 2001-881334	A2 20010614
			US 1998-113636P	P 19981223
			US 1999-164581P	P 19991110
			WO 1999-US30730	A2 19991221
			US 2000-593845	B2 20000614
			WO 2001-US27178	W 20010831

OTHER SOURCE(S): MARPAT 136:232208
 GI




AB The present invention relates to methods of treating parasitic diseases which are mediated by cysteine proteases by administration of 4-aminoazepan-3-one protease inhibitors I (e.g. benzo[1,3]dioxole-5-carboxylic acid [(S)-1-(1-benzyl-3-oxoazepan-4-yl)carbamoyl]-3-

L4 ANSWER 78 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)
 methylbutylamide) and pharmaceutically acceptable salts, hydrates and solvates thereof. In particular, the present invention relates to a method of treating malaria by inhibiting the cysteine protease falcipain. Other diseases against which the claimed compds. are effective include trypanosomiasis (African sleeping sickness, Chagas disease), leishmaniasis, schistosomiasis, onchocerciasis (river blindness) and giardiasis. In I: R1 is R4NR'CHR3C(O)-, RSXCHR3C(O)-, R3CH2C(O)-, R4NR'CHR3C(O)-, II. R2 is H, C1-6alkyl, C3-6cycloalkyl-CO-6alkyl, Ar-CO-6alkyl, Het-CO-6alkyl, R9C(O)-, R9C(S)-, R9SO2-, R9OC(O)-, R9R11NC(O)-, R9R11NC(S)-, R9(R11)NSO2-, 3-(2-pyridyl)benzylcarbonyl, 2-(3-(2-pyridyl)phenyl)ethyl, R7NR6CHR8Z-, and R9SO2R11NC(O)-. R3 is H, C1-6alkyl, C3-6cycloalkyl-CO-6alkyl, C2-6alkenyl, C2-6alkynyl, HetCO-6alkyl and ArCO-6alkyl. R3 and R' may be connected to form a pyrrolidine, piperidine or morpholine ring. R4 is H, C1-6alkyl, C3-6cycloalkyl-CO-6alkyl, Ar-CO-6alkyl, Het-CO-6alkyl, R5C(O)-, R5C(S)-, R5SO2-, R5OC(O)-, R5R12NC(O)-, and R5R12NC(S)-. R5 is H, C1-6alkyl, C2-6alkenyl, C2-6alkynyl, C3-6cycloalkyl-CO-6alkyl, Ar-CO-6alkyl and Het-CO-6alkyl. R6 is H, C1-6alkyl, Ar-CO-6alkyl, and Het-CO-6alkyl. R7 is H, C1-6alkyl, C3-6cycloalkyl-CO-6alkyl, Ar-CO-6alkyl, Het-CO-6alkyl, R10C(O)-, R10C(S)-, R10SO2-, R10OC(O)-, R10R13NC(O)-, and R10R13NC(S)-. R8 is H, C1-6alkyl, C2-6alkenyl, C2-6alkynyl, HetCO-6alkyl and ArCO-6alkyl. R9, R10 independently = C1-6alkyl, C3-6cycloalkyl-CO-6alkyl, Ar-CO-6alkyl and Het-CO-6alkyl. R11, R12, R13, R', R'' independently = H, C1-6alkyl, Ar-CO-6alkyl, and Het-CO-6alkyl. R''' is H, C1-6alkyl, C3-6cycloalkyl-CO-6alkyl, C2-6alkenyl, and Het-CO-6alkyl. R'''' is C1-6alkyl, C3-6cycloalkyl-CO-6alkyl, C2-6alkenyl, C2-6alkynyl, HetCO-6alkyl and ArCO-6alkyl. X is CH2, S, and O; Z is C(O) and CH2; n is 1-5. Although the methods of prepn. are not claimed, 220 example prepn. are included.
 IT 350796-41-7P, Quinoline-8-carboxylic acid [(1S)-1-[[[(4S)-3-oxo-1-(pyridin-2-yl)sulfonyl]azepan-4-yl]carbamoyl]-2-phenylethyl]amide
 RL: PAC (Pharmacological activity); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)
 (preparation of 4-aminoazepan-3-one parasitic cysteine protease inhibitors effective against malaria and other diseases)

RN 350796-41-7 CAPLUS
 CN 8-Quinolincarboxamide, N-[(1S)-2-[[[(4S)-hexahydro-3-oxo-1-(2-pyridinyl)sulfonyl]-1H-azepin-4-yl]amino]-2-oxo-1-(phenylmethyl)ethyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




REFERENCE COUNT: 2 THERE ARE 2 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 79 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
 ACCESSION NUMBER: 2002:142667 CAPLUS
 DOCUMENT NUMBER: 136:200103
 TITLE: Preparation of (thio)urea moiety-containing heterocyclic compounds as VIA-4 antagonists
 INVENTOR(S): Fukui, Hideto; Ikegami, Satoru; Okuyama, Akihiko
 PATENT ASSIGNER(S): Kaken Pharmaceutical Co., Ltd., Japan
 SOURCE: PCT Int. Appl., 43 pp.
 CODEN: PIXXD2
 DOCUMENT TYPE: Patent
 LANGUAGE: Japanese
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2002014272	A1	20020221	WO 2001-JP6833	20010808
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM			
RW:	GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SV, TD, TG			
AU 2001077720	A5	20020225	AU 2001-77720	20010808
PRIORITY APPLN. INFO.:			JP 2000-241657	A 20000809
			WO 2001-JP6833	W 20010808
OTHER SOURCE(S):			MARPAT 136:200103	
GI				

L4 ANSWER 79 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)
 RL: PAC (Pharmacological activity); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)
 (prepn. of (thio)urea moiety-contg. heterocyclic compds. as VIA-4 antagonists)
 RN 401470-80-2 CAPLUS
 CN L-Phenylalanine, 4-[[[(3,5-dichloro-4-pyridinyl)carbonyl]amino]-N-[(3,4-dihydro-1(2H)-quinolinyl)carbonyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




REFERENCE COUNT: 6 THERE ARE 6 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT




AB The title compds. I [R1 = H, alkyl, etc.; X1 = single bond, C.tplbond.C, etc.; Y = O, etc.; Z = NR7R8, etc.; R7, R8 = H, hydrocarbon, etc.; X2 = heterocyclic ring (generic structure given); further details on said heterocyclic ring are given] are prepared. A process for the preparation of I is claimed. In an assay for inhibition of VIA-4/VCAM-1 adhesion, 3-[4-[(3,5-dichloropyridine-4-carbonyl)amino]phenyl]-2-(S)-[3-isobutyl-3-[(1S)-phenylethyl]ureido]propionic acid showed IC50 of 1.1 nM.
 IT 401470-80-2P

L4 ANSWER 80 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
 ACCESSION NUMBER: 2002:116963 CAPLUS
 DOCUMENT NUMBER: 137:163311
 TITLE: A new class of type I protein geranylgeranyltransferase (GGTase I) inhibitor
 AUTHOR(S): Sunami, Satoshi; Ohkubo, Mitsuru; Sagara, Takeshi; Ono, Jun; Asahi, Shuichi; Koito, Seita; Morishima, Hajime
 CORPORATE SOURCE: Banyu Tsukuba Research Institute, Ibaraki, Tsukuba, 300-2611, Japan
 SOURCE: Bioorganic & Medicinal Chemistry Letters (2002), 12(4), 629-632
 CODEN: BMCLEB; ISSN: 0960-894X
 PUBLISHER: Elsevier Science Ltd.
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 GI

L4 ANSWER 80 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)



REFERENCE COUNT: 22 THERE ARE 22 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT



AB Replacement of the thiol groups in I, a potent and highly selective Candida albicans GGTase I inhibitor discovered through screening, with an imidazole ring was achieved by using solid phase synthesis. A non-thiol compound was found as a representative of a new class of potent C. albicans GGTase I inhibitor with high selectivity against human GGTase I. The relation of these results to the possible antifungal activity of these compds. is discussed.



IT 445400-99-7P
 RL: CPN (Combinatorial preparation); PAC (Pharmacological activity); THU (Therapeutic use); BIOL (Biological study); CMRI (Combinatorial study); PREP (Preparation); USES (Uses)
 (new class of type I protein geranylgeranyltransferase (GGTase I) inhibitor in relation to structure and antifungal activity)
 RN 445400-99-7 CAPLUS
 CN 9H-Xanthene-9-carboxamide, N-[(1R)-1-[(3,4-dichlorophenyl)methyl]-2-[[2-(1H-imidazol-4-yl)ethyl]amino]-2-oxoethyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

L4 ANSWER 81 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
 ACCESSION NUMBER: 2002:107157 CAPLUS
 DOCUMENT NUMBER: 136:167388
 TITLE: Preparation and use of quinolone and naphthyridine derivatives as inhibitors of cellular efflux pumps of microbes
 INVENTOR(S): De Souza, Noel J.; Patel, Mahesh V.; Gupta, Shrikant V.; Upadhyay, Dilip J.; Shukla, Milind C.; Chaturvedi, Nishith C.; Bhawsar, Satish B.; Nair, Sheela C.; Jafri, Mohammed A.; Khorakiwala, Mabil F.
 PATENT ASSIGNER(S): Wockhardt Limited, India
 SOURCE: PCT Int. Appl., 149 pp.
 CODEN: PIXXD2
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 7
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2002009758	A2	20020207	WO 2001-1N139	20010731
WO 2002009758	A3	20021227		
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MY, NZ, OL, OM, OS, PA, PE, PG, PH, PI, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM			
RW:	GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LJ, MC, NL, PT, SE, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG			
US 2002165227	A1	20021107	US 2001-850669	20010507
US 6608078	B2	20030819		
AU 2001080091	A5	20020213	AU 2001-80091	20010731
US 2002177559	A1	20021128	US 2001-919347	20010731
EP 1305048	A2	20030502	EP 2001-958373	20010731
R:	AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR			
US 2003144517	A1	20030731	US 2002-303692	20021122
PRIORITY APPLN. INFO.:			US 2000-222201P	P 20000801
			US 2000-640947	A 20000819
			WO 2000-1N111	W 20001121
			US 2001-286291P	P 20010425
			US 2001-850669	A 20010507
			WO 2001-1N100	A 20010508
			US 1999-170676P	P 19991214
			US 2000-202459P	P 20000508
			US 2000-566875	A2 20000508
			US 2001-802793	A3 20010309
			WO 2001-1N139	W 20010731
OTHER SOURCE(S):	MARPAT 136:167388			
GI				

L4 ANSWER 81 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)




AB Title compds. I [R1 = H, (cyclo)alkyl, aryl, aralkyl, arylaminoalkyl, aryloxyalkyl, arylSOO-2 alkyl or when X = C and the nitrogen atom to which R1 is linked forms an (un)substituted 4-7 membered ring with X of the adjacent ring, the ring optionally containing one or more hetero atoms selected from N, O, S, said heteroatom(s) represented by Y; R2 = H, CHO, COOR3, CONHR13, where R13 = H or the NHR13 of CONHR13 is the residue of an amino acid; R3 = H, alkyl, cycloalkyl, aryl, aralkyl, arylaminoalkyl, aryloxyalkyl, arylSOO-2 alkyl, O-carboxy, etc.; R4 = H; R4' = H or R4 and R4' taken together are -O-, -S-, R5 = H, alkyl, amino, alkylamino, acylamino, R6 = H, alkyl, halo, amino, hydroxy, R7 = OH, halo, NR9R10, etc.; R9-10 = H, alkyl, (CH2)nOAc or R9 = H and R10 = 4-7 membered carbocyclic, heterocyclic ring linked to the nitrogen of NR9R10 through an atom of the heterocycle other than the heterocyclic atom, etc.; A = H, alkyl, glycosyl, aralkyl, alkanoyl, aminoalkanoyl wherein the aminoalkanoyl group may be an amino acid residue or A is C6H11O6, SO3H, PO3H2; X = CH, CF, CCl, CCH3, CCF3, COCH3, COCHF2, C-OCF3, N or when X is equal to C it forms together with the nitrogen atom of the adjacent ring an (un)substituted 5-7 membered ring containing carbon atoms and optionally Y atoms representing one or more N, O, S] were prepared. For instance, a mixture of 1-ethyl-6,7-difluoro-1,4-dihydro-4-oxoquinolone-3-carboxylic acid and 1,2,3,4-tetrahydroisoquinoline (DMSO, Et3N 140°C, 24 h) provided, after work-up and trituration II as a solid (62% yield), m.p. 220°C. II with ciprofloxacin had a fractional inhibitory concentration (FIC) index of 0.314 observed against S. aureus 1199 B (Nor A+). I are effective at inhibiting efflux pumps, e.g., MefA, MefE, Bmr, PmrA, etc. 396132-84-6P


IT RL: PAC (Pharmacological activity); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)
 (drug; preparation and use of quinolone and naphthyridine deriva. as inhibitors of cellular efflux pumps of microbes)

L4 ANSWER 81 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)
 RN 396132-84-6 CAPLUS
 CN L-Lysine, N-[[[(5R)-8,9-difluoro-6,7-dihydro-5-methyl-1-oxo-1H,5H-benzo[i,j]quinolinizin-2-yl]carbonyl]-L-phenylalanyl-, methyl ester (9CI) (CA INDEX NAME)

Absolute stereochemistry. Rotation (+).



L4 ANSWER 82 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
 ACCESSION NUMBER: 2002:73778 CAPLUS
 DOCUMENT NUMBER: 136:279671
 TITLE: A CD Exciton Chirality Method for Determination of the Absolute Configuration of threo-β-Aryl-β-hydroxy-α-amino Acid Derivatives
 AUTHOR(S): Lo, Lee-Chiang; Yang, Chun-Tzu; Tsai, Chang-Sheng
 CORPORATE SOURCE: Department of Chemistry, National Taiwan University, Taipei, 106, Taiwan
 SOURCE: Journal of Organic Chemistry (2002), 67(4), 1368-1371
 CODEN: JOCEAH; ISSN: 0022-3263
 PUBLISHER: American Chemical Society
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 OTHER SOURCE(S): CASREACT 136:279671
 GI




AB Threo-β-aryl-β-hydroxy-α-amino acids I (Ar = Ph, C6H4CN-4, C6H4NO2-3, C6H4NO2-2, C6H4Cl-3, C6H4OMe-2, 2,5-dimethoxyphenyl, 2,4-dimethylphenyl) were prepared and their absolute configuration was studied in CH2Cl2 by CD exciton chirality method using 7-diethylaminocoumarin-3-carboxylate as a red-shifted chromophore. By combining the data of CD and NMR coupling const., the authors were able to correlate the preferred conformer (B) and the pos. CD to the corresponding (2S,3R)-absolute configuration. These results are consistent with those obtained from serine and threonine deriva., which represent the simplest form of β-hydroxy-α-amino acids. Thus, this CD method could become a general method for determining the absolute configuration of threo-β-aryl-β-hydroxy-α-amino acids.

IT 405510-93-2P
 RL: BPN (Biosynthetic preparation); PRP (Properties); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation)
 (CD exciton chirality method for determining absolute configuration of chromophore-containing β-aryl-β-hydroxy amino acid esters)

RN 405510-93-2 CAPLUS
 CN 2H-1-Benzopyran-3-carboxylic acid, 7-(diethylamino)-2-oxo-, (1R,2S)-2-[[[7-(diethylamino)-2-oxo-2H-1-benzopyran-3-yl]carbonyl]amino]-3-methoxy-3-oxo-1-phenylpropyl ester (9CI) (CA INDEX NAME)

Absolute stereochemistry.

L4 ANSWER 82 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)



REFERENCE COUNT: 31 THERE ARE 31 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 83 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 2002:46807 CAPLUS
DOCUMENT NUMBER: 137:185793
TITLE: Constrained segment ligation
AUTHOR(S): Tam, James P.; Miao, Zhenwei
CORPORATE SOURCE: Department of Microbiology and Immunology, Vanderbilt University, Nashville, TN, 37232-2363, USA
SOURCE: Innovation and Perspectives in Solid Phase Synthesis & Combinatorial Libraries: Peptides, Proteins and Nucleic Acids--Small Molecule Organic Chemistry Diversity, Collected Papers, International Symposium, 6th, York, United Kingdom, Aug. 31-Sept. 4, 1999 (2001), Meeting Date 1999, 11-14. Editor(s): Epton, Roger. Mayflower Scientific Ltd.: Kingswinford, UK. CODEN: 69CEGV; ISBN: 0-9515735-3-5

DOCUMENT TYPE: Conference
LANGUAGE: English

AB A symposium report. Imine ligation of Leu-Ile-Leu-Asn-Gly-OCH₂CHO with N-terminal (NT) amino acids in pyridine-acetic acid mixts. is highly regioselective. Oxaprolines from NT-Ser and NT-Thr as well as thiaprolines from NT-Cys are useful pseudoprolines (YPro). Differences in ligation rates are useful for an orthogonal tandem ligation strategy to couple multiple unprotected peptide segments without a protection scheme. Model dipeptides were used to study the stereochem. of the YPro derived from imine ligation.

IT 451524-16-6P


RL: SPN (Synthetic preparation); PREP (Preparation)
(ligation of peptide glycoaldehyde esters with N-terminal peptides)

RN 451524-16-6 CAPLUS

CN L-Isoleucinamide, L-leucyl-L-leucyl-L-leucyl-L-asparaginyglycyl-(3S)-2,3,4,9-tetrahydro-1-(hydroxymethyl)-1H-pyrido[3,4-b]indole-3-carbonyl-L-phenylalanyl-L-lysyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

L4 ANSWER 83 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)




REFERENCE COUNT: 10 THERE ARE 10 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 84 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 2002:10473 CAPLUS
DOCUMENT NUMBER: 136:69824
TITLE: Preparation of heterocycle compounds as melanocortin receptor ligands
INVENTOR(S): Carpino, Philip Albert; Cole, Bridget McCarthy; Morgan, Bradley Paul
PATENT ASSIGNEE(S): Pfizer Products Inc., USA
SOURCE: PCT Int. Appl., 108 pp.
CODEN: PIXXD2
DOCUMENT TYPE: Patent
LANGUAGE: English
FAMILY ACC. NUM. COUNT: 1
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2002000654	A1	20020103	WO 2001-1B995	20010531
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MY, NZ, NO, NZ, PL, PT, RO, RU, SD, SE, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM			
RW:	GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG			
EP 1294719	A1	20030326	EP 2001-934254	20010531
R:	AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR			
BR 2001011567	A	20030506	BR 2001-11567	20010531
JP 2004501917	T2	20040122	JP 2002-505778	20010531
US 2002072604	A1	20020613	US 2001-891026	20010625
BG 107268	A	20030630	BG 2002-107268	20021112
NO 2002006280	A	20021230	NO 2002-6280	20021230
PRIORITY APPLN. INFO.:			US 2000-214616P	P 20000628
			WO 2001-1B995	W 20010531

OTHER SOURCE(S): MARPAT 136:69824
GI




II

AB Compds. represented by formula HET-COC(R3R4)-NX4-CO(CR6R7)m-D [I; wherein m = 0, 1 or 2; HET = heterocyclyl; R3, R4 = H, C1-8 alkyl, CH(R8)-aryl, -CH(R8)-heteroaryl, -C(=O)-3 alkyl-C3-8 cycloalkyl (wherein the aryl or

L4 ANSWER 84 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)
 heteroaryl groups are optionally substituted by one or two groups; R8 = H, C1-8 alkyl, -CO-3 alkylaryl, -CO-3 alkylheteroaryl, -C3-6 cycloalkyl; R6, R7 = H, C1-6 alkyl, -CO-3 alkyl-aryl, -CO-3 alkyl-heteroaryl, -CO-3 alkyl-C3-8 cycloalkyl; or R6 and R7 together with the nitrogen atom to which they are attached form a 5- or 6-membered ring optionally contg. an addnl. heteroatom selected from O, S, NR3; D = -CO-6 alkylamino-C(=NR7)-NR15R16, -CO-6 alkylaminopyridyl, -CO-6 alkylaminoimidazolyl, -CO-6 alkylaminothiazolyl, -CO-6 alkylaminopyrimidinyl, -CO-6 alkylaminopiperazinyl-R15, -CO-6 alkylmorpholinyl, etc. (wherein R15, R16 = H, -C1-6 alkyl, -CO-3 alkylaryl, -CO-3 alkylheteroaryl, or -CO-3 alkyl-C3-8 cycloalkyl, wherein the alkyl and aryl groups are optionally substituted with one or two groups); X4 = H or C1-6 alkyl or X4 is taken together with R4 and the nitrogen atom to which X4 is attached and the carbon atom to which R4 is attached and form a five to seven membered ring) are prepd. Melanocortins are peptides derived from pro-opiomelanocortins (POMC) that bind to and activate G protein coupled receptors (GPCR's) of the melanocortin receptor family and regulate a diverse no. of physiol. processes including food intake, metab., and thermogenesis as well as sexual dysfunction. These compds. I are useful for the treatment or prevention of disorders, diseases, or conditions responsive to the activation of melanocortin receptor including obesity, diabetes mellitus, male or female sexual dysfunction, erectile dysfunction, or disorders that cause redbn. in appetite, or feeding behavior and/or body wt.; for modulating appetite and metabolic rates; for acutely stimulating the appetite for the treatment of hepatic lipidosis, cachexia, and other pathologies resulting in/from inappropriate food intake and wt. loss; for acutely stimulating the appetite of livestock for the treatment of ketosis, postpartum anestrus, and other metabolic and reproductive pathologies resulting in/from inappropriate food intake and wt. loss; and for enhancing growth and survivability of neonates in livestock. Thus, esterification of N-Boc-L-Tic-OH with N-hydroxysuccinimide using Et3N and EDC in CH2Cl2 at room temp. for 4 h gave 3,4-dihydro-1H-isoquinoline-2,3-(S)-dicarboxylic acid 2-tert-Bu ester 3-(2,5-dioxopyrrolidin-1-yl) ester which was condensed with D-p-chlorophenylalanine in the presence of Et3N in CH2Cl2 at room temp. overnight to give 3-(S)-[(R)-1-Carboxy-2-(4-chlorophenyl)ethylcarbamoyl]-3,4-dihydro-1H-isoquinoline-2-carboxylic acid tert-Bu ester. The latter compd. was further condensed with 8a-(Pyridin-2-ylmethyl)-2-(2,2,2-trifluoroethyl)tetrahydroimidazo[1,5-a]pyrazine-1,3-dione using Et3N and EDC in CH2Cl2 at 0° for 5 h to give (S)-3-[(R)-1-(4-chlorobenzyl)-2-(1,3-dioxo-8a-(pyridin-2-ylmethyl)-2-(2,2,2-trifluoroethyl)hexahydroimidazo[1,5-a]pyrazin-7-yl)-2-oxoethylcarbamoyl]-3,4-dihydro-1H-isoquinoline-2-carboxylic acid tert-Bu ester which was treated with a mixt. of EtOH and concd. HCl at 0° for 0.5 h to give (S)-1,2,3,4-Tetrahydroisoquinoline-3-carboxylic acid N-[(R)-1-(4-chlorobenzyl)-2-(1,3-dioxo-8a-(pyridin-2-ylmethyl)-2-(2,2,2-trifluoroethyl)hexahydroimidazo[1,5-a]pyrazin-7-yl)-2-oxoethylamide (II) hydrochloride which may be considered as a dipeptide analog heptacyclic amide, N-[N-(1,2,3,4-Tetrahydroisoquinoline-3-carboxyl)-D-p-chlorophenylalanyl]-1,3-dioxo-8a-(pyridin-2-ylmethyl)-2-(2,2,2-trifluoroethyl)hexahydroimidazo[1,5-a]pyrazine.
 IT 252008-71-2P
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)
 (intermediate; preparation of heterocycle compds. as melanocortin receptor ligands and therapeutic agents for treatment of prevention of obesity.

L4 ANSWER 84 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)
 diabetes mellitus, male or female sexual dysfunction)
 RN 252008-71-2 CAPLUS
 CN 2(1H)-Isoquinolinecarboxylic acid, 3-[[[(1R)-1-carboxy-2-(4-chlorophenyl)ethyl]amino]carbonyl]-3,4-dihydro-, 2-(1,1-dimethylethyl) ester, (3R)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




REFERENCE COUNT: 7 THERE ARE 7 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 85 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
 ACCESSION NUMBER: 2001:923616 CAPLUS
 DOCUMENT NUMBER: 136:93691
 TITLE: Preparation of 4-amino-azepan-3-one protease inhibitors
 INVENTOR(S): Marquis, Robert W., Jr.; Ru, Yu; Veber, Daniel F.; Cummings, Maxwell D.; Thompson, Scott K.; Yamashita, Dennis
 PATENT ASSIGNEE(S): Smithkline Beecham Corporation, USA
 SOURCE: PCT Int. Appl., 322 pp.
 CODEN: PIXXD2
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 4
 PATENT INFORMATION:



PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2001095911	A1	20011220	WO 2001-US19062	20010614
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM				
RW: GH, GM, KE, LS, MM, MZ, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SH, TD, TG				
EP 1307204	A1	20030507	EP 2001-946344	20010614
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR				
JP 2004503502	T2	20040205	JP 2002-510089	20010614
BG 107327	A	20030731	BG 2002-107327	20021128
NO 2002005786	A	20030212	NO 2002-5786	20021202
PRIORITY APPLN. INFO.: US 2000-593845 A2 20000614 NO 2001-US19062 W 20010614				
OTHER SOURCE(S): MARPAT 136:93691				
GI				

L4 ANSWER 85 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)
 H, alkyl, arylalkyl, etc.; R11 = H, alkyl, arylalkyl, etc.; R12 = H, alkyl, cycloalkyl, etc.; R13 = H, alkyl, alkenyl, etc.; R15 = H, alkyl, alkenyl, etc.] which inhibit proteases (no data), including cathepsin K, and are useful for treating diseases of excessive bone loss or cartilage or matrix degradn. including osteoporosis, gingival disease including gingivitis and periodontitis, arthritis, more specifically, osteoarthritis and rheumatoid arthritis, Paget's disease, hypercalcemia of malignancy, and metabolic bone disease, were prepd. E.g., a multi-step synthesis of compd. II was given.
 IT 281217-12-7P
 RL: PAC (Pharmacological activity); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)
 (preparation of 4-amino-azepan-3-one protease inhibitors)
 RN 281217-12-7 CAPLUS
 CN 8-Quinolinesulfonyl-1H-azepin-4-yl]amino]-2-oxo-1-(phenylmethyl)ethyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




REFERENCE COUNT: 2 THERE ARE 2 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT



AB The title compds. [I; R1 = COCR13NR11R12, COCR13XR15, COCH2R13; R2 = H, alkyl, cycloalkylalkyl, etc.; R3 = H, alkyl, cycloalkylalkyl, etc.; R4 =

L4 ANSWER 86 OF 261 CAPLUS COPYRIGHT 2004 ACS ON STN
 ACCESSION NUMBER: 2001:868423 CAPLUS
 DOCUMENT NUMBER: 136:5923
 TITLE: Preparation of sulfonamide/carbamide derivatives of 6(5H)phenanthridinones and their use as poly(ADP-ribose) polymerase (PARP) inhibitors
 INVENTOR(S): Li, Jia-He; Kalish, Vincent J.; Zhang, Jie; Serdyuk, Larisa E.; Ferraris, Dana V.; Xiao, Ge; Kletzly, Paul W.
 PATENT ASSIGNEE(S): Guilford Pharmaceuticals, Inc., USA
 SOURCE: PCT Int. Appl., 92 pp.
 CODEN: PIXXD2
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2001090077	A1	20011129	WO 2001-US15571	20010515
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, AM, AZ, BY, BG, BR, CA, CH, CN, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG			
US 2002006927	A1	20020117	US 2001-85455	20010515
PRIORITY APPLN. INFO.:			US 2000-205259P	P 20000519
OTHER SOURCE(S):		MARPAT 136:5923		



AB Title compds. I [R1 = H, halo; R2 = H, OH, NH2, NO, Me, aminomethyl, COOH; one of R3 and R4 = QP and the other of R3 and R4 is one of H, Me, CF3, NO2, amino, halo, piperazinyl, Q = A:O-X(Y), X(Y):A:O; P = Z-amino, Z, (hetero)cycle; A = C, S; O; X = O, S, N, N-substituted amino acid provided that when X = O, S, Y = absent and when X = N, Y = H, alkyl, alkoxy, alkylamino, or Y, Z taken together to form a 5 - 7 membered heterocycle; Z = H, bond, C=O, (cyclo)alkyl, carboxy, etc.] were prepared. Examples include

L4 ANSWER 87 OF 261 CAPLUS COPYRIGHT 2004 ACS ON STN
 ACCESSION NUMBER: 2001:780691 CAPLUS
 DOCUMENT NUMBER: 135:327371
 TITLE: 4-Amino-azepan-3-one inhibitors of cathepsin L, their preparation, and their therapeutic use
 INVENTOR(S): Cummings, Maxwell D.; Marquis, Robert W., Jr.; Ru, Yu; Thompson, Scott K.; Veber, Daniel F.; Yamashita, Dennis S.
 PATENT ASSIGNEE(S): Smithkline Beecham Corporation, USA
 SOURCE: PCT Int. Appl., 38 pp.
 CODEN: PIXXD2
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2001078734	A1	20011025	WO 2001-US12386	20010417
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, AM, AZ, BY, BG, BR, CA, CH, CN, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG			
EP 1303281	A1	20030423	EP 2001-927076	20010417
R:	AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR			
JP 2003531122	T2	20031021	JP 2001-576034	20010417
US 2004034013	A1	20040219	US 2002-258412	20021017
PRIORITY APPLN. INFO.:			US 2000-197717P	P 20000418
			WO 2001-US12386	W 20010417

OTHER SOURCE(S): MARPAT 135:327371
 AB Methods are provided which use 4-amino-azepan-3-one protease inhibitors of cathepsin L in the treatment of diseases in which cathepsin L is implicated, especially treatment or prevention of rheumatoid arthritis; cancer metastasis; diseases requiring inhibition of tissue destruction by macrophages, particularly lung macrophages, such as asthma, chronic obstructive pulmonary disease (COPD), and emphysema; and diseases requiring, for therapy, inhibition of pos. selection of CD4+T- cells by cortical thymic epithelial cells.
 IT 281217-12-7P
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)
 (aminoazepanone inhibitors of cathepsin L, preparation, and therapeutic use)
 RN 281217-12-7 CAPLUS
 CN 8-Quinolinesulfonyl-4-azepin-4-ylamino-2-oxo-1-(phenylmethyl)ethyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.


L4 ANSWER 86 OF 261 CAPLUS COPYRIGHT 2004 ACS ON STN (Continued)
 19 synthetic examples, a PARP assay for 74 compds., models for focal cerebral ischemia, heart ischemia/reperfusion injury (rats) and gout. Examples also include an evaluation of neuroprotective effects on chronic constriction injury (rats). E.g., 2-amino-6(5H)phenanthridinone and 4-Me benzenesulfonyl chloride were reacted (dioxane, Et3N, 40°C, 30 h) to give I (R1, R2, R4 = H; R3 = NHSO2-C6H4-CH3-p; II) as a brown solid in 93% yield. II had IC50 = 0.12 μM for poly(ADP-ribose) polymerase (PARP). I are useful in the treatment of tissue damage resulting from cell damage due to apoptosis, neuronal mediated tissue damage, neurological disorders, neurodegenerative diseases, etc.
 IT 376609-03-9P
 RL: PAC (Pharmacological activity); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)
 (drug, preparation of sulfonamide/carbamide derivs. of 6(5H)phenanthridinones and use as poly(ADP-ribose) polymerase (PARP) inhibitors)
 RN 376609-03-9 CAPLUS
 CN L-Phenylalanine, N-[(5,6-dihydro-6-oxo-3-phenanthridinyl)carbonyl]-4-fluoro- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



REFERENCE COUNT: 3 THERE ARE 3 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT


L4 ANSWER 87 OF 261 CAPLUS COPYRIGHT 2004 ACS ON STN (Continued)



REFERENCE COUNT: 1 THERE ARE 1 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 88 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
 ACCESSION NUMBER: 2001:764908 CAPLUS
 DOCUMENT NUMBER: 136:65517
 TITLE: The pH-Dependent Primary Photoreactions of Ochratoxin A
 AUTHOR(S): Il'ichev, Yuri V.; Perry, Jennifer L.; Manderville, Richard A.; Chignell, Colin F.; Simon, John D.
 CORPORATE SOURCE: Department of Chemistry, Duke University, Durham, NC. 27708, USA
 SOURCE: Journal of Physical Chemistry B (2001), 105(45), 11369-11376
 CODEN: JPCBFK; ISSN: 1089-5647
 PUBLISHER: American Chemical Society
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 AB Steady-state and time-resolved spectroscopies are used to elucidate the primary photoprocesses following the excitation of ochratoxin A (OTA), its dechlorinated derivative ochratoxin B (OTB), and O-Me ether of OTA (MOA). The excited-state dynamics of OTA and OTB depend on the protonation of the isocoumarin moiety. Fluorescence spectra of the protonated forms reveal anomalously large Stokes shifts that are attributed to the enol tautomer formed via an intramolecular excited-state proton transfer. No evidence for "normal" emission of the keto form of OTA and OTB is found even in aqueous solutions. MOA, which lacks a proton on the phenol moiety and exists, therefore, only in the keto form, exhibits weak fluorescence with a substantially smaller Stokes shift. The deprotonated species show relatively strong emission typical for phenolate anions. OTA decomposes slowly upon UV irradiation in aqueous solutions. The photoreaction quantum yield varies significantly with solution pH and O₂ concentration. The highest yield is observed for the deprotonated form of OTA in deoxygenated solutions. The corresponding hydroquinone (OHQ) is identified as a major photoproduct. Monophotonic photoionization of the fully deprotonated OTA in aqueous solution is demonstrated with nanosecond laser flash photolysis. In the absence of O₂ and other scavengers, hydrated electrons are trapped by OTA in the ground state with the diffusion-controlled rate constant. Photoirradiation of OTA in the presence of supercoiled plasmid DNA results in the formation of relaxed circular DNA. The yield of circular DNA correlates with the quantum yield of OTA photodecomposition in these solutions, because the photocleavage efficiency is higher in the absence of O₂ and at basic pH.
 IT 205034-32-8
 RL: CPS (Chemical process); FMU (Formation, unclassified); PEP (Physical, engineering or chemical process); PRP (Properties); FORM (Formation, nonpreparative); PROC (Process)
 (pH-dependent primary photoreactions of ochratoxin A)
 RN 205034-32-8 CAPLUS
 CN L-Phenylalanine, N-[[[(1R)-3,4-dihydro-5,8-dihydroxy-3-methyl-1-oxo-1H-2-benzopyran-7-yl]carbonyl]- (9CI) (CA INDEX NAME)
 Absolute stereochemistry.

L4 ANSWER 88 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)




REFERENCE COUNT: 105 THERE ARE 105 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE REFORMAT

L4 ANSWER 89 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
 ACCESSION NUMBER: 2001:636473 CAPLUS
 DOCUMENT NUMBER: 135:344629
 TITLE: Total Synthesis of Asperazine
 AUTHOR(S): Govek, Steven P.; Overman, Larry E.
 CORPORATE SOURCE: Department of Chemistry, University of California, Irvine, CA. 92697-2025, USA
 SOURCE: Journal of the American Chemical Society (2001), 123(38), 9468-9469
 CODEN: JACSAT; ISSN: 0002-7863
 PUBLISHER: American Chemical Society
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 GI

 I
 AB The first total synthesis of asperazine (I) was accomplished in 22 steps from readily available starting materials. This synthesis confirms the structure of asperazine and provides yet another example of the tremendous utility of intramolecular Heck reactions for forging highly congested quaternary carbon centers.
 IT 370890-28-1P
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)
 (total synthesis of asperazine)
 RN 370890-28-1 CAPLUS
 CN D-Phenylalanine, 7-[[[(2S,3aS,8aS)-1,8-bis[(1,1-dimethylethoxy)carbonyl]-2,3,8,8a-tetrahydro-2-[[[(1R)-2-methoxy-2-oxo-1-(phenylmethyl)ethyl]amino]carbonyl]pyrrolo[2,3-b]indol-3a(1H)-yl]-N-[(1,1-dimethylethoxy)carbonyl]-L-tryptophyl-, methyl ester (9CI) (CA INDEX NAME)
 Absolute stereochemistry.


L4 ANSWER 89 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)



REFERENCE COUNT: 23 THERE ARE 23 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE REFORMAT

L4 ANSWER 90 OF 261 CAPLUS COPYRIGHT 2004 ACS ON STN
 ACCESSION NUMBER: 2001:597979 CAPLUS
 DOCUMENT NUMBER: 135:167035
 TITLE: Preparation of tyrosine derivatives having anti-leukotriene activity
 INVENTOR(S): Makovec, Francesco, Paris, Walter; Rovati, Lucio Claudio
 PATENT ASSIGNEE(S): Rotta Research Laboratorium S.P.A., Italy
 SOURCE: PCT Int. Appl., 27 pp.
 CODEN: PIXXD2
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2001058892	A1	20010816	WO 2001-EP1315	20010207
W: AU, CA, JP, US				
RW: AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR				
IT 1320162	B1	20031118	IT 2000-T0127	20000209
EP 1255749	A1	20021113	EP 2001-905744	20010207
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR				
JP 2003522768	T2	20030729	JP 2001-558442	20010207
US 2003087910	A1	20030508	US 2002-203424	20020808
US 6605722	B2	20030812		
PRIORITY APPLN. INFO.:			IT 2000-T0127	A 20000209
			WO 2001-EP1315	W 20010207
OTHER SOURCE(S):		MARPAT 135:167035		
GI				




AB Comps. I [R1, R2 = H, C1-4 alkyl, halo, MeO, cyano, CF3; R3 = (un)substituted Ph, pyridyl or (iso)quinolinyl, 1- or 2-naphthyl, 2- or 3-indolyl or N-alkyl derivs., 2-, 5- or 6-quinoxalyl, cinnolyl, benzimidazolyl, which may have the L- or D-configuration or be racemic, were prepared and are useful in the treatment of pathol. conditions sensitive to leukotriene inhibition. Thus, O-(2-quinolinylmethyl)-N-quinolaldehyde-DL-tyrosine was prepared by acylation of DL-tyrosine Me ester with quinaldic acid, O-alkylation with 2-chloromethylquinoline hydrochloride, and saponification. The product showed IC50x10-9 M = 20.0 for inhibition of binding of (3H)-LTD4 to guinea pig lung membranes.

L4 ANSWER 91 OF 261 CAPLUS COPYRIGHT 2004 ACS ON STN
 ACCESSION NUMBER: 2001:597958 CAPLUS
 DOCUMENT NUMBER: 135:166827
 TITLE: Preparation of 1H-indole-3-carboxamides, 1H-pyrrolo[4,3-b]indol-1-ones and pyrrolo[1,2,3-de]-1,4-benzoxazine-6-carboxamides as cannabinoid receptor modulators for treating respiratory and non-respiratory diseases
 INVENTOR(S): Leftheris, Katerina; Zhao, Rulin; Chen, Bang-Chi; Kiener, Peter; Wu, Hong; Pandit, Chennagiri R.; Wroblewski, Stephen; Chen, Ping; Hynes, John, Jr.; Longphre, Malinda; Norris, Derek J.; Spergel, Steven; Tokarski, John
 PATENT ASSIGNEE(S): Bristol-Myers Squibb Company, USA; et al.
 SOURCE: PCT Int. Appl., 199 pp.
 CODEN: PIXXD2
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 2
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2001058869	A2	20010816	WO 2001-US4131	20010208
WO 2001058869	A3	20020124		
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MY, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM				
RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG				
EP 1254115	A2	20021106	EP 2001-907144	20010208
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR				
JP 2004502642	T2	20040129	JP 2001-558420	20010208
PRIORITY APPLN. INFO.:			US 2000-181818 P	20000211
			WO 2001-US4131	W 20010208
OTHER SOURCE(S):		MARPAT 135:166827		
GI				

L4 ANSWER 90 OF 261 CAPLUS COPYRIGHT 2004 ACS ON STN (Continued)
 IT 353798-73-99
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)
 (preparation of tyrosine derivs. having anti-leukotriene activity)
 RN 353798-73-9 CAPLUS
 CN Tyrosine, N-(2-quinolinylcarbonyl)-O-(2-quinolinylmethyl)- (9CI) (CA INDEX NAME)




REFERENCE COUNT: 3 THERE ARE 3 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT


L4 ANSWER 91 OF 261 CAPLUS COPYRIGHT 2004 ACS ON STN (Continued)

AB The title compts. [I; A, B = C, N so that ring X = pyrrole, pyrazole or imidazole (wherein when A = N, the group CONR1R2 is attached to atom C-3 and R5 does not exist; and when A = C, one of CONR1R2 and R5 is attached to A and the other to atom C-3; and when B = C, two R4 groups attached to B and atom C-5, resp., form a fused 6-membered heteroaryl]; f = 0-1; g = 1-2; R1, R2 = H, alkyl, heterocycloalkyl, etc.; R2 together with R1 or R5 forms a 5-6 membered heterocyclo; R3 = H, alkyl, aryl, etc.; R4 is attached to atom C-5 and optionally B and is H, alkyl, aryl, etc.; R5 is attached to A or atom C-3 and is H, alkyl, aryl, etc.; R5 together with R2 forms a heterocyclo; useful as cannabinoid receptor modulators (no data given) for treating respiratory and non-respiratory leukocyte-activation associated diseases, were prepared. Thus, reacting the acid chloride II [X = Cl] (multi-step synthesis given) with 2,2,6,6-tetramethylcyclohexylamine afforded the pyrrolo[1,2,3-de]-1,4-benzoxazine-6-carboxamide II [X = 2,2,6,6-tetramethylcyclohexylamino].
 IT 354569-38-3P
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)
 (preparation of 1H-indole-3-carboxamides, 1H-indazole-3-carboxamides, 1H-pyrrolo[4,3-b]indol-1-ones and pyrrolo[1,2,3-de]-1,4-benzoxazine-6-carboxamides as cannabinoid receptor modulators for treating respiratory and non-respiratory diseases)
 RN 354569-38-3 CAPLUS
 CN L-Phenylalanine, N-[(3R)-2,3-dihydro-5-methyl-3-(4-morpholinylmethyl)pyrrolo[1,2,3-de]-1,4-benzoxazin-6-yl]carbonyl]-, methyl ester, monohydrochloride (9CI) (CA INDEX NAME)

Absolute stereochemistry.




● HCl




L4 ANSWER 92 OF 261 CAPLUS COPYRIGHT 2004 ACS ON STN
 ACCESSION NUMBER: 2001:594627 CAPLUS
 DOCUMENT NUMBER: 135:314766
 TITLE: Electrochemical Oxidation of Ochratoxin A: Correlation with 4-Chlorophenol
 AUTHOR(S): Calcutt, M. Wade; Gillman, Ivan G.; Nofhle, Ronald E.; Manderville, Richard A.
 CORPORATE SOURCE: Department of Chemistry, Wake Forest University, Winston-Salem, NC, 27109-7486, USA
 SOURCE: Chemical Research in Toxicology (2001), 14(9), 1266-1272
 CODEN: CRTOEC; ISSN: 0893-228X
 PUBLISHER: American Chemical Society
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 AB Ochratoxin A (OTA) is a mycotoxin implicated in human kidney carcinogenesis, in which oxidative activation is believed to play a key role. To gain an understanding of the oxidative behavior of the toxin, we have carried out an electrochem. study and have observed a direct correlation between the electrochem. of OTA and 4-chlorophenol (4-ClPhOH). Cyclic voltammetry (CV) of OTA in acetonitrile (MeCN) showed that the toxin exhibits an irreversible oxidative half-peak potential (Ep/2) of 1.81 V vs. SCE; the corresponding value for 4-ClPhOH is 1.59 V. For both compds., subsequent scans revealed the appearance of the resp. hydroquinone/benzoquinone couple, which formed from the oxidation of the parent para-chlorophenol moiety. The hydroquinone of OTA (OTHQ) exhibited Ep/2 = 1.21 V in MeCN. Deprotonation of OTA to form the phenolate (OTA-) lowered the potential to Ep/2 = 1.0 V in MeCN. However, from the oxidation of OTA, no evidence for the OTHQ/OTQ redox couple was found. In aqueous phosphate buffer (pH 6-8), the electrochem. behavior of OTA mimicked that observed for OTA- in MeCN; Ep/2 was .apprx.0.8 V vs. SCE and subsequent scans did not generate the OTHQ/OTQ redox couple. From capillary electrophoresis (CE), a diffusion coefficient (D) of 0.48*10⁻⁵ cm² s⁻¹ was determined for OTA in phosphate buffer, pH 7.0. Combining this value with electrochem. data suggested that OTA undergoes a 1H+/1e oxidation in aqueous media. The biol. implications of these findings with respect to the oxidative metabolism of OTA, and other chlorinated phenols, are discussed.
 IT 205034-32-8
 RL: PMU (Formation, unclassified); PRP (Properties); FORM (Formation, nonpreparative)
 (electrochem. oxidation of ochratoxin A and correlation with chlorophenol)
 RN 205034-32-8 CAPLUS
 CN L-Phenylalanine, N-[[[(3R)-3,4-dihydro-5,8-dihydroxy-3-methyl-1-oxo-1H-2-benzopyran-7-yl]carbonyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



L4 ANSWER 93 OF 261 CAPLUS COPYRIGHT 2004 ACS ON STN
 ACCESSION NUMBER: 2001:584421 CAPLUS
 DOCUMENT NUMBER: 135:318403
 TITLE: Asymmetric Pictet-Spengler reactions: synthesis of 1,2,3,4-tetrahydroisoquinoline carboxylic acid (Tic) chimeras
 AUTHOR(S): Spengler, Jan; Schedel, Hartmut; Sieler, Joachim; Quaedflieg, Peter J. L. M.; Broxterman, Quirinus B.; Duchateau, Alexander L. L.; Burger, Klaus
 CORPORATE SOURCE: Department of Organic Chemistry, University of Leipzig, Leipzig, 04103, Germany
 SOURCE: Synthesis (2001), (10), 1513-1518
 CODEN: SYNTEF; ISSN: 0039-7881
 PUBLISHER: Georg Thieme Verlag
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 OTHER SOURCE(S): CASREACT 135:318403
 AB A preparatively simple diastereoselective synthesis of the amino acid chimera (1S,3S)-1,2,3,4-tetrahydroisoquinoline-1,3-dicarboxylic acid from hexafluoroacetone-protected phenylalanine and glyoxylic acid hydrate via Pictet-Spengler reaction is described. The potential of the reaction of hexafluoroacetone-protected phenylalanine with other aldehydes was scrutinized.
 IT 367952-41-8P
 RL: SPN (Synthetic preparation); PREP (Preparation) (preparation of)
 RN 367952-41-8 CAPLUS
 CN 1-Isosquinoilincarboxylic acid, 3-[[[(1S)-2-amino-2-oxo-1-(phenylmethyl)ethyl]amino]carbonyl]-1,2,3,4-tetrahydro-, (1S,3S)- (9CI) (CA INDEX NAME)



Absolute stereochemistry. Rotation (-).



REFERENCE COUNT: 30 THERE ARE 30 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 92 OF 261 CAPLUS COPYRIGHT 2004 ACS ON STN (Continued)
 REFERENCE COUNT: 45 THERE ARE 45 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT


L4 ANSWER 94 OF 261 CAPLUS COPYRIGHT 2004 ACS ON STN
 ACCESSION NUMBER: 2001:584409 CAPLUS
 DOCUMENT NUMBER: 135:288611
 TITLE: Enantioselective Synthesis of a Mitosane Core Assisted by Diversity-Based Catalyst Discovery
 AUTHOR(S): Papaioannou, Nikolaos; Evans, Catherine A.; Blank, Jarred T.; Miller, Scott J.
 CORPORATE SOURCE: Department of Chemistry Merkert Chemistry Center, Boston College, Chestnut Hill, MA, 02467-3860, USA
 SOURCE: Organic Letters (2001), 3(18), 2879-2882
 CODEN: ORLEF7; ISSN: 1523-7060
 PUBLISHER: American Chemical Society
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 OTHER SOURCE(S): CASREACT 135:288611
 GI



AB Synthesis of (-)-mitosane I in optically pure form is reported. A retrosynthetic plan that proceeds through racemic allylic alc. II (R = OH) was carried out. Intermediate II served as a test substrate for a rapid screen of a small library (152 members) of peptide-based kinetic resolution catalysts. Peptide III (R = β-OH) was found to effect kinetic resolution with k_{rel} = 27. Alc. II (R = β-OH) was then converted to optically pure I in eight steps.
 IT 365223-05-8P
 RL: CAT (Catalyst use); PRP (Properties); SPN (Synthetic preparation); PREP (Preparation); USES (Uses) (enantioselective synthesis of a mitosane core assisted by diversity-based catalyst discovery)

L4 ANSWER 94 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)
 RN 36523-05-8 CAPLUS
 CN L-Phenylalanine, N-[(1,1-dimethylethoxy)carbonyl]-3-methyl-L-histidyl-(3R)-1,2,3,4-tetrahydro-3-isoquinolinecarboxyl-(nS)- α -aminobenzenebutanoyl-L-valyl-, methyl ester (9CI) (CA INDEX NAME)

Absolute stereochemistry.



REFERENCE COUNT: 43 THERE ARE 43 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 95 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
 ACCESSION NUMBER: 2001:527755 CAPLUS
 DOCUMENT NUMBER: 135:266637
 TITLE: Is There a Difference between Leads and Drugs? A Historical Perspective
 AUTHOR(S): Oprea, Tudor I.; Davis, Andrew M.; Teague, Simon J.; Leeson, Paul D.
 CORPORATE SOURCE: AstraZeneca R&D Molndal EST Lead Informatics, Molndal, S 431 83, Swed.
 SOURCE: Journal of Chemical Information and Computer Sciences (2001), 41(5), 1308-1315
 CODEN: JCISDH; ISSN: 0095-2338
 PUBLISHER: American Chemical Society
 DOCUMENT TYPE: Journal
 LANGUAGE: English


AB To be considered for further development, lead structures should display the following properties: (1) simple chemical features, amenable for chemical optimization; (2) membership to an established SAR series; (3) favorable patent situation; and (4) good absorption, distribution, metabolism, and excretion (ADME) properties. There are two distinct categories of leads: those that lack any therapeutic use (i.e., "pure" leads), and those that are marketed drugs themselves but have been altered to yield novel drugs. We have previously analyzed the design of leadlike combinatorial libraries starting from 18 lead and drug pairs of structures (S. J. Teague et al. Angew. Chemical, Int. Ed. Engl. 1999, 38, 3743-3748). Here, we report results based on an extended dataset of 96 lead-drug pairs, of which 62 are lead structures that are not marketed as drugs, and 75 are drugs that are not presumably used as leads. We examined the following properties: MW (mol. weight), CMR (the calculated mol. refractivity), RNG (the number of rings),

RTB (the number of rotatable bonds), the number of hydrogen bond donors (HDO) and acceptors (HAC), the calculated logarithm of the n-octanol/water partition (CLogP), the calculated logarithm of the distribution coefficient at pH 7.4 (LogD74), the Daylight-fingerprint druglike score (DFFS), and the property and pharmacophore features score (PPFS). The following differences were observed between the medians of drugs and leads: AMW = 69; ACMR = 1.8; ARNG = AHAC = 1; ARTB = 2; ACLogP = 0.43; ALogD74 = 0.97; AHDO = 0; ADFFS = 0.15; APFFS = 0.12. Lead structures exhibit, on the average, less mol. complexity (less MW, less number of rings and rotatable bonds), are less hydrophobic (lower CLogP and LogD74), and less druglike (lower druglike scores). These findings indicate that the process of optimizing a lead into a drug results in more complex structures. This information should be used in the design of novel combinatorial libraries that are aimed at lead discovery.

IT 174635-53-1, SB 218795 R
 RL: PRP (Properties)
 (drug design and structure-activity relationship between leads and leadlike drugs)
 RN 174635-53-1 CAPLUS
 CN Benzenecetic acid, α -[[(2-phenyl-4-quinolinyl)carbonyl]amino]-, methyl ester, (nR) (9CI) (CA INDEX NAME)

Absolute stereochemistry. Rotation (-).

L4 ANSWER 95 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)



REFERENCE COUNT: 30 THERE ARE 30 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT


L4 ANSWER 96 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
 ACCESSION NUMBER: 2001:514247 CAPLUS
 DOCUMENT NUMBER: 135:257449
 TITLE: Interaction of Ferrocenyl-Dipeptides with 3-Aminopyrazole Derivatives: β -Sheet Models? A Synthetic, Spectroscopic, Structural, and Electrochemical Study
 AUTHOR(S): Sawczko, Pete; Enright, Gary D.; Kraatz, Heinz-Bernhard
 CORPORATE SOURCE: Department of Chemistry, University of Saskatchewan, Saskatoon, SK, S7N 5C9, Can.
 SOURCE: Inorganic Chemistry (2001), 40(17), 4409-4419
 CODEN: INOCAL; ISSN: 0020-1669
 PUBLISHER: American Chemical Society
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 OTHER SOURCE(S): CASREACT 135:257449

AB The use of 3-aminopyrazole deriva. as β -sheet templates is investigated using a series of ferrocenyl-dipeptides [Fc-(Gly)2-OEt, Fc-(Ala)2-OCH2Ph, Fc-Leu-Phe-OMe, Fc-Val-Phe-OMe, Fc-(Phe)2-OMe, Fc-(Leu)2-OMe, Fc-(Val)2-OMe]. The synthesis and full characterization of the ferrocenyl-dipeptides are reported. The solid-state structures of Fc-(Gly)2-OMe and Fc-Leu-Phe-OMe show extensive hydrogen bonding of the podand peptide substituents, resulting in the formation of supramol. Fc-dipeptide assemblies. For Fc-(Gly)2-OMe, this can be described as a parallel β -sheet, whereas intermol. interactions in Fc-Leu-Phe-OMe result in the formation of supramol. helical structures. The saturation titrns. of Fc-dipeptides with 3-amino-5-methylpyrazole (3-AMP) and 3-trifluoroacetamido-5-methylpyrazole (3-TPAC-AMP) show a 1:1 interaction of the Fc-peptide with the aminopyrazole deriva. IR measurements in solution confirm binding to the top face of the Fc-dipeptide and the involvement of the Fc-C=O and the ester C=O groups in establishing H-bonding interactions with the 3-TPAC-AMP. However, binding consts. in chloroform are low and range from 8 to 27 M⁻¹, which correspond to binding energies of 5-7 kJ mol⁻¹. In higher polarity solvents, such as acetonitrile or acetone, the binding consts. are below 5 M⁻¹, emphasizing the limited utility of 3-AMP deriva. as β -sheet templates. Electrochem. measurements confirm the weak interactions between the various Fc-dipeptides and 3-TPAC-AMP. Typical shifts in the redox potential of the Fc moiety are in the range 0-20 mV. Attempts to modify 3-AMP at the 3-position by carbodiimide coupling with Boc-Aib-OH (in order to enhance the binding to the Fc-peptides) resulted in 2-(Boc-Aib)-substituted 3-AMP deriva. Substitution at the 2-position blocks the binding site, and no interactions with Fc-dipeptides are observed.


IT 362056-37-9
 RL: PMU (Formation, unclassified); PRP (Properties); FORM (Formation, nonpreparative)
 (association consts. for the 1:1 interaction complexes of (ferrocenyl)dipeptides with aminopyrazoles as β -sheet templates)
 RN 362056-37-9 CAPLUS
 CN L-Phenylalanine, N-(ferrocenylcarbonyl)-L-phenylalanyl-, methyl ester, compd. with 5-methyl-1H-pyrazol-3-amine (1:1) (9CI) (CA INDEX NAME)

CM 1
 CRN 245123-57-3
 CMP C30 H30 Fe N2 O4
 CCI CCS

L4 ANSWER 96 OF 261 CAPLUS COPYRIGHT 2004 ACS ON STN (Continued)



CM 2

CRN 31230-17-8
CMP C4 H7 N3


REFERENCE COUNT: 70 THERE ARE 70 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 97 OF 261 CAPLUS COPYRIGHT 2004 ACS ON STN
 ACCESSION NUMBER: 2001:436800 CAPLUS
 DOCUMENT NUMBER: 135:178024
 TITLE: Novel sesquiterpenoids from the roots of *Phyllanthus emblica*
 AUTHOR(S): Zhang, Ying-Jun; Tanaka, Takashi; Iwamoto, Yoko; Yang, Chong-Ren; Kouno, Isao
 CORPORATE SOURCE: Faculty of Pharmaceutical Sciences, Nagasaki University, Nagasaki, 852-8521, Japan
 SOURCE: Journal of Natural Products (2001), 64(7), 870-873
 CODEN: JNPRDF; ISSN: 0163-3864
 PUBLISHER: American Chemical Society
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 GI

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

AB Three novel bisabolane-type sesquiterpenoids, phyllaemblic acids B (I) and C (II) and phyllaemblicin D (III), together with two new phenolic glycosides, 2-carboxymethylphenol 1-O-β-D-glucopyranoside (IV) and 2,6-dimethoxy-4-(2-hydroxyethyl)phenol 1-O-β-D-glucopyranoside (V), were isolated from the roots of *Phyllanthus emblica*. The structures of I-V were established by spectral and chemical methods. The absolute stereochem. of I and II was determined by applying the PGME method.
 IT 354813-54-0P
 RL: PRP (Properties); SPN (Synthetic preparation); PREP (Preparation) (preparation and properties of)
 RN 354813-54-0 CAPLUS
 CN Benzeneacetic acid, α-[[[[(2'S,3R,3aS,4R,4'S,5'R,6S,7aR)-3',3a,4,4',5,5',6,6',7,7a-decahydro-3,4,4'-trihydroxy-3,5'-bis(hydroxymethyl)spiro[benzofuran-2(3H),2'-(2H)pyran]-6-yl]carbonyl]amino]-, methyl ester, (aS)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



REFERENCE COUNT: 11 THERE ARE 11 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 98 OF 261 CAPLUS COPYRIGHT 2004 ACS ON STN
 ACCESSION NUMBER: 2001:416769 CAPLUS
 DOCUMENT NUMBER: 135:33476
 TITLE: Preparation of naphth[1,2-d]imidazoles as thrombopoietin mimetics
 INVENTOR(S): Luengo, Juan I.; Duffy, Kevin J.; Price, Alan T.; Zhang, Lihua
 PATENT ASSIGNEE(S): Smithkline Beecham Corporation, USA
 SOURCE: PCT Int. Appl., 90 pp.
 CODEN: PIXXD2
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2001039773	A1	20010607	WO 2000-US33432	20001206
W: AU, CA, JP, US RW: AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR				
EP 1244446	A1	20021002	EP 2000-984123	20001206
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, FI, CY, TR				
JP 2003515560	T2	20030507	JP 2001-541505	20001206
US 2003083361	A1	20030501	US 2002-148945	20020912
PRIORITY APPLN. INFO.: US 1999-169130P P 19991206 WO 2000-US33432 W 20001206				
OTHER SOURCE(S): MARPAT 135:33476				
GI				



AB The title compds. [I; R1-R5 = H, CO2R11, alkyl, etc. (wherein R11 = H, alkyl, cycloalkyl, etc.); R6 = H, alkyl, cycloalkyl, etc.; R7, R8 = absent, H, alkyl, etc.; m = 0-6; X = S, O, NH, etc.; AR = (un)substituted cyclic or polycyclic aromatic ring containing from 3-16 carbon atoms, optionally containing one or more heteroatoms] and their pharmaceutically acceptable salts which are non-peptide TPO mimetics, and are useful in enhancing platelet production, were prepared and formulated. E.g., a 3-step synthesis of the title compound II.HCl was described. Biol. data for compds. I were given.

L4 ANSWER 98 OF 261 CAPLUS COPYRIGHT 2004 ACS ON STN (Continued)
 IT 343603-01-0P
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses) (preparation of naphth[1,2-d]imidazoles as thrombopoietin mimetics)
 RN 343603-01-0 CAPLUS
 CN L-Phenylalanine, N-[[[2-(3',4'-dimethyl[1,1'-biphenyl]-3-yl)-9-hydroxy-1H-naphth[1,2-d]imidazol-7-yl]carbonyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




REFERENCE COUNT: 3 THERE ARE 3 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 99 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
 ACCESSION NUMBER: 2001:356253 CAPLUS
 DOCUMENT NUMBER: 134:366810
 TITLE: Preparation of 7-[(4'-trifluoromethyl-biphenyl-2-carbonyl)amino]-quinoline-3-carboxylic acid amides for inhibiting the secretion of apolipoprotein B
 INVENTOR(S): Bertinato, Peter; Hamanaka, Ernest Seiichi; Ruggeri, Roger Benjamin; Wilson, Douglas Paul
 PATENT ASSIGNEE(S): Pfizer Products Inc., USA
 SOURCE: Eur. Pat. Appl., 124 pp.
 CODEN: EPXXDW
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
EP 1099701	A1	20010516	EP 2000-309947	20001109
R: AT, BR, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO				
BR 2000005322	A	20010717	BR 2000-5322	20001109
US 6369075	B1	20020409	US 2000-711281	20001109
JP 2001139555	A2	20010522	JP 2000-344267	20001110
US 2002132806	A1	20020919	US 2002-54455	20020122
US 6713489	B2	20040330		

PRIORITY APPLN. INFO.:
 US 1999-164803P P 19991110
 US 2000-224956P P 20000811
 US 2000-711281 A3 20001109

OTHER SOURCE(S): MARPAT 134:366810
 GI



AB The title compds. [I, R1 = H, alkyl; R2 = H, alkyl, CHX2, etc.; NR1R2 = 3-7 membered heterocycloalkyl comprising 1-3 heteroatoms; X = (un)substituted aryl, heteroaryl, etc.; Rb = H, alkyl] that inhibit the secretion of apolipoprotein B and/or inhibit microsomal triglyceride transfer protein, and therefore useful in treating and/or preventing atherosclerosis, obesity, diabetes, hyperlipidemia, hyperlipoproteinemia, hypercholesterolemia, hypertriglyceridemia, hypopalipaproteinemia, pancreatitis, myocardial infarction, stroke, restenosis, or Syndrome X.

L4 ANSWER 100 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
 ACCESSION NUMBER: 2001:356207 CAPLUS
 DOCUMENT NUMBER: 134:348283
 TITLE: Methods of administering apolipoprotein B secretion/microsomal triglyceride transfer protein inhibitors
 INVENTOR(S): Chang, George; Vincent, John
 PATENT ASSIGNEE(S): Pfizer Products Inc., USA
 SOURCE: Eur. Pat. Appl., 42 pp.
 CODEN: EPXXDW
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
EP 1099442	A2	20010516	EP 2000-309907	20001108
EP 1099442	A3	20021204		
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO				


PRIORITY APPLN. INFO.:
 US 1999-164579P P 19991110

AB Methods are provided for administration of apoB secretion/MTP inhibitors. The methods comprise administration prior to or during a period of somnolence. Preparation of inhibitors is also described.

IT 339290-34-5P
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)
 (apolipoprotein B secretion/microsomal triglyceride transfer protein inhibitor administration prior to or during somnolence period)


RN 339290-34-5 CAPLUS
 CN 3-Quinolincarboxamide, N-[(1S)-2-amino-2-oxo-1-(phenylmethyl)ethyl]-7-[[[4'-(trifluoromethyl)-1,1'-biphenyl]-2-yl]carbonyl]amino]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



L4 ANSWER 99 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)
 were prepd. E.g., a multi-step synthesis of I [R1 = H; R2 = CH2-pyridyl]2 was given.
 IT 339290-34-5P
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)
 (preparation of 7-[(4'-trifluoromethyl-biphenyl-2-carbonyl)amino]-quinoline-3-carboxylic acid amides for inhibiting the secretion of apolipoprotein B)
 RN 339290-34-5 CAPLUS
 CN 3-Quinolincarboxamide, N-[(1S)-2-amino-2-oxo-1-(phenylmethyl)ethyl]-7-[[[4'-(trifluoromethyl)-1,1'-biphenyl]-2-yl]carbonyl]amino]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




REFERENCE COUNT: 2 THERE ARE 2 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 101 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
 ACCESSION NUMBER: 2001:338483 CAPLUS
 DOCUMENT NUMBER: 134:353176
 TITLE: Preparation of urea derivatives as VLA-4 antagonists
 INVENTOR(S): Okuyama, Akihiko; Ikegami, Satoru; Harada, Tatsuhiko; Maruyama, Tatsuya; Matsumura, Yuzuru; Nagata, Naoya; Fukui, Hideto; Fujimoto, Kyouko
 PATENT ASSIGNEE(S): Kaken Pharmaceutical Co., Ltd., Japan
 SOURCE: PCT Int. Appl., 72 pp.
 CODEN: PIXXD2
 DOCUMENT TYPE: Patent
 LANGUAGE: Japanese
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2001032610	A1	20010510	WO 2000-JP7571	20001027
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MY, NZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM				
RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG				

PRIORITY APPLN. INFO.:
 JP 1999-310316 A 19991029

OTHER SOURCE(S): MARPAT 134:353176
 GI



AB The title compds. I [R1 is hydrogen, alkyl, etc.; X is hydrogen, halogeno, alkyl, aryl, arylamide, etc.; Y is oxygen or sulfur; and Z is a hydrocarbon or heterocyclic group containing a nitrogen atom through which Z is bonded to the carbon atom of CY; the asterisk indicates an asym. carbon] are prepared. Processes for the preparation of I are also claimed. Several compds. of this invention in vitro at 0.01 nM to 3.7 nM gave 50% inhibition of VLA-4/VCAM-1 adhesion.


IT 339001-71-7P
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)
 (preparation of urea derivs. as VLA-4 antagonists)

RN 339001-71-7 CAPLUS
 CN L-Phenylalanine, 4-[(2,6-dichlorobenzoyl)amino]-N-[(4-ethyl-3,4-dihydro-1(2H)-quinolinyl)carbonyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

09/ 964,161

L4 ANSWER 101 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)




REFERENCE COUNT: 8 THERE ARE 8 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 102 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 2001:330818 CAPLUS
DOCUMENT NUMBER: 135:137682
TITLE: The synthesis of amino acid-functionalized β -carbolines as topoisomerase II inhibitors
AUTHOR(S): Deveau, A. M.; Labroli, M. A.; Dieckhaus, C. M.; Barthen, M. T.; Smith, K. S.; Macdonald, T. L.
CORPORATE SOURCE: Department of Chemistry, University of Virginia, Charlottesville, VA, 22901, USA
SOURCE: Bioorganic & Medicinal Chemistry Letters (2001), 11(10), 1251-1255
CODEN: BMCLES; ISSN: 0960-894X
PUBLISHER: Elsevier Science Ltd.
DOCUMENT TYPE: Journal
LANGUAGE: English
OTHER SOURCE(S): CASREACT 135:137682

AB The synthesis and biol. activity of amino acid-functionalized β -carboline derivs., which are structurally related to azatoxin and the tryprostatins, are reported. These compds. were assayed for their growth inhibition properties in H520 and PC3 cell lines and were examined for their abilities to inhibit topoisomerase II-mediated DNA relaxation.
IT 352015-64-6P
RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); BIOL (Biological study); PRSP (Preparation)
(synthesis of amino acid-functionalized β -carbolines as topoisomerase II inhibitors)
RN 352015-64-6 CAPLUS
CN D-Phenylalanine, N-[[[(1R,3S)-2,3,4,9-tetrahydro-1-(4-hydroxy-3,5-dimethoxyphenyl)-1H-pyrido[3,4-b]indol-3-yl]carbonyl]-, methyl ester (9CI) (CA INDEX NAME)

Absolute stereochemistry.



REFERENCE COUNT: 21 THERE ARE 21 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT


L4 ANSWER 103 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 2001:318582 CAPLUS
DOCUMENT NUMBER: 135:120165
TITLE: Potent and selective cathepsin L inhibitors do not inhibit human osteoclast resorption in vitro
AUTHOR(S): James, Ian E.; Marquis, Robert W.; Blake, Simon M.; Hwang, Shing Mei; Gress, Catherine J.; Ru, Yu; Zembryki, Denise; Yamashita, Dennis S.; McQueney, Michael S.; Tomaszek, Thaddeus A.; Oh, Hye-Ja; Gowen, Maxine; Veber, Daniel F.; Lark, Michael W.
CORPORATE SOURCE: Departments of Bone and Cartilage Biology, SmithKline Beecham Pharmaceuticals, King of Prussia, PA, 19406, USA
SOURCE: Journal of Biological Chemistry (2001), 276(15), 11507-11511
CODEN: JBCHA3; ISSN: 0021-9258
PUBLISHER: American Society for Biochemistry and Molecular Biology
DOCUMENT TYPE: Journal
LANGUAGE: English

AB Cathepsins K and L are related cysteine proteases that have been proposed to play important roles in osteoclast-mediated bone resorption. To further examine the putative role of cathepsin L in bone resorption, we have evaluated selective and potent inhibitors of human cathepsin L and cathepsin K in an in vitro assay of human osteoclastic resorption and an in situ assay of osteoclast cathepsin activity. The potent selective cathepsin L inhibitors ($K_i = 0.0099$, 0.034 , and 0.27 nM) were inactive in both the in situ cytochem. assay ($IC_{50} > 1$ μ M) and the osteoclast-mediated bone resorption assay ($IC_{50} > 300$ nM). Conversely, the cathepsin K selective inhibitor was potently active in both the cytochem. ($IC_{50} = 63$ nM) and resorption ($IC_{50} = 71$ nM) assays. A recently reported dipeptide aldehyde with activity against cathepsins L ($K_i = 0.052$ nM) and K ($K_i = 1.57$ nM) was also active in both assays ($IC_{50} = 110$ and 115 nM, resp.) These data confirm that cathepsin K and not cathepsin L is the major protease responsible for human osteoclastic bone resorption.

IT 350796-41-7
RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); BUU (Biological use, unclassified); BIOL (Biological study); USES (Uses)
(potent and selective cathepsin L inhibitors do not inhibit human osteoclast resorption in vitro)
RN 350796-41-7 CAPLUS
CN 8-Quinolincarboxamide, N-[(1S)-2-[[[(4S)-hexahydro-3-oxo-1-(2-pyridinylsulfonyl)-1H-azepin-4-yl]amino]-2-oxo-1-(phenylmethyl)ethyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



L4 ANSWER 103 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)

REFERENCE COUNT: 21 THERE ARE 21 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 104 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
 ACCESSION NUMBER: 2001:228855 CAPLUS
 DOCUMENT NUMBER: 134:252658
 TITLE: Preparation of tyrosine derivatives as inhibitors of $\alpha 4$ containing integrin-mediated binding to ligands VCAM-1 and MadCAM.
 INVENTOR(S): Jackson, David Y.; Sailes, Frederick C.; Sutherland, Daniel P.
 PATENT ASSIGNEE(S): Genentech, Inc., USA
 SOURCE: PCT Int. Appl., 86 pp.
 CODEN: PIXXD2
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2001021584	A1	20010329	WO 2000-US26326	20000925
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM			
RW:	GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG			
EP 1214292	A1	20020619	EP 2000-965417	20000925
R:	AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL			
US 6469047	B1	20021022	US 2000-669779	20000925
JP 2003509488	T2	20030311	JP 2001-524964	20000925
PRIORITY APPLN. INFO.:			US 1999-156062P	P 19990924
			WO 2000-US26326	W 20000925

OTHER SOURCE(S): MARPAT 134:252658
 AB Tyrosine derivs., e.g., $\text{ArCH}_2\text{CH}[\text{N}(\text{A})(\text{Z})]\text{CO}_2\text{Y}$ [Z = H, alkyl; A = B(CH₂)_q-X-, where B = (un)substituted Ph and X = CO, SO₂, null or B = cyanoalkyl, carbocyclyl or heterocyclyl and X = CO; R₆ = H, alkyl, amino, cyano, hydroxy, alkylsulfonyl, etc.; q = 0-3; Y is H, (un)substituted alkoxy, alkoxyalkoxy, aryloxy, alkylaminoalkoxy, dialkylaminoalkoxy, alkylamino, arylamino, heterocyclyl or heteroarylalkyl; Ar is Ph which has hydroxy, carbonate, thiocarbonate, carbamoyloxy or acyloxy groups and optionally (as substituents) were prepared as inhibitors of $\alpha 4$ containing integrin-mediated binding to ligands such as VCAM-1 and MadCAM. Methods of synthesis are described and inhibitory binding data are tabulated for 416 compds., including N-(o-chlorobenzoyl)-O-(allylcarbamoyl)-L-tyrosine, for which IC₅₀ is < 1.0 micromolar.

IT 331470-84-9P
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPH (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)
 (preparation of tyrosine derivs. as inhibitors of $\alpha 4$ containing integrin-mediated binding to ligands VCAM-1 and MadCAM.)

RN 331470-84-9 CAPLUS
 CN L-Tyrosine, N-[(1S)-1,2,3,4-tetrahydro-8-hydroxy-3-isoquinoliny]carbonyl]-, 4-(4-morpholinocarboxylate) (9CI) (CA INDEX

L4 ANSWER 105 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
 ACCESSION NUMBER: 2001:228654 CAPLUS
 DOCUMENT NUMBER: 134:252657
 TITLE: Compounds directed against pilus biogenesis and activity in pathogenic bacteria; methods and compositions for synthesis
 INVENTOR(S): Kihlberg, Jan; Larsson, Andreas; Svensson, Anette; Fex, Tomas; Hultgren, Scott J.; Pinkner, Jerry
 PATENT ASSIGNEE(S): Washington University, USA
 SOURCE: PCT Int. Appl., 85 pp.
 CODEN: PIXXD2
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2001020995	A1	20010329	WO 2000-US26177	20000922
WO 2001020995	C2	20021114		
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM			
RW:	GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG			
PRIORITY APPLN. INFO.:			US 1999-155822P	P 19990923


OTHER SOURCE(S): MARPAT 134:252657
 AB Many Gram-neg. pathogens assemble adhesive structures on their surfaces that allow them to colonize host tissues and cause disease. Novel compns. which inhibit or prevent the formation of a pilus chaperone-subunit complex are disclosed. Interfering with the function of the pilus chaperone neg. affects the chaperone/usher pathway which is one mol. mechanism by which Gram-neg. bacteria assemble adhesive pill structures and thus prevent or inhibit pilus assembly. Also provided are methods for the treatment or prevention of diseases caused by tissue-adhering pilus-forming bacteria by inhibiting the function of pilus chaperones. Also provided are pharmaceutical preps. capable of inhibiting or preventing the formation of a pilus chaperone-subunit complex. Also provided are methods of synthesizing the N-substituted amino acid compds. and compds. useful for the synthesis thereof. In particular, novel fluorinated linker compds. and methods of synthesis are provided. Methods for using the fluorinated linker compds. in methods of solid-phase synthesis of the N-substituted amino acid compds. are also disclosed. Chiral compds. R₄CH₂NR₂COR₃ [R₁-R₃ = (un)substituted alkyl, acyl, aryl, heteroaryl, arylalkyl, heteroarylalkyl, heterocycloalkyl; R₄ = CO₂H, CONH₂, CHO, B(OH)₂, P(O)(OH)₂, alkyl or haloalkyl ketone group] and their salts, esters, and amines are claimed. Thus, N-[2-(1H-indol-3-yl)ethyl]-N-(naphthalene-2-carbonyl)tyrosine was prepared and assayed for affinity for periplasmic chaperones PapD and FimC (K_D estimated as 1-100nM).

IT 331679-48-2P
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPH (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)
 (compds. directed against pilus biogenesis and activity in pathogenic bacteria)

RN 331679-48-2 CAPLUS
 CN L-Phenylalanine, N-[2-(1H-indol-3-yl)ethyl]-N-[(2-oxo-2H-1-benzopyran-3-

L4 ANSWER 104 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)
 NAME)


Absolute stereochemistry.



REFERENCE COUNT: 2 THERE ARE 2 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 105 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)
 yl)carbonyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




REFERENCE COUNT: 1 THERE ARE 1 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 106 OF 261 CAPLUS COPYRIGHT 2004 ACS ON STN
 ACCESSION NUMBER: 2001:222008 CAPLUS
 DOCUMENT NUMBER: 134:252257
 TITLE: Preparation of 2-(indolin-3-yl)quinoline derivatives and compositions in use as antimicrobial agents
 INVENTOR(S): Cuny, Gregory D.; Nauske, James R.; Heefner, Donald L.; Hoemann, Michael Z.; Kumaravel, Gnanasambandam; Melikian-Badalian, Anita; Rossi, Richard F.
 PATENT ASSIGNEE(S): Sepracor, Inc., USA
 SOURCE: U.S., 112 pp., Cont.-in-part of U.S. Ser. No. 878,781, abandoned.
 CODEN: USXXAM
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 7
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 6207679	B1	20010327	US 1998-45051	19980319
WO 9857931	A2	19981223	WO 1998-US12762	19980618
WO 9857931	A3	19990429		
W:	AL, AM, AT, AU, AZ, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, HM, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, RW, GH, GM, KE, LS, MW, SD, SZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG			
EP 991623	A2	20000412	EP 1998-930396	19980618
R:	AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, FI			
US 6172084	B1	20010109	US 1998-99640	19980618
JP 2002505689	T2	20020219	JP 1999-504835	19980618
AU 757059	B2	20030130	AU 1998-79797	19980618
US 6103905	A	20000815	US 1998-213385	19981211
NO 9906269	A	20000216	NO 1999-6269	19991217
US 6376670	B1	20020423	US 2000-658690	20000908
PRIORITY APPLN. INFO.:			US 1997-878781	B2 19970619
			US 1998-45051	A2 19980319
			US 1998-99640	A2 19980618
			WO 1998-US12762	W 19980618
			US 1998-213385	A1 19981211
			US 2000-639622	A2 20000815


OTHER SOURCE(S): MARPAT 134:252257
 GI

L4 ANSWER 106 OF 261 CAPLUS COPYRIGHT 2004 ACS ON STN (Continued)




REFERENCE COUNT: 43 THERE ARE 43 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 106 OF 261 CAPLUS COPYRIGHT 2004 ACS ON STN (Continued)



I



II

AB Title compds. I (wherein; R, R1, R2 and R3 are H, halo, alk(en)(yn)yl, OH, alkoxy, amino, nitro, SH, imine, amide, CO, -(CH2)0-8-R80, etc.; R4 is the same as R-R3 but not H; R5 is the same as R4 except that at least 1(-8) CH2 precede R80; A is (un)substituted with any number of R4 up to the number limited by stability and rules of valence; B is substituted with at least one instance of RS up to the number limited by stability and rules of valence; R80 is (substituted) aryl, cycloalk(en)yl, heterocyclyl or polycyclyl and related quinoline deriva. are prepared as antimicrobial agents. For instance, synthesis of II is accomplished by alkylation of 4-hydroxymethyl-6-trifluoromethyl-2-(N-t-butoxycarbonylindol-3-yl)quinoline with (4-t-butoxycarbonylaminoethyl)benzyl iodide followed by deprotection. There are 282 examples of I provided. The min. inhibitory concentration (MIC) of I against at least one Gram-pos. bacterium is 0.1-10 µg/mL. Certain compds. of formula I have a therapeutic index in primates of at least 10 for the inhibition of infection by at least one Gram-pos. bacterium.

IT 275357-08-9P
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation)
 (preparation and use of quinolinylindole deriva. as antimicrobial agents)

RN 275357-08-9 CAPLUS
 CN 4-Quinolincarboxamide, N-[2-amino-1-[(4-nitrophenyl)methyl]-2-oxoethyl]-2-(5-chloro-1H-indol-3-yl)- (9CI) (CA INDEX NAME)


L4 ANSWER 107 OF 261 CAPLUS COPYRIGHT 2004 ACS ON STN
 ACCESSION NUMBER: 2001:177411 CAPLUS
 DOCUMENT NUMBER: 135:152
 TITLE: A series of quinoline analogues as potent inhibitors of C. albicans prolyl tRNA synthetase
 AUTHOR(S): Yu, X. Y.; Hill, J. M.; Yu, G.; Yang, Y.; Kluge, A. F.; Keith, D.; Finn, J.; Gallant, P.; Silverman, J.; Lim, A.
 CORPORATE SOURCE: Department of Medicinal Chemistry, Cubist Pharmaceuticals, Inc., Cambridge, MA, 02139, USA
 SOURCE: Bioorganic & Medicinal Chemistry Letters (2001), 11(4), 541-544
 CODEN: BMCLEB; ISSN: 0960-894X
 PUBLISHER: Elsevier Science Ltd.
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 OTHER SOURCE(S): CASREACT 135:152

AB A series of quinoline inhibitors of C. albicans prolyl tRNA synthetase was identified. The most potent analog, 2-(4-bromophenyl)-6-chloro-8-methyl-4-quinolinecarboxylic acid, showed IC50=5 nM (Ca. ProRS) with high selectivity over the human enzyme.

IT 342018-11-5
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES (Uses)
 (a series of quinoline analogs as potent inhibitors of C. albicans prolyl tRNA synthetase)

RN 342018-11-5 CAPLUS
 CN L-Phenylalanine, N-[[2-(4-bromophenyl)-6-chloro-8-methyl-4-quinolinyl]carbonyl]-, methyl ester (9CI) (CA INDEX NAME)

Absolute stereochemistry.




REFERENCE COUNT: 16 THERE ARE 16 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 108 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
 ACCESSION NUMBER: 2001:168124 CAPLUS
 DOCUMENT NUMBER: 134:218936
 TITLE: Crystal structure of CDC25 proteins and its use in rational design of inhibitors
 INVENTOR(S): Taylor, Neil R.; Borhani, David; Epstein, David; Rudolph, Johannes; Ritter, Kurt; Fujimori, Taro; Robinson, Simon; Eckstein, Jens; Haupt, Andreas; Walker, Nigel; Dixon, Richard W.; Choquette, Deborah; Blanchard, Jill; Kluge, Arthur; Pal, Kollol; Bockovich, Nicholas; Come, Jon; Hediger, Mark
 PATENT ASSIGNEE(S): Basf Aktiengesellschaft, Germany
 SOURCE: PCT Int. Appl., 314 pp.
 CODEN: PIXXD2
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 2
 PATENT INFORMATION:


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2001016300	A2	20010308	WO 2000-US23473	20000825
WO 2001016300	A3	20020530		
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM				
RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG				
EP 1226237	A2	20020731	EP 2000-959449	20000825
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL				
PRIORITY APPLN. INFO.: US 1999-172215P P 19990831 WO 2000-US23473 W 20000825				

OTHER SOURCE(S): MARPAT 134:218936
 AB The present invention relates to polypeptides which comprise the ligand binding domain of CDC25, crystalline forms of these polypeptides, and the use of these crystalline forms to determine the 3-dimensional structure of the catalytic domain of CDC25 alone and in complexes with pentapeptide inhibitors. Atomic coordinates are provided from x-ray diffraction of crystals of CDC25A and CDC25B catalytic domains in the presence and absence of various inhibitors. The invention also relates to the use of the 3-dimensional structure of the CDC25 catalytic domain in methods of designing and/or identifying potential inhibitors of CDC25 activity, for example, compounds which inhibit the binding of a native substrate to the CDC25 catalytic domain. The method comprises the steps of (1) identifying one or more functional groups capable of interacting with one or more subites of the CDC25 catalytic domain, and (2) identifying a scaffold which presents the functional group or functional groups in a suitable orientation for interacting with one or more subites of the CDC25 catalytic domain. Since CDC25 is a potential target for therapies aimed at controlling proliferative disease, the atomic coordinates allow rational structure-based design of potential agents for the treatment of cancer, restenosis,

L4 ANSWER 108 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)
 reocclusion of coronary artery, or inflammation.
 IT 329274-06-8P
 RL: SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)
 (crystal structure of CDC25 proteins and its use in rational design of inhibitors)
 RN 329274-06-8 CAPLUS
 CN L-Norvalinamide, N-(1-dibenzofuran-2-yl)-4-(sulfomethyl)-L-phenylalanine-L-norvalyl-2-methyl-L-prolyl-3-benzo[b]thien-3-yl-L-alanyl-5-carboxy-N-(1,1-dimethylethyl)- (9CI) (CA INDEX NAME)
 Absolute stereochemistry.



PAGE 1-A



PAGE 1-B


L4 ANSWER 109 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
 ACCESSION NUMBER: 2001:152640 CAPLUS
 DOCUMENT NUMBER: 134:208130
 TITLE: Preparation of substituted ureas as cell adhesion inhibitors
 INVENTOR(S): Delazzo, Stephen E.; Haggmann, William K.; Kamenecka, Theodore M.
 PATENT ASSIGNEE(S): Merck & Co., Inc., USA
 SOURCE: PCT Int. Appl., 60 pp.
 CODEN: PIXXD2
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2001014328	A2	20010301	WO 2000-US22437	20000816
WO 2001014328	A3	20020131		
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM				
RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG				
AU 2000069093	A5	20010319	AU 2000-69093	20000816
US 6353099	B1	20020305	US 2000-641408	20000817
PRIORITY APPLN. INFO.: US 1999-150055P P 19990820 WO 2000-US22437 W 20000816				

OTHER SOURCE(S): MARPAT 134:208130
 AB Compds. R1R2NCONR3CR4R5-Y-COR6 (R1, R2 = H, (un)substituted alkyl, alkenyl, alkynyl, cycloalkyl, heterocyclyl, aryl, or heteroaryl or R1R2N form a mono- or bicyclic ring; R3 is any group given for R1/R2 or R2 and R3 together with the atoms to which they are attached form a heterocyclic ring with the proviso that R1 and R2 do not form a ring; R4 = (un)substituted alkyl, aryl, arylalkyl, biaryl, biarylalkyl, heteroaryl, heteroarylalkyl, heteroarylaryl, heteroarylarylalkyl, arylheteroaryl, or arylheteroarylalkyl; R5 = H, (un)substituted alkyl, alkenyl, or alkynyl; R6 = OH, alkoxy, alkenoxy, alkynoxy, aryloxy, arylalkoxy, or an amino group; Y is a bond or CR7R8, where R7 = H, alkyl, alkenyl, alkynyl, aryl, or arylalkyl; R8 is any group given for R7 plus OH, alkoxy, halo, NO2, amino, etc.] were prepared as antagonists of VLA-4 and/or α4β7 and are useful in the inhibition or prevention of cell adhesion and cell-adhesion mediated pathologies. Thus, treating 4-(2-methoxyphenyl)-L-phenylalanine tert-Bu ester (obtained from 4-iodo-L-phenylalanine and 2-methoxyphenylboronic acid) with pyrrolidine and p-nitrophenyl chloroformate in CH2Cl2 containing diisopropylethylamine and ester cleavage with 50% TFA/CH2Cl2 afforded N-(1-pyrrolidinylcarbonyl)-4-(2-methoxyphenyl)-L-phenylalanine.

IT 328257-46-1P
 RL: SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)
 (preparation of substituted ureas as cell adhesion inhibitors)
 RN 328257-46-1 CAPLUS
 CN [1,1'-Biphenyl]-4-propanoic acid, 2'-cyano-α-[[[1,2-dihydro-1-(methylsulfonyl)spiro[3H-indole-3,4'-piperidin]-1'-yl]carbonyl]amino]-, (αS)- (9CI) (CA INDEX NAME)

L4 ANSWER 109 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)
 Absolute stereochemistry.



L4 ANSWER 110 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
 ACCESSION NUMBER: 2001:137478 CAPLUS
 DOCUMENT NUMBER: 134:188233
 TITLE: Melanocortin metalloprotein constructs, combinatorial libraries, and applications
 INVENTOR(S): Sharma, Shubh D.; Shi, Yi-Qun; Yang, Wei; Cai, Hui-Zhi
 PATENT ASSIGNEE(S): Palatin Technologies, Inc., USA
 SOURCE: PCT Int. Appl., 80 pp.
 CODEN: PIXXD2
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2001013112	A1	20010222	WO 2000-US16396	20000615
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, GR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MY, NZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, AM, AZ, BY, BG, KZ, MD, RU, TJ, TM			
RW:	GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, CA, GN, GW, ML, MR, NE, SN, TD, TG			
EP 1208377	A1	20020529	EP 2000-944681	20000615
R:	AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL			
PRIORITY APPL. INFO.:			US 1999-148994P	P 19990812
			WO 2000-US16396	W 20000615

OTHER SOURCE(S): MARPAT 134:188233
 AB Metallopeptides and metalloprotein combinatorial libraries specific for melanocortin receptors are provided, for use in biol., pharmaceutical and related applications. The metallopeptides and combinatorial libraries are made of peptides, peptidomimetic and peptide-like constructs, in which the peptide, peptidomimetic or construct is conformationally fixed on complexation of a metal ion-binding portion thereof with a metal ion.
 IT 327605-06-1P
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)
 (melanocortin metalloprotein constructs, combinatorial libraries, and applications)
 RN 327605-06-1 CAPLUS
 CN Rhenium, [(2R)-1-acetyl-2-piperidinecarbonyl- κ N1-glycyl- κ N-L-cysteinyl- κ N, κ S-L-phenylalaninamidato(3-)]oxo-, conjugate monoacid, (SP-5-24)- (9CI) (CA INDEX NAME)


L4 ANSWER 111 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
 ACCESSION NUMBER: 2001:111905 CAPLUS
 DOCUMENT NUMBER: 134:291367
 TITLE: A Kinetic Study into the Hydrolysis of the Ochrotoxins and Analogues by Carboxypeptidase A
 AUTHOR(S): Stander, Maria A.; Steyn, Pieter S.; van der Westhuizen, Francois H.; Payne, Barry E.
 CORPORATE SOURCE: Potchefstroom University for Christian Higher Education, Potchefstroom, S. Afr.
 SOURCE: Chemical Research in Toxicology (2001), 14 (3), 302-304
 CODEN: CRTOC; ISSN: 0893-228X
 PUBLISHER: American Chemical Society
 DOCUMENT TYPE: Journal
 LANGUAGE: English

AB The hydrolyses of the ochrotoxins and analogs by carboxypeptidase A were assessed. This was done by measuring the amount of phenylalanine formed with liquid chromatog. coupled to tandem electrospray mass spectrometry. The kinetic data of ochratoxin A, ochratoxin B, and the synthetic bromo-ochratoxin B were compared to the values of a number of synthesized structure analogs, namely, ochratoxin A Me ester, ochratoxin B Me ester, N-(2-hydroxybenzoyl)phenylalanine, N-(5-chloro-2-hydroxybenzoyl)phenylalanine, N-(5-bromo-2-hydroxybenzoyl)phenylalanine, and N-(5-fluoro-2-hydroxybenzoyl)phenylalanine. The halogen-containing analogs had lower turnovers than their des-halo analogs. There are no substantial differences in the kinetic data between the different halogen-containing analogs.

IT 255042-26-3
 RL: BPR (Biological process); BSU (Biological study, unclassified); BIOL (Biological study); PROC (Process)
 (a kinetic study into the hydrolysis of the ochrotoxins and analogs by carboxypeptidase A)


RN 255042-26-3 CAPLUS
 CN L-Phenylalanine, N-[(3R)-5-bromo-3,4-dihydro-8-hydroxy-3-methyl-1-oxo-1H-2-benzopyran-7-yl]carbonyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



REFERENCE COUNT: 22 THERE ARE 22 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 110 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)




● H⁺

REFERENCE COUNT: 5 THERE ARE 5 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 112 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
 ACCESSION NUMBER: 2001:101132 CAPLUS
 DOCUMENT NUMBER: 134:163037
 TITLE: Preparation of 4-[[imidazolylmethyl]amino]chroman-8-carboxamides as protein farnesyltransferase inhibitors
 INVENTOR(S): Baudoin, Bernard; Jimonet, Patrick; Maignan, Sebastien; Achard, Daniel; Mailliet, Patrick; Laoui, Abdelazize; Nemecek, Conception
 PATENT ASSIGNEE(S): Aventis Pharma S.A., Fr.
 SOURCE: PCT Int. Appl., 137 pp.
 CODEN: PIXXD2
 DOCUMENT TYPE: Patent
 LANGUAGE: French
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2001009125	A1	20010208	WO 2000-FR2188	20000728
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, GR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW, AM, AZ, BY, BG, KZ, MD, RU, TJ, TM			
RW:	GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, CA, GN, GW, ML, MR, NE, SN, TD, TG			
FR 2796947	A1	20010202	FR 1999-9892	19990730
PRIORITY APPL. INFO.:			FR 1999-9892	A 19990730
OTHER SOURCE(S):			MARPAT 134:163037	




AB R1R2N2CONR6CR4R5R10 [I; R1 = Z1Z2Z3Z4R9; R2 = H, alkyl, alkanoyl; R4 = (CHR11)nR14; R5 = H, COR15, Z3R16, Z3OCOR; R4R5 = atoms to complete a ring; R, R11 = H, alkyl, aryl; R6 = H or (hydroxy)alkyl; R4R6 = atoms to complete a ring; R9 = H, halo, alkyl, aryl, etc.; R10 = H or Z3OR; R14 = OR, CO2R, alkyl, (hetero)aryl, etc.; R15 = OH, (alkyl)amino, alkyl, alkoxy; R16 = H, halo, OH; Z = (un)substituted 3,4-dihydro-2H-1-benzopyran-4,8-diyl; Z1, Z3 = alkylene; Z2 = imidazolediy; Z4 = (hetero)arylene; n = 0-5) were prepared. Thus, 4-chromanone was converted in 3 steps to 4-(Bocamino)chroman-8-carboxylic acid which was amidated by 4-pyridinemethanamine and the chromatog. resolved product deprotected to give (-) and (+)-4-amino-N-(4-pyridylmethyl)chroman-8-carboxamide. The latter enantiomer was reductively condensed with 1-

L4 ANSWER 112 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)
biphenylmethylimidazole-5-carboxaldehyde (prepn. given) to give title
compd. (+)-II. Data for biol. activity of I were given.

IT 325151-75-5P
RL: BAC (Biological activity or effector, except adverse); BSU (Biological
study, unclassified); SPN (Synthetic preparation); THU (Therapeutic use);
BIOL (Biological study); PREP (Preparation); USES (Uses)
(preparation of 4-((imidazolymethyl)amino)chroman-8-carboxamides as protein
farnesyltransferase inhibitors)

RN 325151-75-5 CAPLUS
CN L-Phenylalanine, N-[[4-[[[1-(1,1'-biphenyl)-4-ylmethyl]-1H-imidazol-5-
yl]methyl]amino]-3,4-dihydro-2H-1-benzopyran-6-yl]carbonyl]-4-
(hydroxymethyl)-, ethyl ester (9CI) (CA INDEX NAME)

Absolute stereochemistry.




REFERENCE COUNT: 2 THERE ARE 2 CITED REFERENCES AVAILABLE FOR THIS
RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 113 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)
compd. (+)-II. Data for biol. activity of I were given.

IT 325144-37-4P
RL: BAC (Biological activity or effector, except adverse); BSU (Biological
study, unclassified); PUR (Purification or recovery); SPN (Synthetic
preparation); THU (Therapeutic use); BIOL (Biological study); PREP
(Preparation); USES (Uses)
(preparation of 4-((imidazolymethyl)amino)chroman-8-carboxamides as protein
farnesyltransferase inhibitors)

RN 325144-37-4 CAPLUS
CN 2H-1-Benzopyran-8-carboxamide, 3,4-dihydro-N-((1S)-1-((4-
methoxyphenyl)methyl)-2-(methylamino)-2-oxoethyl)-4-[[[1-[[4-(5-
thiazolyl)phenyl]methyl]-1H-imidazol-5-yl]methyl]amino]-, (4S)- (9CI) (CA
INDEX NAME)


Absolute stereochemistry.



REFERENCE COUNT: 3 THERE ARE 3 CITED REFERENCES AVAILABLE FOR THIS
RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 113 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
2001:101131 CAPLUS
ACCESSION NUMBER: 134:163036
DOCUMENT NUMBER: Preparation of 4-((imidazolymethyl)amino)chroman-8-
carboxamides as protein farnesyl transferase
inhibitors
TITLE: Baudoin, Bernard; Jimonet, Patrick
INVENTOR(S): Aventis Pharma S.A., Fr.
PCT Int. Appl., 71 pp.
SOURCE: CODEN: PIXXD2
DOCUMENT TYPE: Patent
LANGUAGE: French
FAMILY ACC. NUM. COUNT: 1
PATENT INFORMATION:


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2001009124	A1	20010208	WO 2000-FR2187	20000728
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM			
RW:	GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, CA, GN, GW, ML, MR, NE, SN, TD, TG			
FR 2796946	A1	20010202	FR 1999-9891	19990730
PRIORITY APPLN. INFO.:			FR 1999-9891	A 19990730
OTHER SOURCE(S):		MARPAT 134:163036		
GI				



AB R9C6H4CH2ZCH2NR2Z1CONHCH(CH2R4)CONHMe [I; R2 = H, alkyl, alkanoyl; R4 = (un)substituted aryl; R9 = halo, alkyl, (hetero)aryl, Z2COR12, etc.; R12 = OH, alkoxy, NH2; Z = imidazole-1,5-diyl; Z1 = (un)substituted 3,4-dihydro-2H-1-benzopyran-4,8-diyl; Z2 = alkylene] were prepared. Thus, 4-chromanone was converted in 3 steps to 4-(t-Boc-amino)chroman-8-carboxylic acid which was amidated by (S)-4-ClC6H4CH(NH2)CO2Me and the product amidated by MeNH2 to give, after deprotection, 4-amino-N-((S)-1-methylcarbamoyl-2-(4-chlorophenyl)ethyl)chroman-8-carboxamide. The latter was reductively condensed with 1-(4-methylbenzyl)imidazole-5-carboxaldehyde (preparation given) to give title

L4 ANSWER 114 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
2001:101119 CAPLUS
ACCESSION NUMBER: 134:163035
DOCUMENT NUMBER: Preparation of 4-((imidazolymethyl)amino)chroman-8-
carboxamides as protein farnesyltransferase inhibitors
TITLE: Baudoin, Bernard; Jimonet, Patrick; Laoui, Abdelaziz
INVENTOR(S): Aventis Pharma S.A., Fr.
PCT Int. Appl., 54 pp.
SOURCE: CODEN: PIXXD2
DOCUMENT TYPE: Patent
LANGUAGE: French
FAMILY ACC. NUM. COUNT: 1
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2001009112	A1	20010208	WO 2000-FR2189	20000728
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM			
RW:	GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, CA, GN, GW, ML, MR, NE, SN, TD, TG			
FR 2796948	A1	20010202	FR 1999-9895	19990730
PRIORITY APPLN. INFO.:			FR 1999-9895	A 19990730
OTHER SOURCE(S):		MARPAT 134:163035		
GI				




AB R9C6H4CH2ZCH2NR2Z1CONHCH(R4)CONHMe [I; R2 = H, alkyl, alkanoyl; R4 = CHR13R14; R5 = COR15; R9 = Z2Z3R12; R12, R15 = OH, alkoxy, NH2; R13, R14 = H or (un)substituted aryl; Z = imidazole-1,5-diyl; Z1 = (un)substituted 3,4-dihydro-2H-1-benzopyran-4,8-diyl; Z2 = alk(en)ylene; Z3 = bond or CO] were prepared. Thus, 4-chromanone was converted in 3 steps to 4-(Boc-amino)chroman-8-carboxylic acid which was amidated by (S)-4-ClC6H4CH(NH2)CO2Me and the product amidated by NH3 to give, after deprotection, 4-amino-N-((S)-1-carbamoyl-2-(4-chlorophenyl)ethyl)chroman-8-carboxamide. The latter was reductively condensed with 1-(4-(2-methoxycarbonylvinyl)benzyl)imidazole-5-carboxaldehyde (preparation given) to give title compound (+)-II (R9 = CH3CO2Me). Data for biol. activity of I were given.

IT 324806-29-3P
RL: BAC (Biological activity or effector, except adverse); BSU (Biological
study, unclassified); SPN (Synthetic preparation); THU (Therapeutic use);

L4 ANSWER 114 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)
 BIOL (Biological study); PREP (Preparation); USES (Uses)
 (prepn. of 4-[[[imidazolylmethyl]amino]chroman-8-carboxamides as protein
 farnesyltransferase inhibitors)
 RN 324806-29-3 CAPLUS
 CN 2-Propenoic acid, 3-[4-[[[5-[[[1S]-2-amino-1-[(4-chlorophenyl)methyl]-
 2-oxoethyl]amino]carbonyl]-3,4-dihydro-2H-1-benzopyran-4-yl]amino)methyl]-
 1H-imidazol-1-yl]methyl]phenyl]-, methyl ester (9CI) (CA INDEX NAME)


Absolute stereochemistry.
 Double bond geometry unknown.



REFERENCE COUNT: 2 THERE ARE 2 CITED REFERENCES AVAILABLE FOR THIS
 RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 115 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
 ACCESSION NUMBER: 2001:2798 CAPLUS
 DOCUMENT NUMBER: 134:193689
 TITLE: Synthesis and fluorescence properties of
 intramolecularly quenched fluorogenic p-nitroanilides
 containing coumarin or quinolinone derivatives as
 fluorophores
 AUTHOR(S): Charitos, C.; Tzougraki, C.; Kokotos, G.
 CORPORATE SOURCE: Department of Chemistry, University of Athens, Athens,
 157 71, Greece
 SOURCE: Journal of Peptide Research (2000), 56(6), 373-381
 CODEN: JPERFA; ISSN: 1397-002X
 PUBLISHER: Munksgaard International Publishers Ltd.
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 OTHER SOURCE(S): CASREACT 134:193689
 AB Nine model intramolecularly quenched fluorogenic substrates (IQFS) of the
 general structure F-Phe-NH-Np, containing coumarin or quinolinone derivs. as
 fluorophores (F) and the p-nitroanilide group (Np) as quencher, were
 synthesized. The study of the fluorescence properties of the substrates
 synthesized and the corresponding fluorophores showed that efficient
 quenching of fluorescence (>89%) was observed in all cases. The combination
 of 7-glutaryl-amido-4-methyl-coumarin (Mec-NH-Glt-OH) or
 7-methoxy-4-coumarylacetic acid (Mca) with the p-nitroanilide group gave
 the best results (97.2 and 98.8% quenching, resp.). These fluorophores
 can be used to convert peptide p-nitroanilides into IQFS, which, retaining
 their chromogenic properties, may be applied in both fluorometric and
 colorimetric assays.
 IT 182944-07-6P
 RL: PRP (Properties); SPN (Synthetic preparation); PREP (Preparation)
 (preparation and fluorescence properties of fluorogenic nitroanilides
 containing
 coumarin or quinolinone derivs. as fluorophores)
 RN 182944-07-6 CAPLUS
 CN 2H-1-Benzopyran-3-carboxamide, N-[[[1S]-2-[(4-nitrophenyl)amino]-2-oxo-1-
 (phenylmethyl)ethyl]-2-oxo- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




REFERENCE COUNT: 16 THERE ARE 16 CITED REFERENCES AVAILABLE FOR THIS
 RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT


L4 ANSWER 116 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
 ACCESSION NUMBER: 2000:894620 CAPLUS
 DOCUMENT NUMBER: 134:141358
 TITLE: Development of the first CGRP-antagonist with
 nanomolar affinity
 AUTHOR(S): Beck-Sicking, Annette G.; Rist, Beate; Enzeroth,
 Michael; Lacroix, Silvain
 CORPORATE SOURCE: Department of Pharmacy, ETH Zurich, Zurich, CH 8057,
 Switz.
 SOURCE: Peptides for the New Millennium, Proceedings of the
 American Peptide Symposium, 16th, Minneapolis, MN,
 United States, June 26-July 1, 1999 (2000), Meeting
 Date 1999, 222-223. Editor(s): Fields, Gregg B.; Tam,
 James P.; Barany, George. Kluwer Academic Publishers:
 Dordrecht, Neth.
 CODEN: 69ATHX
 DOCUMENT TYPE: Conference
 LANGUAGE: English
 AB CGRP 27-37, which binds to human CGRP1-receptors with low affinity, has
 been systematically varied. In a stepwise rational optimization the
 undecapeptides FVPTNVGPFAP and FVPTDVGPFAP have been identified, which
 bind to human CGRP-receptors. The replacement of Ser34 by Pro has turned
 out to be crucial for the increase of affinity. Interestingly, neither
 hydroxyproline (Hyp), nor homoproline (Hpr) could fully replace Pro34,
 whereas Aib and Tic were only slightly less active. The increase of
 affinity of single mutations has been additive and correlated with the
 decrease of the min. at $\lambda = 220$ nm by CD spectroscopy. FVPTDVGPFAP
 and FVPTNVGPFAP showed exclusively antagonistic properties in the rat
 vasodilatation assay. Interestingly, the duration of the potency of both
 compds. varied significantly. Whereas FVPTDVGPFAP lost potency after 30
 min, analogs with replacement of Asp31 by Asn31 were active for more than
 2 h. Since both ligands exhibit receptor binding affinities in the same
 range ($K_i = 19/14$ nM), the authors suggested that the analog with Asp31 is
 more rapidly metabolized, perhaps because of an increased susceptibility
 for proteases. This effect could be confirmed by preliminary results with
 other analogs containing either Asp or Asn in position 31. In this case as
 well, the effect of the Asp containing peptides was significantly increased.
 This suggests differences in the metabolism of both compds. and could be
 important for drug design.
 IT 324035-60-1
 RL: BAC (Biological activity or effector, except adverse); BPR (Biological
 process); BSU (Biological study, unclassified); PRP (Properties); BIOL
 (Biological study); PROC (Process)
 (development of first CGRP-antagonist with nanomolar affinity)
 RN 324035-60-1 CAPLUS
 CN L-Phenylalanine, L-phenylalanyl-L-valyl-L-prolyl-L-threonyl-L-
 aspartyl-L-valylglycyl-(3S)-1,2,3,4-tetrahydro-3-isquinolinecarbonyl-L-
 phenylalanyl-L-alanyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

L4 ANSWER 116 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)




REFERENCE COUNT: 3 THERE ARE 3 CITED REFERENCES AVAILABLE FOR THIS
 RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT



L4 ANSWER 119 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
 ACCESSION NUMBER: 2000:830796 CAPLUS
 DOCUMENT NUMBER: 134:101182
 TITLE: Extended TIP(P) analogs as precursors for labeled
 8-opioid receptor ligands
 AUTHOR(S): Kumar, Vivek, Murray, Thomas F.; Aldrich, Jane V.
 CORPORATE SOURCE: Department of Pharmaceutical Sciences School of
 Pharmacy, University of Maryland, Baltimore, MD,
 21201, USA
 SOURCE: Journal of Medicinal Chemistry (2000), 43(26),
 5050-5054
 CODEN: JMCMAR; ISSN: 0022-2623
 PUBLISHER: American Chemical Society
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 AB Peptides H-Tyr-Tic-Phe-Phe-OH (TIPP) and the shorter H-Tyr-Tic-Phe-OH
 (TIP) are potent and highly selective antagonists at the 8-opioid
 receptor and, therefore, are ideal candidates for the attachment of labels
 to assist in the study of 8-opioid receptors. Peptides extended at
 the C-terminus with residues (i.e., Asp, Asn, Glu, Gln and Lys, which can
 be used as handles for further modification and/or labeling) were
 synthesized. The TIPP-D/L-Asx/Glx derivs. (Asx = Asp or Asn; Glx = Glu or
 Gln) exhibited similar 8-receptor affinity to TIPP (K_i = 5-10 nM vs
 K_i = 6 nM), and neither the location of the carboxylic acid moiety nor the
 stereochem. of the C-terminal residue significantly affected the
 8-receptor affinity of these derivs. Extension of TIPP with an
 addnl. residue did not increase μ-receptor affinity, even though the
 position of the acidic group, which imparts 8-receptor selectivity
 to TIPP, was shifted relative to the carboxylic acid moiety of TIPP. The
 8-receptor affinities of the TIP-D/L-Asx/Glx derivs. were found to
 be influenced mainly by the position of the carboxylic acid function
 rather than the stereochem. of the C-terminal residue.
 TIP(P)-D/L-Lys(Ac)-OH derivs. exhibited moderate 8-receptor affinity
 (K_i = 16-28 nM). The most potent compds. found in the extended
 TIP(P) series were TIPP-D-Gln-OH and TIP-D-Gln-OH (K_i = 5 nM),
 which had similar affinities to TIPP.
 IT 319906-09-7P
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological
 study, unclassified); SPN (Synthetic preparation); BIOL (Biological
 study); PREP (Preparation)
 (preparation and biol. activity of TIP(P) analogs, extended at the
 C-terminus, as precursors for labeled 8-opioid receptor ligands)
 RN 319906-09-7 CAPLUS
 CN L-α-Asparagine, L-tyrosyl-(3S)-1,2,3,4-tetrahydro-3-
 isoquinolinecarboxyl-L-phenylalanyl-L-phenylalanyl- (9CI) (CA INDEX NAME)
 Absolute stereochemistry.


L4 ANSWER 119 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)



REFERENCE COUNT: 41 THERE ARE 41 CITED REFERENCES AVAILABLE FOR THIS
 RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 120 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
 ACCESSION NUMBER: 2000:824283 CAPLUS
 DOCUMENT NUMBER: 134:5161
 TITLE: Preparation of phosphonic and carboxylic acid
 derivatives as inhibitors of protein tyrosine
 phosphatase-1B (PTP-1B)
 INVENTOR(S): LeBlanc, Yves; Dufresne, Claude; Roy, Patrick; Wang,
 Zhaoyin
 PATENT ASSIGNEE(S): Merck Frosst Canada & Co., Can.
 SOURCE: PCT Int. Appl., 62 pp.
 CODEN: PIXXD2
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2000069889	A1	20001123	WO 2000-CA567	20000512
W:	AB, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EP, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM			
RW:	GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GH, GW, ML, MR, NE, SN, TD, TG			
EP 1181109	A1	20020227	EP 2000-929175	20000512
R:	AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO			
US 6365592	B1	20020402	US 2000-570092	20000512
PRIORITY APPLN. INFO.:			US 1999-134150P	P 19990514
			US 1999-158778P	P 19991012
			WO 2000-CA567	W 20000512
OTHER SOURCE(S):	MARPAT 134:5161			
GI				




AB Peptides I [B1a, B1b, B2a, B2b = CF₂PO₃H₂ or CF₂CO₂H or one of B1a, B1b, B2a, and B2b is H and the others are H, alkyl, heteroaryl, carbocyclyl, aryl, OH, halo, CHF₂, CF₃, CF₂CO₂H, CH₂PO₃H₂, SO₂NH₂, etc.; X = OH, NH₂; Y = H, alkyl, R1-Z-CO (R1 = alkyl, fluoroalkyl, aryl, heteroaryl, etc.; Z = O, SCH₂, SOCH₂, SO₂CH₂, substituted imino, or CH₂CH), acyl residue of an amino acid which may be substituted, alkyl- or arylsulfonyl] were prepared as inhibitors of PTP-1B. Thus, (4S)-5-[[[(1S)-2-[[[(1S)-2-amino-1-[[4-(difluoro(phosphono)methyl)benzyl]-2-oxoethyl]amino]-1-[[4-(carboxy(difluoromethyl)benzyl]-2-oxoethyl]amino]-4-(benzoylamino)-5-oxopentanoic acid was prepared by the solid-phase method using a TentaGel RAM resin.

L4 ANSWER 120 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)

IT 307990-98-3P
 RL: SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)
 (preparation of phosphonic and carboxylic acid derivs. as inhibitors of protein tyrosine phosphatase-1B)
 RN 307990-98-3 CAPLUS
 CN L-Phenylalaninamide, 4-(carboxy(difluoromethyl)-N-(3-quinolinylcarbonyl)-L-phenylalanyl-4-(difluorophosphonomethyl)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




REFERENCE COUNT: 6 THERE ARE 6 CITED REFERENCES AVAILABLE FOR THIS
 RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 121 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
 ACCESSION NUMBER: 2000:755247 CAPLUS
 DOCUMENT NUMBER: 133:122122
 TITLE: Enantiopure reagents and process for the separation of amino acid enantiomers
 INVENTOR(S): Delplanche, Thierry; Callens, Roland
 PATENT ASSIGNEE(S): Solvay (Societe Anonyme), Belg.
 SOURCE: Eur. Pat. Appl., 18 pp.
 CODEN: EPXXDW
 DOCUMENT TYPE: Patent
 LANGUAGE: French
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
EP 1046627	A2	20001025	EP 2000-201285	20000410
EP 1046627	A3	20001102		
EP 1046627	B1	20040211		
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO				
BE 1012622	A3	20010109	BE 1999-280	19990421
AT 259339	E	20040215	AT 2000-201285	20000410
CA 2305944	AA	20001021	CA 2000-2305944	20000418
JP 2000327594	A2	20001128	JP 2000-120313	20000421

PRIORITY APPLN. INFO.: BE 1999-280 A 19990421
 GI



AB Enantiopure reagents, e.g. I, were prepared and used in resolution of racemic amino acids in presence of NRE3.
 IT 302837-74-7P
 RL: PUR (Purification or recovery); SPN (Synthetic preparation); PREP (Preparation)
 (preparation of enantiopure reagents and process for the resolution of amino acids)
 RN 302837-74-7 CAPLUS
 CN 3-Isquinolinecarboxylic acid, 1,2,3,4-tetrahydro-2-[[[(1S)-2-(2-methoxyethoxy)-2-oxo-1-(phenylmethyl)ethyl]amino]carbonyl]-, (1S)- (9CI)
 (CA INDEX NAME)



Absolute stereochemistry.

L4 ANSWER 122 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
 ACCESSION NUMBER: 2000:752379 CAPLUS
 DOCUMENT NUMBER: 134:50988
 TITLE: Chiral Resolution, Pharmacological Characterization, and Receptor Docking of the Noncompetitive mGlu1 Receptor Antagonist (±)-2-Hydroxyimino-1a,2-dihydro-1H-7-oxacyclopropa[b]naphthalene-7a-carboxylic Acid Ethyl Ester
 AUTHOR(S): Ott, David; Floersheim, Philipp; Inderbitzin, Werner; Stoehr, Natacha; Francotte, Eric; Lecis, Gabrielle; Richert, Paul; Rihs, Grety; Flor, Peter Josef; Kuhn, Rainer; Gasparini, Fabrizio
 CORPORATE SOURCE: Nervous System Research and Core Technologies, Novartis Pharma AG, Basel, CH-4002, Switz.
 SOURCE: Journal of Medicinal Chemistry (2000), 43(23), 4428-4436
 CODEN: JMCMAR; ISSN: 0022-2623
 PUBLISHER: American Chemical Society
 DOCUMENT TYPE: Journal
 LANGUAGE: English

AB Racemic CPCCOEt ((1aRS,7aRS)-2-hydroxyimino-1a,2-dihydro-1H-7-oxacyclopropa[b]naphthalene-7a-carboxylic acid Et ester, (±)-1) derive. have been shown to be subtype-selective metabotropic glutamate (mGlu) 1 receptor antagonists (Annoura et al. Bioorg. Med. Chemical Lett. 1996, 6, 763-766). The optical isomers of (±)-1 have been separated by chromatog. on a chiral stationary phase. The absolute configuration at the C-1a and C-7a positions was determined using x-ray crystallog. of an amide derivative with the Me ester of L-phenylalanine (L-PheOMe) ((+)-6). In a phosphoinositol (PI) turnover assay at the cloned human mGlu1b receptor, (-)-1 and the new amide derivs. (-)-5 and (-)-6, all of which have (1aS,7aS)-stereochem. on the chromane ring system, showed IC50 values of 1.5, 0.43, and 0.93 μM, resp. In contrast, (+)-1 and the new amide derivs. (+)-5 and (+)-6 were found to be inactive up to a concentration of 30 μM indicating a selectivity for the (-)-enantiomers of at least 70-fold. In a previous study (Litschig et al. Mol. Pharmacol. 1999, 55, 453-461) we demonstrated using site-directed mutagenesis that the interaction site of (±)-1 is located in the transmembrane (TM) domain of hmGlu1b. To suggest a plausible binding mode of (-)-1, we have built a mol. mechanics model of the putative seven TM domain of hmGlu1 based on the α-carbon template of the TM helices of rhodopsin. A receptor docking hypothesis suggests that the OH of T815 (TMVII) comes in close contact with the oxime OH of (-)-1 and (-)-5, whereas no such close interactions could be demonstrated by docking of (+)-1.

IT 314021-52-8P
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); PRP (Properties); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation)
 (chiral resolution, pharmacol. characterization, and receptor docking of the noncompetitive mGlu1 receptor antagonist (±)-2-hydroxyimino-1a,2-dihydro-1H-7-oxacyclopropa[b]naphthalene-7a-carboxylic acid Et ester)
 RN 314021-52-8 CAPLUS
 CN L-Phenylalanine, N-[[7,7a-dihydro-7-(hydroxyimino)benzo[b]cyclopropa[e]pyr an-1a(1H)-yl]carbonyl]-, methyl ester (9CI) (CA INDEX NAME)

Absolute stereochemistry.
 Double bond geometry unknown.





REFERENCE COUNT: 23 THERE ARE 23 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 123 OF 261 CAPLUS COPYRIGHT 2004 ACS ON STN
 ACCESSION NUMBER: 2000:707153 CAPLUS
 DOCUMENT NUMBER: 133:266851
 TITLE: Preparation of 4-(imidazolin-2-yl)quinolines as NK-3 and/or GABAA receptor ligands
 INVENTOR(S): Yuan, Jun; Maynard, George D.; Hutchison, Alan; Rachwal, Stanislaw
 PATENT ASSIGNEE(S): Neurogen Corporation, USA
 SOURCE: PCT Int. Appl., 71 pp.
 CODEM: PIXXD2
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 2
 PATENT INFORMATION:


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2000058303	A1	20001005	WO 2000-US8205	20000328
WO 2000058303	C2	20021219		
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM			
RW:	GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG			
EP 1165542	A1	20020102	EP 2000-919753	20000328
EP 1165542	B1	20030820		
R:	AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO			
US 6413982	B1	20020702	US 2000-536922	20000328
JP 20020540203	T2	20021126	JP 2000-608005	20000328
AT 247652	E	20030915	AT 2000-919753	20000328
US 2002198232	A1	20021226	US 2002-140693	20020507
US 6624175	B2	20030923		

PRIORITY APPLN. INFO.:
 US 1999-126926P P 19990329
 US 2000-536922 A1 20000328
 WO 2000-US8205 W 20000328

OTHER SOURCE(S): MARPAT 133:266851
 GI



L4 ANSWER 123 OF 261 CAPLUS COPYRIGHT 2004 ACS ON STN (Continued)
 AB The title compds. [I; R1 = H, halo, OH, etc.; R2, R3 = (un)substituted alkyl, Ph, naphthyl, etc.; R4 = H, halo, OH, etc.; X = NH, O, N(alkyl), Y1 = CR11R12, CR11R12(CH2)p, (CH2)pCR11R12, (CH2)pCO (wherein p = 0-2; R11, R12 = H, (un)substituted Ph, naphthyl, etc.); Y2 = CR11R12, CO (with the proviso that Y2 is not CO when Y1 = (CH2)pCO)] which bind with high affinity to NK-3 receptors and/or GABAA receptors, and therefore are useful in treating patients suffering from certain central nervous system and peripheral diseases or disorders, were prepared. E.g., a multi-step synthesis of imidazoline II which showed IC50 of 19 nM against NK-3 receptor binding, was given. This invention also relates to the use of compds. I in combination with one or more other CNS agents to potentiate the effects of the other CNS agents. The compds. I are also useful as probes for the localization of NK-3 receptors and GABAA receptors.
 IT 298689-34-6P
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)
 [preparation of 4-(imidazolin-2-yl)quinolines as NK-3 and/or GABAA receptor ligands]
 RN 298689-34-6 CAPLUS
 CN 4-Quinolincarboxamide, N-[1-(aminocarbonyl)-1-phenylpropyl]-3-methoxy-2-phenyl- (9CI) (CA INDEX NAME)



REFERENCE COUNT: 4 THERE ARE 4 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 124 OF 261 CAPLUS COPYRIGHT 2004 ACS ON STN
 ACCESSION NUMBER: 2000:645993 CAPLUS
 DOCUMENT NUMBER: 133:238324
 TITLE: Preparation of tyrosine amides and analogs as protein tyrosine phosphatase inhibitors
 INVENTOR(S): Larsen, Scott D.; May, Paul D.; Bleasdale, John E.; Liljebriis, Charlotta; Schostarez, Heinrich Josef; Barf, Tjeerd; Nilsson, Marianne
 PATENT ASSIGNEE(S): Pharmacia and Upjohn AB, Swed.
 SOURCE: PCT Int. Appl., 124 pp.
 CODEM: PIXXD2
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 3
 PATENT INFORMATION:


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2000053583	A1	20000914	WO 2000-US6022	20000309
W:	AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM			
RW:	GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG			
US 6410585	B1	20020625	US 1999-265410	19990310
EP 1161421	A1	20011212	EP 2000-917793	20000309
R:	AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO			
JP 2002539115	T2	20021119	JP 2000-604023	20000309

PRIORITY APPLN. INFO.:
 US 1999-265410 A 19990310
 US 1997-57730P P 19970828
 US 1998-138642 A2 19980824
 WO 2000-US6022 W 20000309


OTHER SOURCE(S): MARPAT 133:238324
 GI

L4 ANSWER 124 OF 261 CAPLUS COPYRIGHT 2004 ACS ON STN (Continued)
 R1 = H, CH2OH, alkylcarbamoyl, etc.; R2 = H or Me; R4 = H or (phenyl)alkyl; Z = (un)substituted 1,4-phenylene; Z1 = CO or SO2 were prepd. Thus, (S)-Me2CO2C(NHCH(CO2H)CH2C6H3(OH)I-4,3 was amidated by Ph(CH2)4NH2 and the product converted in 5 steps to title compd. II. Data for biol. activity of I were given.
 IT 292834-82-3P
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)
 [preparation of tyrosine amides and analogs as protein tyrosine phosphatase inhibitors]
 RN 292834-82-3 CAPLUS
 CN Benzoic acid, 2-(carboxymethoxy)-5-[(2S)-3-oxo-3-[(4-phenylbutyl)amino]-2-[(3-quinolinylcarbonyl)amino]propyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




REFERENCE COUNT: 28 THERE ARE 28 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT



AB R2CH2CR1R2NHZIR3 [I; R = OSO3H, OCH2CO2R4, OCH2CONHOH, N(CH2CO2R4)2, etc.;

L4 ANSWER 125 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
 ACCESSION NUMBER: 2000:609236 CAPLUS
 DOCUMENT NUMBER: 133:310131
 TITLE: Solid-phase synthesis of 1,2,3,4-tetrahydro- β -carboline-containing peptidomimetics
 AUTHOR(S): Li, Xianfeng; Zhang, Lianshan; Zhang, Wei; Hall, Steven E.; Tam, James P.
 CORPORATE SOURCE: Sphinx Pharmaceuticals A Division of Eli Lilly and Company, Cambridge, MA, 02139, USA
 SOURCE: Organic Letters (2000), 2(20), 3075-3078
 CODEN: ORLEF7; ISSN: 1523-7050
 PUBLISHER: American Chemical Society
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 OTHER SOURCE(S): CASREACT 133:310131
 GI




AB A solid-phase method for the synthesis of 1,2,3,4-tetrahydro- β -carboline-containing peptidomimetics I (AA1-AA4 = Ala, Leu, Phe, Pro, Val, Asp, Gly, etc.) has been developed. The key step in the strategy is the Pictet-Spengler condensation of a resin-bound tryptophan-containing fragment with an Fmoc-amino aldehyde.

IT 301850-18-0P
 RL: SPN (Synthetic preparation); PREP (Preparation)
 (solid-phase synthesis of tetrahydro- β -carboline-containing peptidomimetics)
 RN 301850-18-0 CAPLUS
 CN L-Alanine, 2,3,4,9-tetrahydro-1-[[[L-valyl-L-prolyl]amino]methyl]-1H-pyrido[3,4-b]indole-3-carbonyl-L-phenylalanyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

L4 ANSWER 125 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)





REFERENCE COUNT: 6 THERE ARE 6 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 126 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
 ACCESSION NUMBER: 2000:568542 CAPLUS
 DOCUMENT NUMBER: 133:150464
 TITLE: Preparation of quinolinylindole derivatives and compositions in use as antimicrobial agents
 INVENTOR(S): Cuny, Gregory D.; Hauske, James R.; Heefner, Donald L.; Hoemann, Michael Z.; Kumaravel, Gnanasambandam; Melikian-Badalian, Anita; Rossi, Richard F.; Xie, Roger L.
 PATENT ASSIGNEE(S): Sepracor, Inc., USA
 SOURCE: U.S., 228 pp., Cont.-in-part of U.S. Ser. No. 99,640.
 CODEN: USXXAM
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 7
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 6103905	A	20000815	US 1998-213385	19981211
US 6207679	B1	20010327	US 1998-45051	19980319
US 6172084	B1	20010109	US 1998-99640	19980618
WO 2000034265	A2	20000615	WO 1999-US28744	19991203
WO 2000034265	A3	20021003		

W: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM
 RW: GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LJ, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG
 US 6376670 B1 20020423 US 2000-658690 20000908
 PRIORITY APPLN. INFO.: US 1997-878781 B2 19970619
 US 1998-45051 A2 19980319
 US 1998-99640 A2 19980618
 US 1998-213385 A 19981211
 US 2000-639622 A2 20000815
 OTHER SOURCE(S): MARPAT 133:150464
 GI

L4 ANSWER 126 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)




AB Title compds. I; Q = hydrophobic group, H; X = heterocyclyl, amidinyl, formamidinyl, guanidinyl, CN, CSNR2, OR, SR; Z = CC, (E)-CH:CH, (Z)-CH:CH, (CH2)2; L = hydrophobic group, H; R represents independently for each occurrence = H, alkyl, heteroalkyl, aryl, heteroaryl, acyl, sulfonyl; R1 = H, alkyl, aryl, 4-CH3C6H4SO2, (CH2)d; d = 1-6; R2 = H, alkyl, aryl; R3 = H, alkyl, aryl; m = 1-8; n = 1-4 and pharmaceutical preps. using title compds. are prepared as antimicrobial agents. The MIC value of I against at least one Gram-pos. bacterium ranged from 0.1-10 μ g/mL. Thus, the title compound II was prepared and has a therapeutic index in primates of at least 10 for the inhibition of infection by at least one Gram-pos. bacterium.

IT 275357-08-9P
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation)
 (preparation of quinolinylindole derivs. as antimicrobial agents)
 RN 275357-08-9 CAPLUS
 CN 4-Quinolinescarboxamide, N-[2-amino-1-[(4-nitrophenyl)methyl]-2-oxoethyl]-2-(5-chloro-1H-indol-3-yl)- (9CI) (CA INDEX NAME)

09/ 964,161


L4 ANSWER 126 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)



REFERENCE COUNT: 39 THERE ARE 39 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 127 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
 ACCESSION NUMBER: 2000:508920 CAPLUS
 DOCUMENT NUMBER: 133:120243
 TITLE: Preparation of guanidinoisoquinolines as urokinase inhibitors
 INVENTOR(S): Dickinson, Roger Peter; Fish, Paul Vincent; Barber, Christopher Gordon
 PATENT ASSIGNEE(S): UK
 SOURCE: U.S., 94 pp.
 CODEN: USXXAM
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 6093731	A	20000725	US 1999-359439	19990722
PRIORITY APPLN. INFO.:			US 1999-359439	19990722
OTHER SOURCE(S):		MARPAT 133:120243		



AB RZZZNR221R3 [R = N:C(NH2)2 or NHC(:NH)NH2; R2 = H, alkyl, (hetero)aryl, etc.; R3 = CO2H, alkoxycarbonyl, CH2OH, CONH2, CH2NH2, etc.; Z = (4-halo)isoquinoline-1,7-diyl; Z1 = (un)substituted (hetero)(cyclo)alkylene or (un)substituted arylene; Z2 = CO, CH2, SO2] were prepared as urokinase inhibitors (no data). Thus, 1,4-dichloroisoquinoline-7-sulfonyl chloride (preparation given) was amidated by H2NCH2CO2Me3 and the product condensed with guanidine to give, after saponification, title compd I.


IT 256476-58-1P
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)
 (preparation of guanidinoisoquinolines as urokinase inhibitors)

RN 256476-58-1 CAPLUS
 CN Benzenesulfonic acid, α-[[[1-[(aminoiminomethyl)amino]-4-chloro-7-isoquinolinyl]carbonyl]amino]-, mono(trifluoroacetate) (9CI) (CA INDEX NAME)

CM 1

L4 ANSWER 127 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)

CRN 256476-57-0
 CMF C19 H16 Cl N5 O3



CM 2


CRN 76-05-1
 CMF C2 H F3 O2



REFERENCE COUNT: 5 THERE ARE 5 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 128 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
 ACCESSION NUMBER: 2000:501827 CAPLUS
 DOCUMENT NUMBER: 133:120332
 TITLE: Preparation of histamines as inhibitors of protein geranylgeranyl transferase I for use of antifungal agents
 INVENTOR(S): Okubo, Mitsuru; Ono, Jun; Asahi, Shuichi; Sagara, Takeshi; Sato, Toshihiko; Morishima, Hajime
 PATENT ASSIGNEE(S): Banyu Pharmaceutical Co., Ltd., Japan
 SOURCE: Jpn. Kokai Tokkyo Koho, 15 pp.
 CODEN: JKXXAF
 DOCUMENT TYPE: Patent
 LANGUAGE: Japanese
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 2000204078	A2	20000725	JP 1999-5249	19990112
PRIORITY APPLN. INFO.:			JP 1999-5249	19990112
OTHER SOURCE(S):		MARPAT 133:120332		




AB Histamines I [X = O, NH; W = CO, CH2; R = nonarom. heterocyclyl, alicyclyl, (un)substituted C7-12 aralkyl; Y = (halo-substituted) aryl; B = H, carbamoyl, amino-C1-3 alkylcarbamoyl], their pharmacol. acceptable salts, or esters are prepared. Amidation of D-2-hydroxy-3-(1-naphthyl)propionic acid with N(im)-tritylhistamine, esterification of the resulting amide with 1-naphthalenecarboxylic acid, and detritylation of the product gave I (X = O, W = CO, R = Y = 1-naphthyl, B = H), which inhibited geranylgeranyl transferase I of Candida albicans with IC50 of 11 nM.

IT 285980-03-2P
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)
 (preparation of histamines as inhibitors of protein geranylgeranyl transferase I for use of antifungal agents)

RN 285980-03-2 CAPLUS
 CN 9H-Xanthene-9-carboxamide, N-[(1R)-1-[(3,5-dichlorophenyl)methyl]-2-[(2-(1H-imidazol-4-yl)ethyl)amino]-2-oxoethyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

L4 ANSWER 128 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)




L4 ANSWER 129 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 2000:456887 CAPLUS
 DOCUMENT NUMBER: 133:89444
 TITLE: Preparation of 4-amino-azepan-3-one protease inhibitors
 INVENTOR(S): Marquis, Robert Wells, Jr.; Ru, Yu; Veber, Daniel Frank; Cummings, Maxwell David; Thompson, Scott Kevin; Yamashita, Dennis
 PATENT ASSIGNEE(S): Smithkline Beecham Corp., USA
 SOURCE: PCT Int. Appl., 273 pp.
 CODEN: PIXXD2
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 4
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2000038687	A1	20000706	WO 1999-US30730	19991221
W: AE, AL, AU, BA, BB, BG, BR, CA, CM, CZ, EE, GE, GR, GM, HR, HU, ID, IL, IN, IS, JP, KP, KR, LC, LK, LR, LT, LV, MA, MG, MK, MN, MX, NO, NZ, PL, RO, SG, SI, SK, SL, TR, TT, UA, US, UZ, VN, YU, ZA, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM				
RW: GH, GM, KE, LG, MW, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CO, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG				
CA 2356671	AA	20000706	CA 1999-2356671	19991221
BR 9916488	A	20011009	BR 1999-16488	19991221
EP 1158986	A1	20011205	EP 1999-963112	19991221
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO				
JP 2002533397	T2	20021008	JP 2000-590640	19991221
AU 768565	B2	20031218	AU 2000-19411	19991221
NZ 511710	A	20031219	NZ 1999-511710	19991221
EP 1384713	A1	20040128	EP 2003-76211	19991221
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, FI, RO, CY				
ZA 2001004208	A	20020523	ZA 2001-4208	20010523
US 2003144175	A1	20030731	US 2001-881334	20010614
NO 2001003124	A	20010622	NO 2001-3124	20010622
US 2002147188	A1	20021010	US 2002-74940	20020213
US 2003044399	A1	20030306	US 2002-74639	20020213
US 2003225061	A1	20031204	US 2003-404142	20030401
US 2004002487	A1	20040101	US 2003-404716	20030401
PRIORITY APPLN. INFO.:			US 1998-113636P	P 19981223
			US 1999-164581P	P 19991110
			EP 1999-963112	A3 19991221
			WO 1999-US30730	W 19991221
			US 2000-593845	B2 20000614
			US 2000-653815	A1 20000901
			US 2001-881334	A1 20010614
			US 2002-74940	A1 20020213

OTHER SOURCE(S): MARPAT 133:89444
 GI

L4 ANSWER 129 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)



L4 ANSWER 130 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 2000:452347 CAPLUS
 DOCUMENT NUMBER: 133:89798
 TITLE: Preparation of peptidyl boronic ester and acid compounds as proteasome inhibitors
 INVENTOR(S): Adams, Julian; Ma, Yu-Ting; Stein, Ross; Baevsky, Matthew; Grenier, Louis; Plamondon, Louis
 PATENT ASSIGNEE(S): Leukosite, Inc., USA
 SOURCE: U.S., 38 pp., Cont.-in-part of U.S. Ser. No. 330,525, abandoned.
 CODEN: USXXAM
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 3
 PATENT INFORMATION:


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 6083903	A	20000704	US 1995-442581	19950516
CA 2203936	AA	19960509	CA 1995-2203936	19951027
WO 9613266	A1	19960509	WO 1995-US14117	19951027
W: AL, AM, AT, AU, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK				
RW: KE, LS, MW, SD, SZ, UG, AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG				
AU 9641398	A1	19960523	AU 1996-41398	19951027
AU 710564	B2	19990923		
ZA 9509119	A	19960527	ZA 1995-9119	19951027
EP 788360	A1	19970813	EP 1995-939670	19951027
EP 788360	B1	20030528		
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LI, LU, MC, NL, PT, SE				
CN 1168633	A	19971224	CN 1995-196590	19951027
US 5780454	A	19980714	US 1995-549318	19951027
JP 10510245	T2	19981006	JP 1995-514834	19951027
NZ 337211	A	20001222	NZ 1995-337211	19951027
IL 115790	A1	20021201	IL 1995-115790	19951027
EP 1312609	A1	20030521	EP 2003-4280	19951027
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE				
AT 241631	E	20030615	AT 1995-939670	19951027
FI 9701746	A	19970606	FI 1997-1746	19970423
NO 9701929	A	19970612	NO 1997-1929	19970425
US 6066730	A	20000523	US 1998-85404	19980526
US 6297217	B1	20011002	US 2000-490511	20000125
US 6465433	B1	20021015	US 2001-953540	20010914
US 2002173488	A1	20021121	US 2002-100295	20020318
US 6548668	B2	20030415		
US 6617317	B1	20030909	US 2002-125997	20020419
US 2003199561	A1	20031023	US 2003-392165	20030319
PRIORITY APPLN. INFO.:			US 1994-330525	B2 19941028
			US 1995-442581	A 19950516
			EP 1995-939670	A3 19951027
			NZ 1995-296717	A1 19951027
			US 1995-549318	A3 19951027
			WO 1995-US14117	W 19951027
			US 1998-85404	A3 19980526
			US 2000-490511	A1 20000125
			US 2001-953540	A1 20010914
			US 2002-100295	A1 20020318

AB The title compds. [I; R1 = COCR13NR11R12, COCR13XR15, COCH2R13; R2 = H, alkyl, cycloalkylalkyl, etc.; R3 = H, alkyl, cycloalkylalkyl, etc.; R4 = H, alkyl, arylalkyl, etc.; R11 = H, alkyl, arylalkyl, etc.; R12 = H, alkyl, cycloalkyl, etc.; R13 = H, alkyl, alkenyl, etc.; R15 = H, alkyl, alkenyl, etc.] which inhibit proteases (no data), including cathepsin K, and are useful for treating diseases of excessive bone loss or cartilage or matrix degradation including osteoporosis, gingival disease including gingivitis and periodontitis, arthritis, more specifically, osteoarthritis and rheumatoid arthritis, Paget's disease, hypercalcemia of malignancy, and metabolic bone disease, were prepared. E.g., a multi-step synthesis of compound II was given. Compds. I are effective at 0.4-400 mg/kg/day.

IT 281217-12-7P
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)
 (preparation of 4-amino-azepan-3-one protease inhibitors)

RN 281217-12-7 CAPLUS
 CN 8-Quinolincarboxamide, N-[(1S)-2-[(hexahydro-3-oxo-1-(2-pyridinylsulfonyl)-1H-azepin-4-yl)amino]-2-oxo-1-(phenylmethyl)ethyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



REFERENCE COUNT: 3 THERE ARE 3 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT


L4 ANSWER 130 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)
US 2002-125997 A1 20020419

OTHER SOURCE(S): MARPAT 133:89798
AB Peptidyl boronic acid and ester compds. P-NRCHR2-X2-CHR3BZ122 [P = 2- or 8-quinolinyl-, 2-quinoxaliny-, 2- or 3-pyridyl-, piperazinyl-, 3-furanyl-, or 3-pyrrolylcarbonyl, or -sulfonyl, or morpholinylcarbonyl; X2 = CONH, CH2NH, CH(OH)CH2, CH(OH)CH(OH), CH(OH)CH2NH, CH:CH, COCH2, SO2NH, SO2CH2, or CH(OH)CH2CONH; R = H or alkyl; R2, R3 = H, alkyl, cycloalkyl, aryl, heterocyclyl, CH2-R5 (R5 = aryl, aralkyl, alkaryl, cycloalkyl, heterocyclyl) or alkyl-chalcogen; Z1, Z2 = alkyl, hydroxy, alkoxy, aryloxy, or together form a dihydroxy compound] were prepared as proteasome inhibitors. Thus, coupling of (1S,2S,3R,5S)-pinanediol leucine boronate trifluoroacetate salt with N-Boc-β-(1-naphthyl)-L-alanine, followed by deprotection, acylation with 4-morpholinylcarbonyl chloride and cleavage of the pinanediol moiety afforded N-(4-morpholine)carbonyl-β-(1-naphthyl)-L-alanine-L-leucine boronic acid [MG-273], which inhibited 20S proteasome with Ki = 0.18 nM.

IT 179324-53-99, 314
RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)
(preparation of peptidyl boronic ester and acid compds. as proteasome inhibitors)

RN 179324-53-9 CAPLUS
CN Boronic acid, [(1R)-3-methyl-1-[(2S)-1-oxo-3-phenyl-2-[(2-quinolinylcarbonyl)amino]propyl]amino]butyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




REFERENCE COUNT: 7 THERE ARE 7 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 131 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
ACCESSION NUMBER: 2000:441762 CAPLUS
DOCUMENT NUMBER: 133:74323
TITLE: Preparation of N-acylphenylalanine derivatives and analogs as inhibitors of α4β1 mediated cell adhesion
INVENTOR(S): Teegarden, Bradley R.; Jayakumar, Honnappa; Matsuki, Kenji; Chrusciel, Robert A.; Fisher, Jed F.; Tanis, Steven P.; Thomas, Edward W.; Blinn, James R.
PATENT ASSIGNEE(S): Tanabe Seiyaku Co., Ltd., Japan; Pharmacia & Upjohn Company
SOURCE: PCT Int. Appl., 215 pp.
CODEN: PIXXD2
DOCUMENT TYPE: Patent
LANGUAGE: English
FAMILY ACC. NUM. COUNT: 1
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2000037429	A2	20000629	WO 1999-US30665	19991220
WO 2000037429	A3	20030522		
W:	AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM			
RW:	GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LJ, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG			
EP 1144365	A2	20011017	EP 1999-966584	19991220
EP 1144365	A3	20030709		
EP 1144365	B1	20040317		
R:	AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO			
JP 2003524614	T2	20030819	JP 2000-589501	19991220
PRIORITY APPLN. INFO.:			US 1998-113501P P	19981222
			WO 1999-US30665 W	19991220

OTHER SOURCE(S): MARPAT 133:74323
GI




L4 ANSWER 131 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)

AB Title compds. 1 [X = halo, CF3, NO2, OH, alkoxy, NH2, alkyl; n = 1-3; Z1, Z2 = CH or N; Y = OCH2 or NHCO; R = OH or alkoxy; R1 = acyl group] or their pharmaceutically acceptable salts were prepared as inhibitors of α4β1 mediated adhesion to either the vascular cell adhesion mol. (VCAM-1) or the CS-1 domain of fibronectin and are useful in the treatment of inflammatory diseases. Approx. 200 invention compds. and their intermediates were prepared by various coupling methods and purified by chromatog. on silica gel. Thus, 4-[(2,6-dichlorobenzoyl)amino]-N-[(1S)-7-hydroxy-1,2,3,4-tetrahydro-3-isoquinolyl]carbonyl]-L-phenylalanine was prepared by deprotection of resin-bound N-(tert-butoxycarbonyl)-4-[(2,6-dichlorobenzoyl)amino]-L-phenylalanine with 50% TFA/CH2Cl2, followed by treatment with (3S)-2-(tert-butoxycarbonyl)-7-hydroxy-1,2,3,4-tetrahydro-3-isoquinolinecarboxylic acid, deprotection, and hydrolysis with 2N LiOH. In vitro cell adhesion inhibitory and/or modulatory activities are reported for > 100 invention compds. tested in Jurkat CS-1 and/or Jurkat endothelial cell (EC) adhesion inhibition assays. Ten compds. showed IC50 values ≤ 0.8 μM in both assays.

IT 279239-07-5P
RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)
(preparation of N-acylphenylalanine derivs. and analogs as inhibitors of α4β1 mediated cell adhesion)

RN 279239-07-5 CAPLUS
CN L-Phenylalanine, 4-[(2,6-dichlorobenzoyl)amino]-N-[(1S)-1,2,3,4-tetrahydro-7-hydroxy-3-isoquinolyl]carbonyl]- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



L4 ANSWER 132 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
ACCESSION NUMBER: 2000:401813 CAPLUS
DOCUMENT NUMBER: 133:43453
TITLE: Preparation of 2-(3-indolyl)quinolines as antibacterial agents
INVENTOR(S): Cuny, Gregory D.; Hauske, James R.; Heefner, Donald L.; Hoemann, Michael Z.; Kumaravel, Gnanasambandam; Melikian-Badalian, Anita; Rossi, Richard P.; Xie, Roger L.
PATENT ASSIGNEE(S): Sepracor, Inc., USA
SOURCE: PCT Int. Appl., 155 pp.
CODEN: PIXXD2
DOCUMENT TYPE: Patent
LANGUAGE: English
FAMILY ACC. NUM. COUNT: 7
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2000034265	A2	20000615	WO 1999-US28744	19991203
WO 2000034265	A3	20021003		
W:	AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM			
RW:	GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LJ, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG			
US 6103905	A	20000815	US 1998-213385	19981211
PRIORITY APPLN. INFO.:			US 1997-878781	B2 19970619
			US 1998-45051	A2 19980319
			US 1998-99640	A2 19980618

OTHER SOURCE(S): MARPAT 133:43453
GI



AB The title compds. (I) [wherein L and Q = independently a hydrophobic group or is absent; X = heterocyclyl, (form)amidinyl, guanidinyl, CN, C(S)NR₂, N(R)C(S)R, OR, SR, NR₂, or PR₂; Z = C.tplbond.C, CH:CH, or CH₂CH₂; R = independently H, (hetero)alkyl, (hetero)aryl, acyl, sulfonyl, etc.; R₁ = H, alkyl, aryl, p-toluenesulfonyl, phthalimidoalkyl, or aminoalkyl; R₂ and R₃ = independently H, alkyl, or acyl] were prepared by standard synthetic and solid phase combinatorial methods. For example, II was synthesized in a 3-step sequence involving: (1) reduction of 2-[5-bromo-1-(tert-butoxycarbonyl)indol-3-yl]-6-(trifluoromethyl)-4-quinolinecarboxylic acid to the alc. with LiAlH₄ (44%), (2) addition of 4-iodo-N-(tert-butoxycarbonyl)benzylamine (preparation given) to the alc. (82%), and (3) indolyl and amine deprotection using TFA (78%). Nearly two-thirds of the 534 indolylquinolines tested in assays against cultures of methicillin-resistant Staphylococcus aureus (MRSA), ciprofloxacin-resistant Staphylococcus aureus (CRSA), vancomycin-resistant Enterococcus spp. (VRE), and/or penicillin-resistant Pseudomonas (PRP) had in vitro min. inhibitory concns. (MICs) ≤ 10 μ M. For 12 of the 15 compds. tested in vivo for toxicity, all mice were surviving 7 days after administration of 40 mg/kg doses.

IT 275357-08-9P

RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)

RN 275357-08-9 CAPLUS

CN 4-Quinolonecarboxamide, N-[2-amino-1-[(4-nitrophenyl)methyl]-2-oxoethyl]-2-(5-chloro-1H-indol-3-yl)- (9CI) (CA INDEX NAME)

ACCESSION NUMBER: 2000:358667 CAPLUS
DOCUMENT NUMBER: 133:177433
TITLE: P-C bond formation: synthesis of phosphino amino acids by palladium-catalyzed cross-coupling
AUTHOR(S): Kratz, Heinz-Bernhard; Pletsch, Andreas
CORPORATE SOURCE: Department of Chemistry, University of Saskatchewan, Saskatoon, SK, S7N 5C9, Can.
SOURCE: Tetrahedron: Asymmetry (2000), 11(7), 1617-1621
CODEN: TASYE3; ISSN: 0957-4166
PUBLISHER: Elsevier Science Ltd.
DOCUMENT TYPE: Journal
LANGUAGE: English
OTHER SOURCE(S): CASREACT 133:177433


AB (4-Diethylphosphinyl)- and (4-diphenylphosphinyl) deriva. of D- and L-phenylalanine were synthesized using a Pd-catalyzed cross-coupling giving the desired products in very high yields and without racemization.

IT 288263-21-8


RL: RCT (Reactant); RACT (Reactant or reagent)
(preparation of phosphino amino acids by palladium-catalyzed cross-coupling)

RN 288263-21-8 CAPLUS

CN Ferrocene, [[[(1S)-1-[(4-iodophenyl)methyl]-2-methoxy-2-oxoethyl]amino]carbonyl]- (9CI) (CA INDEX NAME)



REFERENCE COUNT: 15 THERE ARE 15 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT



ACCESSION NUMBER: 2000:351382 CAPLUS
DOCUMENT NUMBER: 132:330369
TITLE: Treatment of tumors by administration of growth hormone releasing compounds and their antagonists
INVENTOR(S): Muccioli, Giampiero; Papotti, Mauro; Ghigo, Ezio; Deghenghi, Romano
PATENT ASSIGNEE(S): Asta Medica Aktiengesellschaft, Germany
SOURCE: PCT Int. Appl. 32 pp.
CODEN: PIXXD2
DOCUMENT TYPE: Patent
LANGUAGE: English
FAMILY ACC. NUM. COUNT: 2
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2000029011	A1	20000525	WO 1999-EP8662	19991111
W: AU, BG, BR, BY, CA, CN, CZ, DE, GE, HR, HU, ID, IL, IN, IS, JP, KG, KR, KZ, LT, LV, MK, MX, NO, NZ, PL, RO, RU, SG, SI, SK, TR, UA, UZ, YU, ZA, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM				
RW: AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE				
US 6124263	A	20000926	US 1998-192406	19981116
BR 9915390	A	20010807	BR 1999-15390	19991111
EP 1131083	A1	20010912	EP 1999-955974	19991111
EP 1131083	B1	20040121		
R: AT, BR, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO				
JP 2002529512	T2	20020910	JP 2000-582057	19991111
NZ 511280	A	20021025	NZ 1999-511280	19991111
AU 768516	B2	20031218	AU 2000-12706	19991111
ZA 2001003182	A	20010907	ZA 2001-3182	20010419
NO 2001002367	A	20010709	NO 2001-2367	20010514
BG 105572	A	20020131	BG 2001-105572	20010607
PRIORITY APPL. INFO.:			US 1998-192406	A 19981116
			AU 1999-43717	A3 19990528
			WO 1999-EP8662	W 19991111

OTHER SOURCE(S): MARPAT 132:330369

AB A method for treating a tumor in a mammal by administering a growth hormone releasing compound or an antagonist thereof in an amount effective to reduce or inhibit proliferation of tumorigenic cells in the mammal. In particular, the tumors to be treated include lung, mammary, thyroid or pancreas tumors. The preferred compds. are certain peptides that contain Me tryptophan and lysine units.

IT 268545-46-6

RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES (Uses)


(treatment of tumors by administration of growth hormone releasing compds. and antagonists)

RN 268545-46-6 CAPLUS


CN L-Lysinamide, 2-methyl-N-[(2S)-4-methyl-1-oxo-2-[(5S)-6-oxo-1,7-diazaspiro[4.4]non-7-yl]pentyl]-D-tryptophyl-(3R)-2,3,4,4a,5,9b-hexahydro-5-(methylsulfonyl)-1H-pyrido[4,3-b]indole-3-carbonyl-L-phenylalanyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

PAGE 1-A




PAGE 2-A




REFERENCE COUNT: 5 THERE ARE 5 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 135 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
 ACCESSION NUMBER: 2000:288696 CAPLUS
 DOCUMENT NUMBER: 133:12871
 TITLE: Opioid peptide analogs containing 2'-hydroxy,6'-methyltyrosine in place of Tyr1 display greatly enhanced δ antagonist potency but unchanged μ agonist potency
 AUTHOR(S): Berezowska, Irena; Lemieux, Carole; Nguyen, Thi M.
 CORPORATE SOURCE: -D.; Chung, Nga N.; Schiller, Peter W.
 SOURCE: Clinical Research Institute of Montreal, Montreal, QC, H2W 1K7, Can.
 Peptides 1998, Proceedings of the European Peptide Symposium, 25th, Budapest, Aug. 30-Sept. 4, 1998 (1999), Meeting Date 1998, 718-719. Editor(s): Bajusz, Sandor; Hudecz, Ferenc.
 Akademiai Kiado: Budapest, Hung.
 CODEN: 68WKAY
 DOCUMENT TYPE: Conference
 LANGUAGE: English
 AB The authors report the syntheses and in vitro opioid activity profiles of the hmt1-analogs of the δ antagonists TIP (H-Tyr-Tic-Phe-OH) and TIPP (H-Tyr-Tic-Phe-Phe-OH) and of the μ agonists TAPP (H-Tyr-D-Ala-Phe-Phe-NH₂) and DALDA (H-Tyr-D-Arg-Phe-Lys-NH₂). In vitro opioid activities of the compds. were determined in the μ -receptor-representative guinea pig ileum assay and in the δ receptor-representative mouse vas deferens (MVD) assay, and their μ and δ receptor affinities were measured in binding assays based on displacement of [³H]DAMGO and [³H]DSLET, resp., from rat brain membrane binding sites. The tripeptide H-Hmt-Tic-Phe-OH was an about 15 times more potent δ antagonist against the δ agonist DEDPE than its parent TIP, showing δ antagonist potency (MVD) and δ receptor binding affinity in the subnanomolar range. Furthermore, this compound showed greatly improved δ receptor selectivity as compared to TIP. The hmt1-analog of the tetrapeptide TIPP, H-Hmt-Tic-Phe-Phe-OH, displayed very high δ antagonist potency in the MVD assay, comparable to that of H-Dmt-Tic-Phe-Phe-OH. In the binding assays, it showed slightly higher δ receptor affinity than H-Dmt-Tic-Phe-Phe-OH and 20-fold higher δ selectivity. Thus, (Hmt1)TIPP ranks among the most potent and most specific δ opioid antagonists reported to date. Substitution of Hmt for Tyr1 in the μ agonist peptides TAPP and DALDA resulted in μ -agonist potencies comparable to those of their resp. parent peptides. In conclusion, replacement of Tyr1 in opioid peptides with Hmt produced a potency increase in the case of the δ antagonists but not in the case of the μ agonists.
 IT RL: BAC (Biological activity or effector, except adverse); BPR (Biological process); BSU (Biological study, unclassified); PRP (Properties); BIOL (Biological study); PROC (Process)
 (opioid peptide analogs δ antagonist and μ agonist activity in relation to structure)
 RN 271795-36-9 CAPLUS
 CN L-Phenylalanine, 2-hydroxy-6-methyl-L-tyrosyl-(3S)-1,2,3,4-tetrahydro-3-isoquinolinecarboxyl- (9CI) (CA INDEX NAME)
 Absolute stereochemistry.



REFERENCE COUNT: 6 THERE ARE 6 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT


L4 ANSWER 136 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
 ACCESSION NUMBER: 2000:269112 CAPLUS
 DOCUMENT NUMBER: 133:37726
 TITLE: N-acyl phenylalanine analogues as potent small molecule VLA-4 antagonists
 AUTHOR(S): Chen, Li; Tilley, Jefferson W.; Huang, Tai-Nan; Miklowick, Dorota; Trilles, Richard; Guthrie, Robert W.; Luk, Kin; Hanglow, Angela; Rowan, Karen; Schwinge, Virginia; Wolitzky, Barry
 CORPORATE SOURCE: Roche Research Center, Hoffmann-La Roche, Inc., Nutley, NJ, 07110, USA
 SOURCE: Bioorganic & Medicinal Chemistry Letters (2000), 10(8), 725-727
 CODEN: BMCLE8; ISSN: 0960-894X
 PUBLISHER: Elsevier Science Ltd.
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 AB We have identified a series of low mol. weight (Mr <500) N-acylphenylalanines that are effective inhibitors of the VCAM-VLA-4 interaction. Investigation of the SAR of the N-acyl moiety led to the identification of N-benzylpyroglutamyl deriva. as being particularly potent.
 IT 275802-12-59
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation)
 (N-acyl phenylalanine analogues as VLA-4 antagonists)
 RN 275802-12-5 CAPLUS
 CN L-Tyrosine, N-(9-acridinylcarbonyl)-O-pentyl- (9CI) (CA INDEX NAME)
 Absolute stereochemistry.



REFERENCE COUNT: 7 THERE ARE 7 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 137 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
 ACCESSION NUMBER: 2000:236481 CAPLUS
 DOCUMENT NUMBER: 133:28301
 TITLE: Influence of Halogen Salts on the Production of the Ochratoxins by *Aspergillus ochraceus* Wilh.
 AUTHOR(S): Stander, Maria A.; Steyn, Pieter S.; Luebben, Annelie; Miljkovic, Ana; Mantle, Peter G.; Marais, Gert J.
 CORPORATE SOURCE: School of Chemistry and Biochemistry, University of Potchefstroom, Potchefstroom, 2520, S. Afr.
 SOURCE: Journal of Agricultural and Food Chemistry (2000), 48(5), 1865-1871
 CODEN: JAFCAU; ISSN: 0021-8561
 PUBLISHER: American Chemical Society
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 AB The first report of the biol. production of bromo ochratoxin B by *Aspergillus ochraceus* Wilh. is presented as well as a study of the influence of potassium bromide, potassium iodide, potassium fluoride, and potassium chloride on the production of ochratoxin A and ochratoxin B. Potassium fluoride and potassium iodide inhibited the growth of the fungus, whereas potassium chloride substantially stimulated the production of ochratoxin A in shaken solid substrate fermentation on whole wheat or shredded wheat, generally giving a high yield of ochratoxins. Increasing levels of potassium bromide led to a decline in ochratoxin A production and an increase in bromo-ochratoxin B, ochratoxin B, and 4-hydroxy ochratoxin B. Nevertheless, *A. ochraceus* was much less versatile in the bromo analogs than other fungi, which produce metabolites containing chlorine. Anal. included aminopropyl solid-phase extraction column cleanup followed by quant. anal. on reversed-phase HPLC using fluorescence detection and employing N-(5-chloro-2-hydroxybenzoyl)phenylalanine as an internal standard
 IT 255042-26-3P
 RL: PRP (Properties); PUR (Purification or recovery); PREP (Preparation) (Influence of potassium halides on production of ochratoxins by *Aspergillus ochraceus*)
 RN 255042-26-3 CAPLUS
 CN L-Phenylalanine, N-[[[(3R)-5-bromo-3,4-dihydro-8-hydroxy-3-methyl-1-oxo-1H-2-benzopyran-7-yl]carbonyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



REFERENCE COUNT: 25 THERE ARE 25 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 138 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
 ACCESSION NUMBER: 2000:144899 CAPLUS
 DOCUMENT NUMBER: 132:189658
 TITLE: Amino acid derivative and peptide anti-cancer compounds and methods
 INVENTOR(S): Stewart, John M.; Chan, Daniel C. F.; Gera, Lojos; York, Eunice; Bunn, Paul
 PATENT ASSIGNEE(S): USA
 SOURCE: PCT Int. Appl., 55 pp.
 CODEN: PIXXD2
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2000011022	A1	20000302	WO 1999-US19381	19990820
W:	AR, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, GR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM			
RW:	GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG			
US 6388054	B1	20020514	US 1999-378019	19990819
AU 2000015959	A1	20000314	AU 2000-15959	19990820
US 2002183252	A1	20021205	US 2001-35662	20011228
PRIORITY APPLN. INFO.:			US 1998-97210P	P 19980820
			US 1999-141169P	P 19990625
			US 1999-378019	A 19990819
			WO 1999-US19381	W 19990820

OTHER SOURCE(S): MARPAT 132:189658
 AB The invention provides amino acid derivative and peptidic compds. useful to inhibit tumor growth and to induce apoptosis. In general, the anti-cancer agents (ACA) are described by the formula [ACA]-X [X = linker group with 2-5 functional groups or is absent; n = 1; ACA as described in the invention (Markush included)].

IT 259883-84-6P
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); PRP (Properties); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)
 (peptide and non-peptide anti-cancer compds. and methods)
 RN 259883-84-6 CAPLUS
 CN L-Arginine, N2-[(2S)-1-oxo-4-phenyl-2-[[[(3R)-1,2,3,4-tetrahydro-2-(tricyclo[3.3.1.1.3,7]dec-1-ylacetyl)-3-isquinolinyl]carbonyl]amino]butyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

L4 ANSWER 138 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)



REFERENCE COUNT: 4 THERE ARE 4 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 139 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
 ACCESSION NUMBER: 2000:84777 CAPLUS
 DOCUMENT NUMBER: 132:137288
 TITLE: Preparation of isoquinolinylguanidines as urokinase inhibitors
 INVENTOR(S): Barber, Christopher Gordon; Dickinson, Roger Peter; Fish, Paul Vincent
 PATENT ASSIGNEE(S): Pfizer Inc., USA; Pfizer Limited
 SOURCE: PCT Int. Appl., 222 pp.
 CODEN: PIXXD2
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2000005214	A2	20000203	WO 1999-IB1289	19990715
WO 2000005214	A3	20010531		
W:	AR, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, GR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM			
RW:	GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG			
CA 2337247	AA	20000203	CA 1999-2337247	19990715
AU 9945289	A1	20000214	AU 1999-45289	19990715
AU 745545	B2	20020321		
EP 1077945	A1	20010228	EP 1999-928177	19990715
EP 1077945	B1	20030108		
R:	AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO			
BR 9912374	A	20010417	BR 1999-12374	19990715
EE 200100049	A	20020617	EE 2001-49	19990715
JP 2002521367	T2	20020716	JP 2000-561170	19990715
AT 230730	T	20030115	AT 1999-928177	19990715
PT 1077945	E	20030331	PT 1999-99928177	19990715
ES 2189434	T3	20030701	ES 1999-928177	19990715
EW 530049	B	20030501	TW 1999-88112548	19990723
ZA 2001000230	A	20020301	ZA 2001-230	20010109
NO 2001000400	A	20010326	NO 2001-400	20010123
HR 2001000059	A1	20020228	HR 2001-59	20010123
BG 105252	A	20011231	BG 2001-105252	20010214
PRIORITY APPLN. INFO.:			GB 1998-16228	A 19980724
			GB 1999-88229	A 19990416
			WO 1999-IB1289	W 19990715

OTHER SOURCE(S): MARPAT 132:137288
 AB R31R2Z2Z2R [I; R = N:C(NH2)2 or NHC:(NH)NH2; R2 = H, alkyl, (hetero)aryl, etc.; R3 = CO2R7, CH2OH, CONRR8R9, CH2NRR8R9; R7 = H or alkyl; R8, R9 = H, (hydroxy)alkyl, etc.; NR8R9 = heterocyclyl; Z = (4-chloro)isoquinoline-7,1-diyl; Z1 = (cyclo)alkylene, heterocyclyl; Z2 = CO, etc.; Z1NR2 = azetidine-, pyrrolidine-, or (homopiperidinediyl); Z2 = CO, CH2, SO2) were prepared. Thus, 2-(H2N)C6H4CO2CMe3 was condensed with 7-bromo-1,4-dichloroisoquinoline (preparation each given) and SOCl2 and the product iminated by (H2N)2C:NH to give, after saponification, 2-(HO2C)C6H4NHSO2ZN:C(NH2)2.HCl (R = 4-chloroisoquinoline-7,1-diyl). Data for biol. activity of I were given.

IT 256476-58-1P

L4 ANSWER 139 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)
 (prepn. of isoquinolinylguanidines as urokinase inhibitors)
 RN 256476-58-1 CAPLUS
 CN Benzeneacetic acid, α -[[[1-[(aminoiminomethyl)amino]-4-chloro-7-isoquinolinyl]carbonyl]amino]-, mono(trifluoroacetate) (9CI) (CA INDEX NAME)

CM 1
 CRN 256476-57-0
 CMF C19 H16 Cl N5 O3




CM 2
 CRN 76-05-1
 CMF C2 H F3 O2



L4 ANSWER 140 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
 ACCESSION NUMBER: 2000:814 CAPLUS
 DOCUMENT NUMBER: 132:207466
 TITLE: Phenylglycine Methyl Ester, a Useful Tool for Absolute Configuration Determination of Various Chiral Carboxylic Acids
 AUTHOR(S): Yabuuchi, Tetsuya; Kusumi, Takenori
 CORPORATE SOURCE: Faculty of Pharmaceutical Sciences, Tokushima University, Tokushima, 770-8505, Japan
 SOURCE: Journal of Organic Chemistry (2000), 65(2), 397-404
 CODEN: JOCEAH, ISSN: 0022-3263
 PUBLISHER: American Chemical Society
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 AB A new chiral anisotropic reagent, phenylglycine Me ester (PGME), developed for the elucidation of the absolute configuration of chiral α,α -disubstituted acetic acids, has turned out to be applicable to other substituted carboxylic acids, such as chiral α -hydroxy-, α -alkoxy-, and α -acyloxy α,α -disubstituted acetic acids, as well as to chiral β,β -disubstituted propionic acids. Because a carboxylic moiety is convertible from other functional groups, e.g., ozonolysis of an olefin and oxidative cleavage of a glycol, the present findings can expand the utility of the PGME method to the absolute configuration determination of various types of organic compds., even those which initially lack oxygen functions. Several examples of the combination of chemical reactions and the PGME method are described.
 IT 259877-12-8P
 RL: PRP (Properties); SPN (Synthetic preparation); PREP (Preparation)
 (determination of absolute configuration of chiral carboxylic acids using phenylglycine Me ester as anisotropic reagent)
 RN 259877-12-8 CAPLUS
 CN Benzeneacetic acid, α -[[[2S]-3,4-dihydro-6-hydroxy-2,5,7,8-tetramethyl-2H-1-benzopyran-2-yl]carbonyl]amino]-, methyl ester, (aS)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.





REFERENCE COUNT: 16 THERE ARE 16 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 141 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
 ACCESSION NUMBER: 1999:795654 CAPLUS
 DOCUMENT NUMBER: 132:22957
 TITLE: Preparation of spiroperidine derivatives as melanocortin receptor agonists
 INVENTOR(S): Nargund, Ravi P.; Ye, Zhixiong; Palucki, Brenda L.; Bakshi, Raman K.; Patchett, Arthur A.; Van Der Ploeg, Leonardus H. T.
 PATENT ASSIGNEE(S): Merck & Co., Inc., USA
 SOURCE: PCT Int. Appl., 77 pp.
 CODEN: PIXXD2
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 9964002	A1	19991216	WO 1999-US13252	19990610
W: AE, AL, AM, AU, AZ, BA, BB, BG, BR, BY, CA, CN, CU, CZ, EE, GD, GE, HR, HU, ID, IL, IN, IS, JP, KG, KR, KZ, LC, LR, LT, LV, MD, MG, MK, MW, MX, NO, NZ, PL, RO, RU, SG, SI, SK, TJ, TM, TR, TT, UA, US, UZ, VN, YU, ZA, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, RW: GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG				
CA 2334551	AA	19991216	CA 1999-2334551	19990610
AU 9946801	A1	19991230	AU 1999-46801	19990610
AU 742425	B2	20020103		
EP 1085869	A1	20010328	EP 1999-930220	19990610
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, PT, IE, SI, LT, LV, FI, RO				
US 6294534	B1	20010925	US 1999-329814	19990610
JP 2002517444	T2	20020618	JP 2000-553071	19990610
US 2001029259	A1	20011011	US 2001-781373	20010212
US 6410548	B2	20020625		
PRIORITY APPL. INFO.:				
			US 1998-88908P	P 19980611
			GB 1998-17179	A 19980806
			US 1999-123260P	P 19990308
			US 1999-329814	A3 19990610
			WO 1999-US13252	W 19990610
OTHER SOURCE(S):			MARPAT 132:22957	
GI				

L4 ANSWER 141 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)



AB Certain novel spiroperidine compds. I [Cy2 = six-membered aromatic ring containing 0 or 1 N; X = O, CH2, etc.; Q = Q1; Y = CO, SO2, etc.; R1, Rb = H, C1-8 alkyl, etc.; R2 = H or halo, R3 = Rb, halo, ORb, N(Rb)2, SO2Rb, CF3, OCF3; Cy = aryl, 5 or 6 membered heteroaryl, 5 or 6 membered heterocyclyl, 5 or 6 membered carbocyclyl; m, p, q independently = 0, 1, or 2] are agonists of melanocortin receptors (no data) and are useful for the treatment, control or prevention of diseases and disorders responsive to the activation of melanocortin receptors. The compds. of the present invention are therefore useful for treatment of diseases and disorders such as obesity, diabetes, sexual dysfunction including erectile dysfunction and female sexual dysfunction.
 IT 252008-71-2P
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)
 (preparation of spiroperidine derivs. as melanocortin receptor agonists)
 RN 252008-71-2 CAPLUS
 CN 2-[[[1H]-isoquinolin-1-yl]carboxylic acid, 3-[[[1R]-1-carboxy-2-(4-chlorophenyl)ethyl]amino]carbonyl]-3,4-dihydro-, 2-(1,1-dimethylethyl) ester, (3R)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




L4 ANSWER 141 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)

REFERENCE COUNT: 3 THERE ARE 3 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 142 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 1999:753019 CAPLUS
 DOCUMENT NUMBER: 132:12506
 TITLE: Preparation of peptides for treating diseases of excessive bone loss or cartilage or matrix degradation as cysteine protease inhibitors
 INVENTOR(S): Bondinell, William Edward; Desjarlais, Renee Louise; Veber, Daniel Frank; Yamashita, Dennis Shinji
 PATENT ASSIGNEE(S): Smithkline Beecham Corporation, USA
 SOURCE: PCT Int. Appl., 128 pp.
 CODEN: PIAXD2
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 9959526	A2	19991125	WO 1999-US11266	19990520
WO 9959526	A3	20000120		
W: AE, AL, AU, BA, BB, BG, BR, CA, CN, CZ, EE, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KP, KR, LC, LK, LR, LT, LV, MG, MK, MN, MX, NO, NZ, PL, RO, SG, SI, SK, SL, TR, TT, UA, US, UZ, VN, YU, ZA, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM				
RW: GH, GM, KE, LS, MM, SD, SL, SZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG				
CA 2332531	AA	19991125	CA 1999-2332531	19990520
EP 1067894	A2	20010117	EP 1999-924421	19990520
R: BE, CH, DE, ES, FR, GB, IT, LI, NL				
JP 2002515411	T2	20020528	JP 2000-549192	19990520
US 6518267	B1	20030211	US 2000-700828	20001121
PRIORITY APPLN. INFO.:			US 1998-86557P	P 19980521
			WO 1999-US11266	W 19990520
OTHER SOURCE(S):			MARPAT 132:12506	
GI				



AB The present invention provides peptides bis-aminomethyl carbonyl protease


L4 ANSWER 142 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)

inhibitors I (R1, R2 = alkyl, P = X, Y; R3 selected from the group consisting of: CH2CH(CH3)2, CH2CH2CH3, CH2CH=CH2, or CH2Ph; R4 is selected from the group consisting of: alkyl, N-piperazine, N-tetrahydroisoquinoline, substituted alkyl, Ph, benzofuran, benzothiazole, quinoline, naphthyl, and benzoxazole; R5 = Ph and Ph substituted with alkyl, N-piperidine, benzofuran, pyridine; Q = arylacyl) and pharmaceutically acceptable salts, hydrates and solvates thereof which inhibit proteases, including cathepsin K, pharmaceutical compns. of such compds., and methods for treating diseases of excessive bone loss or cartilage or matrix degradn., including osteoporosis; gingival disease including gingivitis and periodontitis; arthritis, more specifically, osteoarthritis and rheumatoid arthritis; Paget's disease; hypercalcemia of malignancy; and metabolic bone disease, comprising inhibiting said bone loss or excessive cartilage or matrix degradn. by administering to a patient in need thereof a compd. of the present invention. Thus, (S)-3N-(N-(thianaphthenyl-2-carbonyl)-leucinyll)-amino-1N-(3-[2-(1-oxo-pyridyl)phenylacetyl]-amino-butan-2-one was prepd. for treating diseases of excessive bone loss or cartilage or matrix degradn. as cysteine protease inhibitor. Detn. of cathepsin K proteolytic catalytic activity of these compds. are reported.

IT 251458-61-4P
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)
 (preparation of peptides for treating diseases of excessive bone loss or cartilage or matrix degradation as cysteine protease inhibitors)

RN 251458-61-4 CAPLUS
 CN 8-Quinolincarboxamide, N-[(1S)-2-oxo-2-[[2-oxo-3-[[[3-(2-pyridinyl)phenyl]acetyl]amino]butyl]amino]-1-(phenylmethyl)ethyl]- (9CI)
 (CA INDEX NAME)

Absolute stereochemistry.




L4 ANSWER 143 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 1999:750286 CAPLUS
 DOCUMENT NUMBER: 132:137250
 TITLE: Preparation and antibacterial activity of sulfonamide derivatives and amides of coumarin compounds
 AUTHOR(S): Shah, Sonal; Desai, Devki; Mehta, R. H.
 CORPORATE SOURCE: Department of Chemistry, Faculty of Science, M. S. University of Baroda, Vadodara, 390 002, India
 SOURCE: Journal of the Indian Chemical Society (1999), 76(10), 507-508
 CODEN: JICSAH; ISSN: 0019-4522
 PUBLISHER: Indian Chemical Society
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 GI

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

AB Coumarin sulfonamide deriva. I and II (R = Me, MeO, AcNH, Cl, Br; R1 = H, Br) were prepared by sulfonylation of an aminophenylaminocarbonyl coumarin, and by nucleophilic displacement of a chloromethyl coumarin with o-phenylenediamine followed by sulfonylation. Coumarin deriva. III (R2 = Me, Me2CH, MeSCH2CH2, HOCH2; X = bond, CH2) and their racemic deriva (R2 = H, Me, Me2CH, MeSCH2CH2, PhCH2; X = bond, CH2) were also prepared from a coumarin acid chloride and racemic and L-amino acid deriva. I, II, and III were tested for their antibiotic activity against E. coli, S. aureus, S. typhosa, and S. albus; I and II showed moderate to low activity against the tested bacteria, while the coumarins III showed no activity against any of the bacteria.

IT 256658-76-1P
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation)
 (preparation and antibacterial activity of coumarin sulfonamide and amino acid deriva.)
 RN 256658-76-1 CAPLUS
 CN Phenylalanine, N-[(8-methoxy-2-oxo-2H-1-benzopyran-3-yl)carbonyl]- (9CI)
 (CA INDEX NAME)



REFERENCE COUNT: 16 THERE ARE 16 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT


L4 ANSWER 144 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
 ACCESSION NUMBER: 1999:735984 CAPLUS
 DOCUMENT NUMBER: 132:107853
 TITLE: Cycloaddition of homophthalic anhydrides with aldehydes and ketones: a route to 3,4-dihydroisocoumarin-4-carboxylic acid derivatives
 AUTHOR(S): Yu, Neifang; Poulain, Rebecca; Tartar, Andre; Gesquiere, Jean-Claude
 CORPORATE SOURCE: Faculte de Pharmacie, Institut Pasteur and UMR CNRS, Lille, 59006, Fr.
 SOURCE: Tetrahedron (1999), 55(48), 13735-13740
 CODEN: TETRAB; ISSN: 0040-4020
 PUBLISHER: Elsevier Science Ltd.
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 OTHER SOURCE(S): CASREACT 132:107853

AB Homophthalic anhydride (1H-2-benzopyran-1,3(4H)-dione) reacts with benzaldehyde, in the presence of boron trifluoride-di-Et ether complex to give the cycloadduct 3-phenyl-3,4-dihydroisocoumarin-4-carboxylic acid in good to excellent yield. Under these conditions, we did not observe the formation of Perkin-type products. The reaction can be extended to a wide variety of aldehydes and to some ketones in synthetically useful yields. Amides can be obtained in good yields after activation of the carboxylic function with DCC at 0°C.

IT 255841-53-3P
 RL: SPN (Synthetic preparation); PREP (Preparation)
 (preparation of dihydroisocoumarin-4-carboxylic acid via cycloaddn. of homophthalic anhydrides with aldehydes or ketones)


RN 255841-53-3 CAPLUS
 CN Benzeneacetic acid, α -[[(3R,4R)-3,4-dihydro-1-oxo-3-phenyl-1H-2-benzopyran-4-yl]carbonyl]amino]-, methyl ester, rel- (9CI) (CA INDEX NAME)

Relative stereochemistry.



REFERENCE COUNT: 15 THERE ARE 15 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 145 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
 ACCESSION NUMBER: 1999:735231 CAPLUS
 DOCUMENT NUMBER: 132:93616
 TITLE: The synthesis of bromo- and iodo-ochratoxin B
 AUTHOR(S): Steyn, Pieter S.; Payne, Barry E.
 CORPORATE SOURCE: School of Chemistry and Biochemistry, Potchefstroom University for CHE, Potchefstroom, 2520, S. Afr.
 SOURCE: South African Journal of Chemistry (1999), 52(2/3), 69-70
 CODEN: SAJCDG; ISSN: 0379-4350
 PUBLISHER: South African Chemical Institute
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 OTHER SOURCE(S): CASREACT 132:93616
 GI




AB The effective synthesis of 5-bromo- and 5-iodo-ochratoxin B I (R = H, R1 = Br, iodo) by halogenation of ochratoxin B I (R = R1 = H) with pyridinium hydrobromide perbromide, and iodine and mercury(II) oxide, resp., was reported. 5-Iodo-ochratoxin B was further converted to 5-iodo-ochratoxin B Me ester I (R = Me, R1 = iodo) using thionyl chloride and methanol.

IT 255042-27-4P, 5-Iodo-ochratoxin B
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)
 (synthesis of bromo- and iodo-ochratoxin B)

RN 255042-27-4 CAPLUS
 CN L-Phenylalanine, N-[[[(3R)-3,4-dihydro-8-hydroxy-5-iodo-3-methyl-1-oxo-1H-2-benzopyran-7-yl]carbonyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




REFERENCE COUNT: 22 THERE ARE 22 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 145 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)

L4 ANSWER 146 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
 ACCESSION NUMBER: 1999:708602 CAPLUS
 DOCUMENT NUMBER: 131:310837
 TITLE: Preparation of phenylalanine sulfonamide derivatives and related compounds as CCR-3 receptor antagonists
 INVENTOR(S): Dhanak, Dashyant; Widdowson, Katherine L.; White, John R.
 PATENT ASSIGNEE(S): Smithkline Beecham Corporation, USA
 SOURCE: PCT Int. Appl., 23 pp.
 CODEN: PIXXD2
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 9955330	A1	19991104	WO 1999-US8950	19990427
W: CA, JP, US RW: AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE				
CA 2329821	AA	19991104	CA 1999-2329821	19990427
EP 1073434	A1	20010207	EP 1999-921462	19990427
R: BE, CH, DE, ES, FR, GB, IT, LI, NL				
JP 2002512960	T2	20020508	JP 2000-545529	19990427
PRIORITY APPLN. INFO.: US 1998-83229P P 19980427 WO 1999-US8950 W 19990427				
OTHER SOURCE(S): MARPAT 131:310837 GI				




AB The title compds. I [R1 = alkyl, aryl, heteroaryl, etc.; R2 = alkyl, benzyl; R3 = OH, alkoxy, NO2, NH2, etc.; m = 1-3; n = 0-3] and EtO2CCH(R)NHCOPh (R = indolylmethyl, Ph, CH2CH2Ph), CCR-3 receptor antagonists (no data), were prepared E.g., (S)-Et 2-benzoylamino-3-(4-nitrophenyl)propionate was prepared

IT 247580-60-5P
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USBS (Uses)
 (preparation of phenylalanine sulfonamide derivs. and related compds. as CCR-3 receptor antagonists)

RN 247580-60-5 CAPLUS
 CN L-Tyrosine, N-(8-quinolinylcarbonyl)-, ethyl ester (9CI) (CA INDEX NAME)

Absolute stereochemistry.

L4 ANSWER 146 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)




REFERENCE COUNT: 3 THERE ARE 3 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 147 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 1999:655304 CAPLUS
 DOCUMENT NUMBER: 132:64499
 TITLE: Angiotensin II analogs encompassing 5,9- and 5,10-fused thiazabicycloalkane tripeptide mimetics
 AUTHOR(S): Johansson, Petra; Lindeberg, Gunnar; Tong, Weimin; Gogoll, Adolf; Synnergren, Barbro; Nyberg, Fred; Karlén, Anders; Hallberg, Anders
 CORPORATE SOURCE: Department of Organic Pharmaceutical Chemistry, Uppsala University, Uppsala, SE-751 23, Swed.
 SOURCE: Journal of Medicinal Chemistry (1999), 42(22), 4524-4537
 CODEN: JMCMAR; ISSN: 0022-2623
 PUBLISHER: American Chemical Society
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 AB A simple exptl. procedure on solid phase for the construction of new tripeptidic 5,9- and 5,10-fused thiazabicycloalkane scaffolds that adopt β -turns has been developed. This N terminal-directed bicyclization, relying on masked aldehyde precursors derived from glutamic acid as key building blocks, provides a complement to the related bicyclization previously reported, where an aspartic acid-derived precursor was employed to induce cyclization toward the C-terminal end of the peptide. Thus, the regioselectivity of the bicyclization can be altered simply by varying the chain length of the incorporated aldehyde precursor. Four analogs of the hypertensive octapeptide angiotensin II, comprising the new scaffolds in the 3-5- and 5-7-positions, were synthesized. One of these conformationally constrained angiotensin II analogs exhibited AT₁ receptor affinity (K_i = 750 nM). Results from theor. conformational anal. of model compds. of the bicyclic tripeptide mimetics are presented, and they demonstrate that subtle differences in geometry have a strong impact on the affinity to the AT₁ receptor.
 IT 253277-45-1P
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation)
 (preparation and biol. activity of angiotensin II analogs containing thiazabicycloalkane tripeptide mimetics)
 RN 253277-45-1 CAPLUS
 CN L-Phenylalanine, L- α -aspartyl-L-arginyl-L-valyl-L-tyrosyl-L-cysteiny-L-histidyl-(5R)-5-mercapto-L-prolyl-, cyclic (5-7)-thioether (9CI) (CA INDEX NAME)

Absolute stereochemistry.

L4 ANSWER 147 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)




REFERENCE COUNT: 66 THERE ARE 66 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 148 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 1999:629552 CAPLUS
 DOCUMENT NUMBER: 132:9841
 TITLE: Oxidation of Ochratoxin A by an Fe-Porphyrin System: Model for Enzymatic Activation and DNA Cleavage
 AUTHOR(S): Gillman, Ivan G.; Clark, T. Nicole; Manderville, Richard A.
 CORPORATE SOURCE: Department of Chemistry, Wake Forest University, Winston-Salem, NC, 27109-7486, USA
 SOURCE: Chemical Research in Toxicology (1999), 12(11), 1066-1076
 CODEN: CRTOEC; ISSN: 0893-228X
 PUBLISHER: American Chemical Society
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 AB Ochratoxin A (OTA) is a fungal toxin that facilitates single-strand DNA cleavage, DNA adduction, and lipid peroxidn. when metabolically activated. To model the enzymic activation of OTA, we have employed the water-soluble iron(III) meso-tetrakis(4-sulfonatophenyl)porphyrin (FeTPPS) oxidation system. In its presence, OTA has been found to facilitate single-strand cleavage of supercoiled plasmid DNA through production of reactive oxygen species (ROS) (i.e., the hydroxyl radical, HO \cdot). The reaction of OTA with the FeTPPS oxidation system also generated three hydroxylated products (chlorine atom still attached), which was taken as evidence for production of the known hydroxylated metabolites of OTA. This result suggested that the FeTPPS system served as a reasonable model for the enzymic activation of OTA. When the reaction of OTA with FeTPPS was carried out in the presence of excess hydrogen peroxide (H₂O₂) and sodium ascorbate, a hydroquinone species (OTHQ) was detected in which an OH group has replaced the chlorine atom of OTA. The production of OTHQ was dependent on the presence of the reducing agent, sodium ascorbate, which suggested that the oxidation catalyst furnished the quinone derivative OTQ that was subsequently reduced to OTHQ by ascorbate. Utilizing a synthetic sample of OTHQ, the hydroquinone was found to undergo autoxidn. with a $t_{1/2}$ of 11.1 h at pH 7.4, and to possess a pK_a value of 8.03 for the phenolic oxygen ortho to the carbonyl groups. Our findings imply that the hydroquinone (OTHQ) and quinone (OTQ) metabolites of OTA have the ability to cause alkylation/redox damage and have allowed us to propose a viable pathway for oxidative damage by OTA.
 IT 205034-32-8P
 RL: BPR (Biological process); BSU (Biological study, unclassified); MFM (Metabolic formation); SPN (Synthetic preparation); BIOL (Biological study); FORM (Formation, nonpreparative); PREP (Preparation); PROC (Process)
 (preparation of ochratoxin hydroquinone and oxidation of ochratoxin A by Fe-porphyrin system)
 RN 205034-32-8 CAPLUS
 CN L-Phenylalanine, N-[[[(3R)-3,4-dihydro-5,8-dihydroxy-3-methyl-1-oxo-1H-2-benzopyran-7-yl]carbonyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




L4 ANSWER 148 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)

REFERENCE COUNT: 47 THERE ARE 47 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 149 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 1999:613947 CAPLUS
 DOCUMENT NUMBER: 131:243287
 TITLE: Preparation of dioxopiperazinoacetamides as fructose-1,6-bisphosphatase inhibitors
 INVENTOR(S): Mjalli, Adnan M. M.; Mason, James Christopher; Arienti, Kristen Lee; Short, Kevin Michael; Kimmich, Rachel Denise Anne; Jones, Todd Kevin
 PATENT ASSIGNEE(S): Ontogen Corporation, USA
 SOURCE: PCT Int. Appl., 74 pp.
 CODEN: PIXXD2
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 9947549	A1	19990923	WO 1999-US5552	19990315
W: AU, CA, JP				
RW: AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE				
CA 2289621	AA	19990923	CA 1999-2289621	19990315
AU 9930870	A1	19991011	AU 1999-30870	19990315
US 6107274	A	20000822	US 1999-270121	19990315
EP 1070084	A1	20010124	EP 1999-912505	19990315
R: DE, FR, GB				
JP 2001294586	A2	20011023	JP 2000-386045	19990315
PRIORITY APPLN. INFO:			US 1998-78065P	P 19980316
			WO 1999-US5552	W 19990315
OTHER SOURCE(S):		MARFAT 131:243287		
GI				




AB Title compds. [I; R1 = cycloalkyl or aralkyl; R2 = cycloalkylmethyl or (ar)alkyl; R3 = H, F, alkyl, substituted Ph; R4 = H, alkyl, acyl, substituted Ph; R5 = H; R1R5 = atoms to complete a ring] were prepared. Thus, L-R2CH(NH2)CO2Me.HCl (R2 = cyclohexyl), 4-(NC)C6H4CHO, N-Fmoc-1,2,3,4-tetrahydroisoquinoline-3-carboxylic acid, and 4-(CNH2CH2C)C6H4OCH2Ph were subjected to Ugi condensation and the product cyclized to give, after deprotection, I (R1R5 = 2-(H2C)C6H4CH2, R2 = cyclohexylmethyl, R3 = 4-(NC)C6H4, R4 = CH2CH2C6H4(OH)-4). Data for biol.

L4 ANSWER 149 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)

IT 244220-68-6P
 RL: BYP (byproduct); PREP (Preparation)
 (preparation of dioxopiperazinoacetamides as fructose-1,6-bisphosphatase inhibitors)
 RN 244220-68-6 CAPLUS
 CN 2(1H)-Isoquinolinecarboxylic acid, 3-[[[(1R)-1-(4-cyanophenyl)-2-oxo-2-[[2-(4-(phenylmethoxy)phenyl]ethyl]amino]ethyl]][(1S)-1-(cyclohexylmethyl)-2-methoxy-2-oxoethyl]amino]carbonyl]-3,4-dihydro-7-hydroxy-, 9H-fluoren-9-ylmethyl ester, (3S)-(9CI) (CA INDEX NAME)

Absolute stereochemistry.



REFERENCE COUNT: 1 THERE ARE 1 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT


L4 ANSWER 150 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 1999:583907 CAPLUS
 DOCUMENT NUMBER: 131:294930
 TITLE: Characterization of spatially addressable libraries: stereoisomer analysis of tetrahydro-β-carbolines as an example
 AUTHOR(S): Cheng, Cesar C.; Chu, Yen-Ho
 CORPORATE SOURCE: Department of Chemistry, The Ohio State University, Columbus, OH, 43210, USA
 SOURCE: Journal of Combinatorial Chemistry (1999), 1(6), 461-466
 CODEN: JCCHFF; ISSN: 1520-4766
 PUBLISHER: American Chemical Society
 DOCUMENT TYPE: Journal
 LANGUAGE: English

AB Combinatorial chemical approaches have facilitated the process of lead discovery and optimization for new drugs, catalysts, and materials. A successful combinatorial program typically includes high throughput library synthesis, characterization, and screening. One of many challenges in this program is to develop high throughput characterization methods to address issues concerning reaction stereoselectivity and racemization during the library synthesis. Using the core structure of tetrahydro-β-carboline as an example, the authors demonstrate the usefulness of capillary electrophoresis in enantiomeric separation of stereoisomers generated by parallel synthesis and rapid quant. measurement of the isomer ratios, in addition to assessing possible racemization during reactions for its overall potential in library characterization.


IT 246137-76-8P
 RL: ANT (Analyte); PEP (Physical, engineering or chemical process); PRP (Properties); SPN (Synthetic preparation); ANST (Analytical study); PREP (Preparation); PROC (Process)
 (characterization of spatially addressable libraries: stereoisomer anal. of tetrahydro-β-carbolines)
 RN 246137-76-8 CAPLUS
 CN L-Tyrosine, N-[[[(1S,3S)-1-ethyl-2,3,4,9-tetrahydro-1-methyl-1H-pyrido[3,4-b]indol-3-yl]carbonyl]]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



REFERENCE COUNT: 37 THERE ARE 37 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 151 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
 ACCESSION NUMBER: 1999-567712 CAPLUS
 DOCUMENT NUMBER: 131:271976
 TITLE: Synthesis and electrochemical anion recognition by novel redox-responsive ferrocenyl dipeptide ester derivatives; 1H NMR anion complexation studies
 AUTHOR(S): Gallagher, John F.; Kenny, Peter T. M.; Sheehy, Michael J.
 CORPORATE SOURCE: Dublin City University, Dublin, 9, Ire.
 SOURCE: Inorganic Chemistry Communications (1999), 2(8), 327-330
 CODEN: ICCOFP; ISSN: 1387-7003
 PUBLISHER: Elsevier Science S.A.
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 AB The synthesis of six N-ferrocenyl dipeptide ester deriva. FeCOR [Fc = (η⁵-C₅H₅)Fe(η⁵-C₅H₄); R = Gly-Gly(OMe) (1), Leu-Leu(OEt) (2), Ala-Phe(OEt) (3), Phe-Phe(OMe) (4), Phe-Leu(OBn) (5) and Phe-Ser(OEt) (6)] is described. Derivatives 1-3 were prepared by coupling reactions of FcCOOH with the free N-terminal dipeptide esters. The electrochem. anion sensing behavior (using a Pt microdisk working electrode) and 1H NMR anion coordination studies of these novel pendant N-ferrocenyl dipeptide ester deriva. with a range of anions are reported. Best results were observed with 1, which sensed halides and dihydrogen phosphate guest anions with marked selectivity, the trend being H₂PO₄⁻ > Cl⁻ > Br⁻ > HS₄⁻.
 IT 245123-57-3P
 RL: PEP (Physical, engineering or chemical process); PRP (Properties); RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); PROC (Process); RACT (Reactant or reagent)
 (preparation and complexation of anions by pendant N-ferrocenyl dipeptide ester deriva.)
 RN 245123-57-3 CAPLUS
 CN L-Phenylalanine, N-(ferrocenylcarbonyl)-L-phenylalanyl-, methyl ester (9CI) (CA INDEX NAME)




REFERENCE COUNT: 16 THERE ARE 16 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 151 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)


L4 ANSWER 152 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
 ACCESSION NUMBER: 1999-529142 CAPLUS
 DOCUMENT NUMBER: 131:170268
 TITLE: Condensed heterocyclic system derivatives, namely 4-amino(thio)chroman-8-carboxamides, useful as farnesyl transferase inhibitors, and their preparation and pharmaceutical compositions
 INVENTOR(S): Baudoin, Bernard; Clerc, Francois; Dereu, Norbert; El-Ahmad, Youssef; Hardy, Jean-Claude; Jimonet, Patrick; Le Brun, Alain
 PATENT ASSIGNEE(S): Rhone-Poulenc Rorer S.A., Fr.
 SOURCE: FCT Int. Appl., 228 pp.
 CODEN: FIHXD2
 DOCUMENT TYPE: Patent
 LANGUAGE: French
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 9941248	A1	19990819	WO 1999-FR298	19990211
W: AL, AU, BA, BB, BG, BR, CA, CN, CU, CZ, EE, GD, GE, HR, HU, ID, IL, IN, IS, JP, KP, KR, LC, LR, LT, LV, MG, MK, MN, MX, NO, NZ, PL, RO, SG, SI, SK, SL, TR, TT, UA, US, UZ, VN, YU, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM				
RW: CH, GM, KE, LS, MW, SD, SZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG				
FR 2774987	A1	19990820	FR 1998-1762	19980213
FR 2774987	B1	20000317		
ZA 5901073	A	19990810	ZA 1999-1073	19990210
CA 2321218	AA	19990819	CA 1999-2321218	19990211
AU 5924287	A1	19990830	AU 1999-24287	19990211
EP 1054882	A1	20001129	EP 1999-903732	19990211
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, PT, IE, FI				
JP 2002503659	T2	20020205	JP 2000-531443	19990211
PRIORITY APPL. INFO.:			FR 1998-1762 A	19980213
			US 1998-81577P	P 19980414
			WO 1999-FR298	W 19990211

OTHER SOURCE(S):
 GI MARPAT 131:170268



I



II


AB The invention concerns novel title compds. I, their preparation, pharmaceutical compns., and use for preparing medicines (wherein R1 = COCH(NH₂)CH₂SH,

L4 ANSWER 152 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)
 CH₂CH(NH₂)CH₂SH, or CHR1R12; R2 = H, alkyl, aralkyl, aryl, alkylcarbonyl, aralkylcarbonyl, arylcarbonyl, or heteroaralkyl; R3 = H, halo, alkyl, aryl, aralkyl; R4 = CHR13R14; R5 = H, COR15; R6 = H, alkyl, aryl, aralkyl; R7, R8 = H, alkyl, aryl, or aralkyl; X = O, S, S(O), S(O)₂; R11 = (un)substituted heterocyclyl; R12 = H, alkyl, aryl, aralkyl; R13, R14 = H, (un)substituted aryl or heteroaryl (both = H); R15 = OH, alkoxy, NH₂, aralkylamino, alkylamino, NHCH(CO₂R16)CH₂CH₂SM₆; R16 = H, alkyl, including racemates stereoisomers, and salts). The compds. are inhibitors of farnesyl transferase, and as such are potent antitumor and antileukemic agents. Examples include 86 syntheses, evaluation of the inhibition of farnesylation of K-ras in vitro, activity against human tumor cells (HCT116) in vitro [IC₅₀ = 0.1 nM to 100 μM in both cases], and 3 pharmaceutical formulations. For instance, 4-chromanone underwent a sequence of reductive amination to the 4-amino compd. (50.4%), N-protection by BOC (95.3%), 8-lithiation and carboxylation (53.2%), peptide coupling with H-L-Phe-NH₂ (93%), removal of BOC (28%), reductive amination with S-(triphenylmethyl)-N-BOC-L-cysteine (87%), and final deprotection with Et₃SiH and CF₃CO₂H, to give 2 diastereomers of title compd. II, isolated as the di(trifluoroacetate) salts.

IT 238764-09-5P
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)
 (intermediate; preparation of aminochromancarboxamides and -thiochromancarboxamides as farnesyl transferase inhibitors)

RN 238764-09-5 CAPLUS
 CN Carbamic acid, [8-[[[(1S)-2-amino-2-oxo-1-(phenylmethyl)ethyl]amino]carbon yll]-3,4-dihydro-2H-1-benzopyran-4-yl]-, 1,1-dimethylethyl ester (9CI) (CA INDEX NAME)

Absolute stereochemistry.



REFERENCE COUNT: 1 THERE ARE 1 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 153 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
 ACCESSION NUMBER: 1999:505653 CAPLUS
 DOCUMENT NUMBER: 131:153742
 TITLE: Bradykinin antagonists containing pentafluorophenylalanine, and therapeutic use
 INVENTOR(S): Stewart, John M.; Gera, Lajos
 PATENT ASSIGNEE(S): University Technology Corporation, USA
 SOURCE: U.S.; 6 pp.
 CODEN: USXXAM
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 5935932	A	19990810	US 1998-96716	19980612
PRIORITY APPL. INFO.:			US 1997-49571P	P 19970613

OTHER SOURCE(S): MARPAT 131:153742

AB Bradykinin antagonists containing pentafluorophenylalanine which are therapeutically useful are provided. Also provided are methods to antagonize bradykinin receptors in a mammal in need of such antagonism, comprising administering a bradykinin antagonist containing pentafluorophenylalanine. Further provided are methods to treat inflammation in a mammal in need of such inhibition, comprising administering a bradykinin antagonist containing pentafluorophenylalanine. Lastly, a method to treat cancer in a mammal in need of such inhibition comprises administering a bradykinin antagonist containing pentafluorophenylalanine.

IT 236729-63-8P

RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)
 (bradykinin antagonists containing pentafluorophenylalanine, and therapeutic use)


RN 236729-63-8 CAPLUS

CN L-Arginine, D-arginyl-L-arginyl-L-prolyl- (4R)-4-hydroxy-L-prolylglycyl-3- (2-thienyl)-L-alanyl-L-seryl- (3R)-1,2,3,4-tetrahydro-3-isoquinolinecarboxyl-2,3,4,5,6-pentafluoro-L-phenylalanyl- (9CI) (CA INDEX NAME)


Absolute stereochemistry.

L4 ANSWER 153 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)

PAGE 1-A




PAGE 1-B



L4 ANSWER 153 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)

PAGE 2-A



REFERENCE COUNT: 32 THERE ARE 32 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 154 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
 ACCESSION NUMBER: 1999:487274 CAPLUS
 DOCUMENT NUMBER: 131:116520
 TITLE: Preparation of phenylalanine derivatives as pharmaceutical agents
 INVENTOR(S): Head, John Clifford; Archibald, Sarah Catherine; Warrelow, Graham John; Porter, John Robert
 PATENT ASSIGNEE(S): Celltech Therapeutics Limited, UK
 SOURCE: PCT Int. Appl., 65 pp.
 CODEN: PIXXD2
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 9937618	A1	19990729	WO 1999-GB279	19990127
W:	AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM			
RW:	GH, GM, KE, LS, MW, SD, SZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG			
US 6329372	B1	20011211	US 1999-237060	19990126
AU 9924320	A1	19990809	AU 1999-24320	19990127
EP 1051399	A1	20001115	EP 1999-903798	19990127
R:	AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, FI			
JP 2002501051	T2	20020115	JP 2000-528542	19990127
US 2002035127	A1	20020321	US 2001-964161	20010926
PRIORITY APPL. INFO.:			GB 1998-1674	A 19980127
			GB 1998-26669	A 19981203
			US 1999-237060	A1 19990126
			WO 1999-GB279	W 19990127

OTHER SOURCE(S): MARPAT 131:116520

AB Phenylalanine derivs. 4-[R1(Alk1)R1a]C6H2RaRb(Alk2)mCHRR2NR3COHet [R is a carboxylic acid or derivative; R1 = H, OH, alkoxy or optionally substituted cycloaliph., polycycloaliph., heterocycloaliph., polyheterocycloaliph., arom. or heteroarom. group; Alk1 = optionally substituted aliphatic or heteroaliph. chain; L1 is a linker atom or group; r, s = 0, 1; Ra, Rb = -L2(CH2)pL3Rc, where L2, L3 = a covalent bond or linker atom or group; p = 0, 1; q = 1-3; Rc = H, halo, alkyl, OH, alkoxy, etc.; Alk2 = alkylene; m = 0, 1; R2 = H, Me; R3 = H, alkyl; Het is an optionally substituted heteroarom. group] and their salts, solvates, hydrates and N-oxides were prepared as pharmaceutical agents. Thus, N-(2-chloronicotinoyl)-N'-(3,5-dichloro-4-picoyl)-L-4-aminophenylalanine was prepared by coupling reaction of N-(3,5-dichloro-4-picoyl)-L-4-aminophenylalanine Me ester with 2-chloronicotinoyl chloride followed by ester hydrolysis. Title compds. were tested for inhibition of integrin-dependent cell adhesion and generally have IC50 values in the α4β1 and α4β7 assays of 1μM and below.


IT 232617-85-5P

RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)
 (preparation of phenylalanine derivs. as pharmaceutical agents)

RN 232617-85-5 CAPLUS

L4 ANSWER 154 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)
 CN L-Tyrosine, O-[(2,6-dichlorophenyl)methyl]-N-(4-quinolinylcarbonyl)- (9CI)
 (CA INDEX NAME)

Absolute stereochemistry.




REFERENCE COUNT: 4 THERE ARE 4 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 155 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
 ACCESSION NUMBER: 1999:484863 CAPLUS
 DOCUMENT NUMBER: 131:266894
 TITLE: The Opioid μ Agonist/ δ Antagonist DIPP-NH2[Y] Produces a Potent Analgesic Effect, No Physical Dependence, and Less Tolerance than Morphine in Rats
 AUTHOR(S): Schiller, Peter W.; Fundytus, Marian E.; Merovitz, Lissa; Weltrowska, Grazyna; Nguyen, Thi M.-D.; Lemieux, Carole; Chung, Nga H.; Coderre, Terence J.
 CORPORATE SOURCE: Laboratory of Chemical Biology and Peptide Research and Pain Mechanisms Laboratory, Clinical Research Institute of Montreal, Montreal, QC, H2W 1R7, Can.
 SOURCE: Journal of Medicinal Chemistry (1999), 42(18), 3520-3526
 CODEN: JMCMAR; ISSN: 0022-2623
 PUBLISHER: American Chemical Society
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 AB Opioid compds. with mixed μ agonist/ δ antagonist properties are expected to be analgesics with low propensity to produce tolerance and dependence. In an effort to strengthen the μ agonist component of the mixed μ agonist/ δ antagonist H-Tyr-Tic-Phe-Phe-NH2 (TIPP-NH2), analogs containing structurally modified tyrosine residues in place of Tyr1 were synthesized. Among the prepared compds., H-Dmt-Tic-Phe-Phe-NH2 (DIPP-NH2; Dmt = 2',6'-dimethyltyrosine) and H-Dmt-Ticf[CH2NH]Phe-Phe-NH2 (DIPP-NH2[Y]) retained a mixed μ agonist/ δ antagonist profile, as determined in the guinea pig ileum and mouse vas deferens assays, whereas H-Tmt-Tic-Phe-Phe-NH2 (Tmt = N,2',6'-trimethyltyrosine) was a partial μ agonist/ δ antagonist and H-Tmt-Ticf[CH2NH]Phe-Phe-NH2 was a μ antagonist/ δ antagonist. DIPP-NH2[Y] showed binding affinities in the subnanomolar range for both μ and δ receptors in the rat brain membrane binding assays, thus representing the first example of a balanced μ agonist/ δ antagonist with high potency. In the rat tail flick test, DIPP-NH2[Y] given i.c.v. produced a potent analgesic effect (ED50 = 0.04 μ g), being about 3 times more potent than morphine (ED50 = 0.11 μ g). It produced less acute tolerance than morphine but still a certain level of chronic tolerance. Unlike morphine, DIPP-NH2[Y] produced no phys. dependence whatsoever upon chronic administration at high doses (54.5 μ g/h) over a 7-day period. In conclusion, DIPP-NH2[Y] fulfills to a large extent the expectations based on the mixed μ agonist/ δ antagonist concept with regard to analgesic activity and the development of tolerance and dependence.
 IT 245538-28-7P
 RL: BAC (Biological activity or effector, except adverse); BPR (Biological process); BSU (Biological study, unclassified); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); PROC (Process); USES (Uses)
 (opioid μ agonist/ δ antagonist DIPP-NH2[Y] produces a potent analgesic effect and no phys. dependence and less tolerance than morphine in Rats in relation to structure)
 RN 245538-28-7 CAPLUS
 CN L-Phenylalaninamide, N,2,6-trimethyl-L-tyrosyl-(3S)-1,2,3,4-tetrahydro-3-isoquinolinecarbonyl-L-phenylalanyl- (9CI) (CA INDEX NAME)


Absolute stereochemistry.

L4 ANSWER 155 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)




REFERENCE COUNT: 25 THERE ARE 25 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 156 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
 ACCESSION NUMBER: 1999:470677 CAPLUS
 DOCUMENT NUMBER: 131:228622
 TITLE: Replacement of the quinoline system in 2-phenyl-4-quinolinecarboxamide NK-3 receptor antagonists
 AUTHOR(S): Giardina, G. A. M.; Artico, M.; Cavignera, S.; Cerri, A.; Consolandi, E.; Gagliardi, S.; Graziani, D.; Grugni, M.; Hay, D. W. P.; Luttman, M. A.; Mena, R.; Raveglia, L. F.; Rigolio, R.; Sarau, H. M.; Schmidt, D. B.; Zanon, G.; Farina, C.
 CORPORATE SOURCE: Department of Medicinal Chemistry, SmithKline Beecham S.p.A., Milan, 20021, Italy
 SOURCE: Farmaco (1999), 54(6), 364-374
 CODEN: FRMCES; ISSN: 0014-827X
 PUBLISHER: Elsevier Science S.A.
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 OTHER SOURCE(S): CASREACT 131:228622
 GI



AB Results from a medicinal chemical approach aimed at replacing the quinoline ring system in the potent and selective human neurokinin-3 (hNK-3) receptor antagonists (RS)-I (R = MeOCO, Et; X1 = C; X2 = N), (R)-II (R = MeOCO; X1 = C; X2 = N) and (S)-I (R = Et; X1 = C; X2 = N) are discussed. The data give further insight upon the potential NK-3 pharmacophore. In particular, it is highlighted that both the benzene-condensed ring and the quinoline nitrogen are crucial determinants for optimal binding affinity to the hNK-3 receptor. Some novel compds. I (R = MeOCO; X1 = X2 = C (II); R = MeOCO; X1 = X2 = N) and III (R1R2 = N:CHCH:N, CH2CH2CH2CH2) maintained part of the binding affinity to the receptor and compound II, featuring the naphthalene ring system, appears to be suitable for further modifications; it offers the option to introduce electron-withdrawing groups at position 2 and 4, conferring on the ring an overall electron-deficiency similar to that of the quinoline.
 IT 174635-51-9
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); BIOL (Biological study)
 (preparation, binding affinity and structure-activity relationship of NK-3 receptor antagonists)
 RN 174635-51-9 CAPLUS
 CN Benzenecetic acid, α -[[(2-phenyl-4-quinolinyl)carbonyl]amino]-, methyl ester (9CI) (CA INDEX NAME)

L4 ANSWER 156 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)



REFERENCE COUNT: 26 THERE ARE 26 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 157 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN


ACCESSION NUMBER: 1999:454977 CAPLUS
DOCUMENT NUMBER: 131:194017
TITLE: Cyclic AMP phosphodiesterase inhibition by coumarins and furanocoumarins
AUTHOR(S): Sardari, Soroush; Nishibe, S.; Horita, K.; Nikaido, T.; Daneshmandi, M.
CORPORATE SOURCE: School Pharmaceutical Sciences, Health Sciences Univ. Hokkaido, Hokkaido, Japan
SOURCE: Pharmazie (1999), 54(7), 554-556
CODEN: PHARAT; ISSN: 0031-7144
PUBLISHER: Govi-Verlag Pharmazeutischer Verlag
DOCUMENT TYPE: Journal
LANGUAGE: English

AB The phosphodiesterase inhibitory property of several synthetic as well as natural coumarins is reported. In comparison to caffeine, many of the coumarins tested showed strong inhibitory activity on phosphodiesterase. A comparison among the coumarins revealed that lipophilicity and the presence of a phenolic hydroxyl group at position 6 or 7 are 2 important factors in the phosphodiesterase inhibitory activity of coumarins.

IT 222640-31-5
RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); BIOL (Biological study)
(CAMP phosphodiesterase inhibition by coumarins and furanocoumarins)

RN 222640-31-5 CAPLUS
CN L-Phenylalanine, N-[(2-oxo-2H-furo[2,3-h]-1-benzopyran-8-yl)carbonyl]-, ethyl ester (9CI) (CA INDEX NAME)

Absolute stereochemistry.




REFERENCE COUNT: 14 THERE ARE 14 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT


L4 ANSWER 158 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 1999:439520 CAPLUS
DOCUMENT NUMBER: 131:102538
TITLE: Preparation of quinoline, isoquinoline, cinnoline and tyrosine derivatives as antiinflammatory and anti-allergy agents
INVENTOR(S): Nakao, Toyoo; Takei, Masao; Fukamachi, Hiromi; Ohashi, Hiroshi
PATENT ASSIGNEE(S): Kirin Brewery Co., Ltd., Japan
SOURCE: Jpn. Kokai Tokkyo Koho, 40 pp.
CODEN: JKXXAF
DOCUMENT TYPE: Patent
LANGUAGE: Japanese
FAMILY ACC. NUM. COUNT: 1
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 11189586	A2	19990713	JP 1997-358639	19971225
PRIORITY APPLN. INFO.: JP 1997-358639 19971225				
OTHER SOURCE(S): MARPAT 131:102538				
GI				



I




II

L4 ANSWER 158 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)

study, unclassified); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation)
(prepn. of quinoline, isoquinoline, cinnoline and amino acid derivs. as antiinflammatory and anti-allergy agents)

RN 231634-29-0 CAPLUS
CN L-Tyrosine, O-(6,7-dimethoxy-4-quinolinyl)-N-(3-quinolinylcarbonyl)-, methyl ester (9CI) (CA INDEX NAME)


Absolute stereochemistry.



AB Title compds. [I; T = O at 4, 3 position; R1 = H, 3-I, 3-Cl, 3-F, 3-OMe; R4 = H, OMe; R5 = H, OMe; G = N, CH; E = CH, N; D = CH, N; R2 = Et, Me; n = 0, 1; p = 1; O; X = Bond, CH2CH2, CH:CH, O, CH2, (CH2)10, (CH2)3, (CH2)6; Z = H, Bu-t, (un)substituted benzene, 1-naphthyl, 2-naphthyl, 3-quinolinyl] are prepared as antiinflammatory agents and anti-allergy agents. Thus, title compound II was prepared from reaction product of isovanillic acid, cyclopentyl bromide, acetic acid, (triphenylphosphoranylidene)-, Me ester, and L-Tyrosine, O-(1,1-dimethylethyl)-, Me ester with addition cyclization reaction product of 3,4-dimethoxyaniline and propanedioic acid, (ethoxymethylene)-, di-Et ester (EtOCCOC(=CHOC(=O)Et)COOEt).

IT 231634-29-0
RL: BAC (Biological activity or effector, except adverse); BSU (Biological


L4 ANSWER 159 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
 ACCESSION NUMBER: 1999:405937 CAPLUS
 DOCUMENT NUMBER: 131:199816
 TITLE: Metal complexes of biologically important ligands. CXIV. Ferrocenyl-oxazolones as N and C donors in Pd(II), Pt(II) and Ir(III) complexes and ferrocenyl-dipeptides
 AUTHOR(S): Bauer, Werner; Polborn, Kurt; Beck, Wolfgang
 CORPORATE SOURCE: Institut für Anorganische Chemie, Ludwig-Maximilians-Universität, München, D-81377, Germany
 SOURCE: Journal of Organometallic Chemistry (1999), 579(1-2), 269-279
 CODEN: JORCAI; ISSN: 0022-328X
 PUBLISHER: Elsevier Science S.A.
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 GI



I
 II

AB 2-Ferrocenyl-4R-5(4H)-oxazolones, e.g. I, were obtained from N-ferrocenyl-α-amino acids and function as N donors in dichloro-phosphine-palladium(II) and platinum(II) complexes. The reaction of ferrocenyl-oxazolone and ferrocenyl-bis(oxazolone) with [Cp*IrCl2]2 afforded trimetallic and pentametallic complexes, e.g. II, with a C,N bridging oxazolone. Ring opening of the ferrocenyl-oxazolones with α-amino acid esters gave N-ferrocenyl-dipeptide esters. In the ferrocene bis(dipeptides) the two peptide esters are aligned parallel by hydrogen bonding. The structures of platinum complex of ferrocenyl-oxazolone and II were determined by x-ray diffraction.
 IT 181589-81-1P
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)
 (preparation and saponification of)
 RN 181589-81-1 CAPLUS
 CN Ferrocene, [1-((1S)-2-methoxy-2-oxo-1-(phenylmethyl)ethylamino)carbonyl]-

L4 ANSWER 159 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)
 (9CI) (CA INDEX NAME)



REFERENCE COUNT: 56 THERE ARE 56 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT


L4 ANSWER 160 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
 ACCESSION NUMBER: 1999:404983 CAPLUS
 DOCUMENT NUMBER: 131:45107
 TITLE: Preparation of peptidyl antipicornaviral compounds
 INVENTOR(S): Webber, Stephen E.; Dragovich, Peter S.; Prins, Thomas J.; Littlefield, Ethel S.; Marakovits, Joseph T.; Babine, Robert E.
 PATENT ASSIGNEE(S): Agouron Pharmaceuticals, Inc., USA
 SOURCE: PCT Int. Appl., 187 pp.
 CODEN: PIXXD2
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 9931122	A1	19990624	WO 1998-US26583	19981215
W: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, RW: GH, GM, KE, LS, MW, SD, SZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, CA, GN, GW, ML, MR, NE, SN, TD, TG				
US 5962407	A	19991005	US 1997-991739	19971216
CA 2312940	AA	19990624	CA 1998-2312940	19981215
AU 9918262	A1	19990705	AU 1999-18262	19981215
AU 762682	B2	20030703		
EP 1037905	A1	20000927	EP 1998-963184	19981215
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, FI				
BR 9813651	A	20001003	BR 1998-13651	19981215
JP 2002508389	T2	20020319	JP 2000-539045	19981215
NO 2000003067	A	20000815	NO 2000-3067	20000615
PRIORITY APPLN. INFO.: US 1997-991739 A 19971216 WO 1998-US26583 W 19981215				

OTHER SOURCE(S): MARPAT 131:45107
 AB Picornaviral 3C protease inhibitors R8R4NCR3R6C((M)NR7CR2R5CR1:CZ21 [M = O, S; R1 = H, F, alkyl, OH, SH, O-alkyl group; R2, R5 = H, alkyl, X-Y1-A1(B1)D1, X-Y2-A2(B2)D2 (X = :CH, :CF, CH2, CF2, CHF, S; Y1, Y2 = :CH, :CF, or X and Y1 or Y2 may form a ring; A1, A2 = C, CH, CF, S, P, Se, N, etc.; D1 and D2 are moieties with a lone pair of electrons capable of forming a hydrogen bond; R1, R2 = H, F, alkyl, cycloalkyl, heterocycloalkyl, aryl, heteroaryl, etc.); R3, R6 = H, F, alkyl, cycloalkyl, heterocycloalkyl, aryl, heteroaryl, CHO, OH, SH, etc.; R4 is any suitable organic moiety or R4 and R3 or R6 may form a ring; R7, R8 = H, alkyl, cycloalkyl, heterocycloalkyl, aryl, heteroaryl, etc. or R4 and R8 may form a ring; Z, Z1 are H, F, alkyl, cycloalkyl, heterocycloalkyl, aryl, heteroaryl, etc.] were prepared. Thus, Et 3-(Cbz-L-N-Me-Phe-L-Gln)-E-propenone (Cbz = benzyloxycarbonyl) was prepared and showed Ki >100 μM for inhibition of Rhinovirus protease.
 IT 227613-70-99
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)
 (preparation of peptidyl antipicornaviral compds.)
 RN 227613-70-9 CAPLUS
 CN 2-Heptenoic acid, 7-amino-4-[[[(2S)-2-[[[(3S)-2-[(ethylthio)carbonyl]-1,2,3,4-tetrahydro-3-isoquinolinyl]carbonyl]methylamino]-1-oxo-3-

L4 ANSWER 160 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)
 phenylpropylamino]-7-oxo-, ethyl ester, (2E,4S)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.
 Double bond geometry as shown.



REFERENCE COUNT: 3 THERE ARE 3 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 161 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
 ACCESSION NUMBER: 1999:396525 CAPLUS
 DOCUMENT NUMBER: 131:200018
 TITLE: Synthesis of conformationally constrained peptides using the Heck reaction
 AUTHOR(S): Wright, David E.
 CORPORATE SOURCE: R.W. Johnson Pharmaceutical Research Institute, San Diego, CA, 92121, USA
 SOURCE: Peptides: Frontiers of Peptide Science, Proceedings of the American Peptide Symposium, 15th, Nashville, June 14-19, 1997 (1999), Meeting Date 1997, 279-280.
 Editor(s): Tam, James P.; Kaumaya, Pravin T. P.
 Kluwer: Dordrecht, Neth.
 CODEN: 67UCAR
 CONFERENCE: Conference
 LANGUAGE: English
 DOCUMENT TYPE: Conference
 AB A symposium report. Heck reaction was used to develop cyclic opiate peptides. Cyclization was carried out using a p-iodophenylalanine residue and an alkene group.
 IT 241814-55-1
 RL: RCT (Reactant); RACT (Reactant or reagent)
 (preparation of conformationally constrained cyclopeptides via Heck cyclization)
 RN 241814-55-1 CAPLUS
 CN L-Phenylalaninamide, N-(1-oxo-4-pentenyl)-L-tyrosyl-(3S)-1,2,3,4-tetrahydro-3-isoquinolinecarbonyl-L-phenylalanyl-4-iodo- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



REFERENCE COUNT: 5 THERE ARE 5 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT


L4 ANSWER 162 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
 ACCESSION NUMBER: 1999:368538 CAPLUS
 DOCUMENT NUMBER: 131:153427
 TITLE: Proteasome inhibitors: a novel class of potent and effective antitumor agents
 AUTHOR(S): Adams, Julian; Palombella, Vito J.; Sausville, Edward A.; Johnson, Jill; Destree, Antonia; Lazarus, Douglas D.; Maas, Jochen; Pien, Christine S.; Prakash, Samuel; Elliott, Peter J.
 CORPORATE SOURCE: ProScript, Inc., Cambridge, MA, 02139, USA
 SOURCE: Cancer Research (1999), 59(11), 2615-2622
 CODEN: CNREAS; ISSN: 0008-5472
 PUBLISHER: AACR Subscription Office
 DOCUMENT TYPE: Journal
 LANGUAGE: English

AB The ubiquitin-proteasome pathway plays a critical role in the regulated degradation of proteins involved in cell cycle control and tumor growth. Dysregulating the degradation of such proteins should have profound effects on tumor growth and cause cells to undergo apoptosis. To test this hypothesis, we developed a novel series of proteasome inhibitors, exemplified by PS-341, which we describe here. As determined by the National Cancer Institute in vitro screen, PS-341 has substantial cytotoxicity against a broad range of human tumor cells, including prostate cancer cell lines. The PC-3 prostate cell line was, therefore, chosen to further examine the antitumor activity of PS-341. In vitro, PS-341 elicits proteasome inhibition, leading to an increase in the intracellular levels of specific proteins, including the cyclin-dependent kinase inhibitor, p21. Moreover, exposure of such cells to PS-341 caused them to accumulate in the G2-M phase of the cell cycle and subsequently undergo apoptosis, as indicated by nuclear condensation and poly(ADP-ribose) polymerase cleavage. Following weekly i.v. treatment of PS-341 to mice bearing the PC-3 tumor, a significant decrease (60%) in tumor burden was observed in vivo. Direct injection of PS-341 into the tumor also caused a substantial (70%) decrease in tumor volume with 40% of the drug-treated mice having no detectable tumors at the end of the study. Studies also revealed that i.v. administration of PS-341 resulted in a rapid and widespread distribution of PS-341, with highest levels identified in the liver and gastrointestinal tract and lowest levels in the skin and muscle. Modest levels were found in the prostate, whereas there was no apparent penetration of the central nervous system. An assay to follow the biol. activity of the PS-341 was established and used to determine temporal drug activity as well as its ability to penetrate tissues. As such, PS-341 was shown to penetrate PC-3 tumors and inhibit intracellular proteasome activity 1.0 h after i.v. dosing. These data illustrate that PS-341 not only reaches its biol. target but has a direct effect on its biochem. target, the proteasome. Importantly, the data show that inhibition of this target site by PS-341 results in reduced tumor growth in murine tumor models. Together, the results highlight that the proteasome is a novel biochem. target and that inhibitors such as PS-341 represent a unique class of antitumor agents. PS-341 is currently under clin. evaluation for advanced cancers.

IT 179324-59-5
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); PRP (Properties); THU (Therapeutic use); BIOL (Biological study); USRS (Uses)
 (development of potent, selective and reversible dipeptide boronic acid proteasome inhibitors as antitumor agents)

L4 ANSWER 162 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)
 RN 179324-59-5 CAPLUS
 CN Boronic acid, [(1R)-3-methyl-1-[[[(2S)-1-oxo-4-phenyl-2-[(2-quinolinylcarbonyl)amino]butyl]amino]butyl]- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



REFERENCE COUNT: 32 THERE ARE 32 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 163 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
 ACCESSION NUMBER: 1999:332252 CAPLUS
 DOCUMENT NUMBER: 131:88160
 TITLE: Enantioselective solid-phase synthesis of α-amino acid derivatives
 AUTHOR(S): O'Donnell, Martin J.; Delgado, Francisca; Pottorf, Richard S.
 CORPORATE SOURCE: Department of Chemistry, Indiana University-Purdue University at Indianapolis, Indianapolis, IN, 46202, USA
 SOURCE: Tetrahedron (1999), 55(20), 6347-6362
 CODEN: TETRAE; ISSN: 0040-4020
 PUBLISHER: Elsevier Science Ltd.
 DOCUMENT TYPE: Journal
 LANGUAGE: English

AB Wang-resin bound deriva. of glycine Schiff base esters are alkylated in the presence of quaternary ammonium salts derived from cinchonidine or cinchonine using phosphazene bases to give either enantiomer of the product α-amino acid deriva. in 51-89% ee.
 IT 197392-59-9
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or Reagent)
 (preparation and reaction of in enantioselective solid-phase synthesis of α-amino acid deriva.)
 RN 197392-59-9 CAPLUS
 CN Phenylalanine, N-(2-quinolinylcarbonyl)- (9CI) (CA INDEX NAME)



REFERENCE COUNT: 77 THERE ARE 77 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT


L4 ANSWER 164 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
 ACCESSION NUMBER: 1999:317178 CAPLUS
 DOCUMENT NUMBER: 130:325390
 TITLE: Preparation of amino acid spiro-piperidine derivatives as somatostatin agonists
 INVENTOR(S): Guo, Liangqun; Mosley, Ralph T.; Pasternak, Alexander; Patchett, Arthur A.; Yang, Lihu
 PATENT ASSIGNER(S): Merck & Co., Inc., USA
 SOURCE: PCT Int. Appl., 101 pp.
 CODEN: PIXXD2
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 9922735	A1	19990514	WO 1998-US22917	19981028
W: AL, AM, AU, AZ, BA, BB, BG, BR, BY, CA, CN, CU, CZ, EE, GD, GE, HR, HU, ID, IL, IS, JP, KG, KR, KZ, LC, LK, LR, LT, LV, MD, MG, MK, MN, MX, NO, NZ, PL, RO, RU, SG, SI, SK, SL, TJ, TM, TR, TT, UA, US, UZ, VN, YU, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM				
RW: GH, GM, KE, LS, MW, SD, SZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG				
AU 9912854	A1	19990524	AU 1999-12854	19981028
US 6117880	A	20000912	US 1998-181590	19981028
PRIORITY APPL. INFO.:				
US 1997-64422P P 19971030				
GB 1998-6697 A 19980327				
WO 1998-US22917 W 19981028				

OTHER SOURCE(S): MARPAT 130:325390
 AB Title compds. R3R4NCOCR1R1aZ1ENR2 [NR2 represents spiro(indene-1,4'-piperidine) or spiro(indole-1,4'-piperidine) or related 2,3-dihydro deriva. in which the residue at the 3-position is CH2, CHCO2R2, CO, CHCH2CO2R2, CHCONR22, NSO2R2, or CR5 (R2 and R5 = H, alkyl, arylalkyl, cycloalkyl, etc.); R1 = alkyl, aryl, arylalkyl, heteroaryl, CO2R2, etc.; R1a = H, alkyl; R3 = H, alkyl, arylalkyl, heteroarylalkyl; R4 = CHCO2R2, CHCONR2, etc.; E = SO2, CO(CR22)n (n = 0-5), C(:NCR), C(:NNO2), C(:NSO2NR22); Z1 = NH, alkylimino, hydroxyalkylimino] were prepared as somatostatin agonists. Thus, R2NCO-D-Trp-NH(CH2)5NH2 [R2N = spiro(indene-1,4'-piperidine)] was prepared from reactions of D-tryptophan Me ester, spiro(indene-1,4'-piperidine), and 1,5-pentanediamine. The compds. of the invention inhibit the binding of somatostatin to its receptor at at IC50 of about 30 pM to about 3 pM.
 IT 223770-91-0P
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)
 RN [preparation of amino acid spiro-piperidine derivs. as somatostatin agonists]
 CN 223770-91-0 CAPLUS
 L-Lysine, N-(spiro[1H-indene-1,4'-piperidin]-1'-ylcarbonyl)-D-phenylalanyl-, methyl ester (9CI) (CA INDEX NAME)

Absolute stereochemistry.


L4 ANSWER 164 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)



REFERENCE COUNT: 2 THERE ARE 2 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 165 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
 ACCESSION NUMBER: 1999:314009 CAPLUS
 DOCUMENT NUMBER: 131:129781
 TITLE: Synthesis and antipseudomonal activity of fluoroquinolonyl-penicillin derivatives
 AUTHOR(S): Tsou, Tai-Li; Tang, Shang-Tao; Wu, Jing-Ran; Hung, Yao-Wen; Liu, Yu-Tien
 CORPORATE SOURCE: Institute of Preventive Medicine, National Defense Medical Center, Taipei, Taiwan
 SOURCE: European Journal of Medicinal Chemistry (1999), 34(3), 255-259
 CODEN: EJMCAS; ISSN: 0223-5234
 PUBLISHER: Editions Scientifiques et Medicales Elsevier
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 AB A series of ampicillin and amoxicillin derivs. containing an N1-substituted-(6-fluoro-1,4-dihydro-4-oxoquinolin-3-yl) carbonyl moiety at the α-amino group were prepared and their antibacterial activities were evaluated. These derivs. displayed a broad spectrum of antibacterial activity against Gram-(+) and Gram-(-) bacteria. In comparison with the original antibiotics, some of the derivs. were more active against Pseudomonas aeruginosa strains. However, their antibacterial activities decrease when N-1 substitution was replaced with alkyl substituents. Interestingly, several products induced filamentation in three strains of P. aeruginosa.
 IT 190902-80-8P
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation)
 RN [synthesis and antipseudomonal activity of fluoroquinolonyl-penicillin derivs.]
 CN 190902-80-8 CAPLUS
 4-Thia-1-azabicyclo[3.2.0]heptane-2-carboxylic acid, 6-[[[(2R)-[[[(6,7-difluoro-1,4-dihydro-4-oxo-3-quinolinyl)carbonyl]amino]phenylacetyl]amino]-3,3-dimethyl-7-oxo-, (2S,5R,6R)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



REFERENCE COUNT: 18 THERE ARE 18 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 166 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
 ACCESSION NUMBER: 1999:250927 CAPLUS
 DOCUMENT NUMBER: 131:44415
 TITLE: Solid-Phase Synthesis of a Combinatorial Array of 1,3-Bis(acylamino)-2-butanones, Inhibitors of the Cysteine Proteases Cathepsins K and L
 AUTHOR(S): Yamashita, Dennis S.; Dong, Xiaoyang; Oh, Hye-Ja; Brook, Christopher S.; Tomaszek, Thaddeus A.; Szewczuk, Lawrence; Tew, David G.; Veber, Daniel F.
 CORPORATE SOURCE: Departments of Medicinal Chemistry Molecular Recognition and Synthetic Chemistry, SmithKline Beecham Pharmaceuticals, King of Prussia, PA, 19406, USA
 SOURCE: Journal of Combinatorial Chemistry (1999), 1(3), 207-215
 CODEN: JCCHFF; ISSN: 1520-4766
 PUBLISHER: American Chemical Society
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 AB To more rapidly prepare members of the 1,3-bis(acylamino)-2-butanone class of cysteine protease inhibitors, a solid-phase synthesis was developed. 1-Azido-3-amino-2,2-dimethoxybutane, which has two amino groups differentiated and the ketone protected as a ketal, served as a surrogate for the 1,3-diamino-2-butanone core. Thus, 1-azido-3-amino-2,2-dimethoxybutane was coupled to the BAL-resin-linked carboxylic acids derived from α-amino acid esters. Evaluation of a small combinatorial array by measuring inhibition consts. (Ki,apps) against cathepsins K, L, and B provided some structure-activity relationship trends with respect to selectivity and potency. Novel, potent inhibitors of cathepsins K and L were identified. Novel, potent inhibitors of cathepsins K and L were identified.
 IT 227178-23-6P
 RL: SPN (Synthetic preparation); PREP (Preparation)
 RN [preparation of]
 CN 227178-23-6 CAPLUS
 2-Quinolonecarboxamide, N-[[[(1S)-2-[[[1-methyl-2-oxo-3-[[[3-(2-pyridinyl)phenyl]acetyl]amino]propyl]amino]-2-oxo-1-(phenylmethyl)ethyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




REFERENCE COUNT: 33 THERE ARE 33 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 167 OF 261 CAPLUS COPYRIGHT 2004 ACS ON STN
 ACCESSION NUMBER: 1999:184268 CAPLUS
 DOCUMENT NUMBER: 130:223587
 TITLE: 1-amino-7-isoquinoline derivatives as serine protease inhibitors
 INVENTOR(S): Liebeschuetz, John Walter; Wylie, William Alexander; Wauzkwycz, Bohdan; Murray, Christopher William; Rimmer, Andrew David; Welsh, Pauline Mary; Jones, Stuart Donald; Roscoe, Jonathan Michael Ernest; Young, Stephen Clinton; Morgan, Phillip John; Camp, Nicholas Paul; Crew, Andrew Philip Austin
 PATENT ASSIGNER(S): Proteus Molecular Design Ltd., UK
 SOURCE: PCT Int. Appl., 89 pp.
 CODEN: PIXXD2
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 13
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 9911657	A1	19990311	WO 1998-GB2600	19980828
W: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GR, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UK, US, UZ, VN, YU, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, RW: GH, GM, KE, LS, MW, SD, SZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG				
AU 9888753	A1	19990322	AU 1998-88753	19980828
EP 1012166	A1	20000620	EP 1998-940425	19980828
EP 1012166	B1	20031029		
R: CH, DE, ES, FR, GB, IT, LI, NL				
US 6262069	B1	20010717	US 2000-485677	20000225
US 2002040144	A1	20020404	US 2001-865418	20010529
US 6420438	B1	20020716	US 2000-865418	20010529
US 2003216403	A1	20031120	US 2003-296245	20030514
PRIORITY APPLN. INFO.:			GB 1997-18392	A 19970829
			GB 1998-3173	A 19980213
			WO 1998-GB2600	W 19980828
			US 2000-485677	A1 20000225
			WO 2001-GB2566	W 20010612

OTHER SOURCE(S): MARPAT 130:223587
 GI



L4 ANSWER 167 OF 261 CAPLUS COPYRIGHT 2004 ACS ON STN (Continued)


AB 1-aminoisoquinoline amino acid deriva. I (R1 = H, halo, cyano, nitro, hydroxy, amino, alkoxy, alkyl, aminoalkyl, hydroxyalkyl, thiol, alkylthio, aminosulfonyl, alkoxyalkyl, alkoxyacetyl, acyloxymethoxycarbonyl or alkylamino (optionally substituted); R2 = H, halo, Me, amino, hydroxy, or oxo; and R is X-X-Y(R7)-L-Lp(D)n, where each X independently is a C, N, O or S atom or a CO, CR1, CR12 or NR1 group; Y is a nitrogen atom or a CR1 group or Y and L taken together form a cyclic group; R7 is a lipophilic group selected from alkyl, alkenyl, mono- or bi-cycloalkyl, aryl, heteroaryl, mono- or bicycloalkyl, mono- or bicycloalkylalkenyl, aralkyl, heteroaryl-alkyl, arylalkenyl, heteroarylalkenyl, all optionally substituted by a group R1; L is an organic linker group containing 1 to 5 backbone atoms selected from C, N, O and S, or a branched alkyl or cyclic group; Lp is a lipophilic organic group selected from alkyl, heterocyclic, alkenyl, alkaryl, cycloalkyl, polycycloalkyl, cycloalkenyl, aryl, aralkyl or haloalkyl group or a combination of two or more such groups optionally substituted by one or more of oxo, thio, aza or R1 groups; D is a hydrogen bond donor group; and n is 0, 1, or 2) or their 3,4-dihydro deriva. were prepared as serine protease inhibitors. Thus, 1-aminoisoquinolin-7-oyl-D-phenylglycine-4-methoxybenzylamide was prepared by amidation of Boc-D-phenylglycine with 4-methylbenzylamine, followed by deprotection and coupling with 1-aminoisoquinoline-7-carboxylic acid trifluoroacetate.

IT 221049-79-2P
 RL: SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)
 (preparation of aminoisoquinoline peptidyl deriva. as serine protease inhibitors)

RN 221049-79-2 CAPLUS

CN Glycine, (2R)-N-[(1-amino-7-isoquinolinyl)carbonyl]-2-phenylglycyl-3-(2-naphthalenyl)-D-alanyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



REFERENCE COUNT: 12 THERE ARE 12 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 169 OF 261 CAPLUS COPYRIGHT 2004 ACS ON STN
 ACCESSION NUMBER: 1999:148078 CAPLUS
 DOCUMENT NUMBER: 130:279176
 TITLE: Antifungal activity of 2'-substituted furanocoumarins and related compounds
 AUTHOR(S): Sardari, Soroush; Able, M.; Micetich, R. G.; Danesh-Talab, M.
 CORPORATE SOURCE: Faculty Pharmacy Pharmaceutical Sciences, University Alberta, Edmonton, AB, Can.
 SOURCE: Pharmazie (1999), 54(2), 156-158
 CODEN: PHARAT; ISSN: 0031-7144
 PUBLISHER: Gavi-Verlag Pharmazeutischer Verlag
 DOCUMENT TYPE: Journal
 LANGUAGE: English


AB The antifungal activity of several 2'-substituted furanocoumarins and 7-substituted coumarins against Candida albicans, Cryptococcus neoformans, Saccharomyces cerevisiae, and Aspergillus niger was tested. Strong antifungal activities were shown by angelicin, 2'-NO2-furanocoumarin as well as the coumarins containing propylamine, decylamine, hexadecylamine, and 11-amino-undecanoic acid Et ester. The 2'-carboxamido-furanocoumarins containing propylamine, decylamine, and hexadecylamine as amino component had also antifungal activity. The min. inhibitor concns. were listed.

IT 22640-31-5
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); BIOL (Biological study)
 (antifungal activity of 2'-substituted furanocoumarins and related compds.)

RN 22640-31-5 CAPLUS

CN L-Phenylalanine, N-[(2-oxo-2H-furo[2,3-h]-1-benzopyran-8-yl)carbonyl]-, ethyl ester (9CI) (CA INDEX NAME)

Absolute stereochemistry.



REFERENCE COUNT: 12 THERE ARE 12 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 168 OF 261 CAPLUS COPYRIGHT 2004 ACS ON STN
 ACCESSION NUMBER: 1999:150027 CAPLUS
 DOCUMENT NUMBER: 130:311633
 TITLE: Synthesis and antibacterial activity of penicillin derivatives containing an (N1-substituted-6-fluoro-4-oxoquinolin-3-yl)carbonyl moiety
 AUTHOR(S): Tsou, Tai-Li; Tang, Shang-Tao; Chang, Shu-Lin; Wu, Jing-Ran; Tarn, Lih-Jeng; Wu, Ching-Shei
 CORPORATE SOURCE: Institute of Preventive Medicine, National Defense Medical Center, Taipei, Taiwan
 SOURCE: Chinese Pharmaceutical Journal (Taipei) (1998), 50(6), 369-383
 CODEN: CPHJRP; ISSN: 1016-1015
 PUBLISHER: Pharmaceutical Society of Republic of China
 DOCUMENT TYPE: Journal
 LANGUAGE: English


AB A series of ampicillin and amoxicillin deriva. containing an N1-substituted (6-fluoro-1,4-dihydro-4-oxoquinolin-3-yl)carbonyl moiety at the α-amino group were prepared and their antibacterial activities were evaluated. These deriva. displayed a broad spectrum of antibacterial activity against Gram (+) and Gram (-) bacteria. When comparison to the original antibiotics, some of the deriva. were more active against Pseudomonas aeruginosa, Streptococcus pyogenes, and β-lactamase producing Staphylococcus aureus strains. The MICs ranged from 3.2 to 13 μg/mL for P. aeruginosa and from 0.0062 to 0.05 μg/mL for St. pyogenes. Their antibacterial activities decreased when the N-1 position was replaced with alkyl substituents.

IT 190902-80-8P
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); PRP (Properties); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation)
 (preparation and antibacterial activity of 6-fluoroquinolonyl penicillin deriva.)

RN 190902-80-8 CAPLUS

CN 4-Thia-1-azabicyclo[3.2.0]heptane-2-carboxylic acid, 6-[[[(2R)-[(6,7-difluoro-1,4-dihydro-4-oxo-3-quinolinyl)carbonyl]amino]phenylacetyl]amino]-3,3-dimethyl-7-oxo-, (2S,5R,6R)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.





REFERENCE COUNT: 27 THERE ARE 27 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 170 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
 ACCESSION NUMBER: 1999:139868 CAPLUS
 DOCUMENT NUMBER: 130:196958
 TITLE: Preparation of 3-tert-butyl-L-tyrosinamide-containing peptides and related compounds exhibiting a motilin receptor antagonism
 INVENTOR(S): Kotake, Ken-ichiro; Kozono, Toshiro; Sato, Tsutomu; Takanashi, Hisanori
 PATENT ASSIGNEE(S): Chugai Sanyaku Kabushiki Kaisha, Japan
 SOURCE: PCT Int. Appl., 144 pp.
 CODEN: PIXXD2
 DOCUMENT TYPE: Patent
 LANGUAGE: Japanese
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 9909053	A1	19990225	WO 1998-JP3627	19980814
W:	AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, ES, FI, GB, GE, GH, GM, HR, HU, ID, IL, IS, KE, KG, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM			
RW:	GH, GM, KE, LS, MW, SD, SZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, CA, GN, GW, ML, MR, NE, SN, TD, TG			
TM 460478	B	20011021	TM 1998-87113211	19980811
CA 2301687	AA	19990225	CA 1998-2301687	19980814
AU 9886490	A1	19990208	AU 1998-86490	19980814
AU 741216	B2	20011129		
JP 2000044595	A2	20000215	JP 1998-229586	19980814
EP 1006122	A1	20000607	EP 1998-937826	19980814
R:	AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LI, LU, NL, SE, MC, PT, IE, FI			
US 6255285	B1	20010703	US 2000-485620	20000215
PRIORITY APPL. INFO.:			JP 1997-255879	A 19970815
			JP 1998-186802	A 19980528
			WO 1998-JP3627	W 19980814
OTHER SOURCE(S):		MARPAT 130:196958		
GI				

L4 ANSWER 170 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)



AB Phenethylamine deriva. represented by general formula [I]; wherein A represents an amino acid or α -substituted amino acid residue; R1 represents R6CO, (un)substituted C2-7 linear or branched alkyl, C3-8 alkenyl, or C3-8 alkynyl; R2 represents hydrogen, C1-3 linear or branched alkyl; R3 represents COR7, (un)substituted C1-5 linear or branched alkyl, C2-5 alkenyl, or C2-5 alkynyl; R4 represents H, C1-6 linear or branched alkyl, C2-6 alkenyl, C2-6 alkynyl, etc.; R5 represents hydroxy or C1-4 n-alkoxy; R6 represents (un)substituted C1-6 linear or branched alkyl, C2-7 alkenyl, or C2-7 alkynyl, optionally benzene- or heterocyclic ring-condensed C3-7 cycloalkyl, (un)substituted C6-12 aromatic ring, (un)substituted C3-12 (un)saturated heterocyclic ring, (un)substituted NH2, (un)substituted linear or branched C1-5 alkoxy, C2-6 alkenyloxy, C2-6 alkynoxy, etc.; and R7 represents H, (un)substituted C1-5 linear or branched alkyl, C3-7 cycloalkyl, (un)substituted NH2, OH, linear or branched alkyl C1-6 alkoxy, or C3-7 cycloalkyloxy are prepared. Also claimed are a motilin receptor antagonist, an inhibitor of digestive tract motility, and a remedy for high blood motilin. They are also useful for the treatment of irritable bowel syndrome. Thus, Na-methyl-N-[2-(3-tert-butyl-4-hydroxyphenyl)-1-methylethyl]-L-valinamide was condensed with Boc-Phe-OH using HOBt and DIC in DMF at room temperature for 2.5 days followed by deprotection with CF3CO2H in CH2Cl2 to give the title compound (II). II in vitro showed IC50 of 1.9 nM for inhibiting the binding of [125I]motilin motilin receptor preparation from rabbit ileum mucus membrane.


IT RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)
 (preparation of 3-tert-butyl-L-tyrosinamide-containing peptide compds. as motilin receptor antagonists, inhibitors of digestive tract motility,

L4 ANSWER 170 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)
 and remedy for high blood motilin)
 RN 220806-26-8 CAPLUS
 CH L-Tyrosinamide, L-phenylalanyl-(1S)-1,2,3,4-tetrahydro-3-isoguinolinecarbonyl-3-(1,1-dimethylethyl)-, mono(trifluoroacetate) (salt) (9CI) (CA INDEX NAME)

CM 1

CRN 220806-25-7
 CMF C32 H38 N4 O4

Absolute stereochemistry.



CM 2

CRN 76-05-1
 CMF C2 H F3 O2




REFERENCE COUNT: 6 THERE ARE 6 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 171 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
 ACCESSION NUMBER: 1999:137688 CAPLUS
 DOCUMENT NUMBER: 130:346848
 TITLE: Discovery of a Novel Class of Selective Non-Peptide Antagonists for the Human Neurokinin-3 Receptor. 2. Identification of (S)-N-(1-Phenylpropyl)-3-hydroxy-2-phenylquinoline-4-carboxamide (SB 223412)
 Giardina, Giuseppe A. M.; Raveglia, Luca F.; Grugni, Mario; Sarau, Henry M.; Farina, Carlo; Medhurst, Andrew D.; Graziani, Davide; Schmidt, Dulcie B.; Rigolio, Roberto; Luttmann, Mark; Cavignera, Stefano; Foley, James J.; Vecchiotti, Vittorio; Hay, Douglas W. P.
 CORPORATE SOURCE: Department of Medicinal Chemistry, SmithKline Beecham S.p.A., Baranzate, 20021, Italy
 SOURCE: Journal of Medicinal Chemistry (1999), 42(6), 1053-1065
 CODEN: JMCMAR; ISSN: 0022-2623
 PUBLISHER: American Chemical Society
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 AB Optimization of the previously reported 2-phenyl-4-quinolinecarboxamide hNK-3 receptor antagonist with regard to potential metabolic instability of the ester moiety and affinity and selectivity for the human neurokinin-3 (hNK-3) receptor, is described. The ester functionality could be successfully replaced by the ketone or by lower alkyl groups (Et or n-Pr). Investigation of the substitution pattern of the quinoline ring resulted in the identification of position 3 as a key position to enhance hNK-3 binding affinity and selectivity for the hNK-3 vs. the hNK-2 receptor. All of the chemical groups introduced at this position, with the exception of halogens, increased the hNK-3 binding affinity, and SB 223412 (3-OH derivative, hNK-3-CHO binding Ki = 1.4 nM) and the 3-NH2 derivative (hNK-3-CHO binding Ki = 1.2 nM) were the most potent compds. of this series. Selectivity studies vs. the other neurokinin receptors (hNK-2-CHO and hNK-1-CHO) revealed that SB 223412 is about 100-fold selective for the hNK-3 vs. hNK-2 receptor, with no affinity for the hNK-1 at concns. ≤ 100 μ M. In vitro studies demonstrated that SB 223412 is a potent functional antagonist of the hNK-3 receptor (reversal of senktide-induced contractions in rabbit isolated iris sphincter muscles and reversal of NKB-induced Ca2+ mobilization in CHO cells stably expressing the hNK-3 receptor), while in vivo this compound showed oral and i.v. activity in NK-3 receptor-driven models (senktide-induced behavioral responses in mice and senktide-induced miosis in rabbits). Overall, the biol. data indicate that SB 223412 [(S)-N-(1-phenylpropyl)-3-hydroxy-2-phenylquinoline-4-carboxamide] may serve as a pharmacol. tool in animal models of disease to assess the functional and pathophysiol. role of the NK-3 receptor and to establish therapeutic indications for non-peptide NK-3 receptor antagonists.

IT RL: BPR (Biological process); BSU (Biological study, unclassified); RCT (Reactant); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation); PROC (Process); RACT (Reactant or reagent)
 (discovery of a novel class of selective non-peptide antagonists for human neurokinin-3 receptor and identification of (S)-N-(1-phenylpropyl)-3-hydroxy-2-phenylquinoline-4-carboxamide (SB 223412))

RN 174635-51-9 CAPLUS
 CN Benzeneacetic acid, α -[[(2-phenyl-4-quinolinyl)carbonyl]amino]-, methyl ester (9CI) (CA INDEX NAME)


L4 ANSWER 171 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)



REFERENCE COUNT: 37 THERE ARE 37 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 172 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
 ACCESSION NUMBER: 1999:136877 CAPLUS
 DOCUMENT NUMBER: 130:209514
 TITLE: Preparation of hydroxyindanylbutanediarnides and related compounds as inhibitors of aggrecanase and matrix metalloproteinases for the treatment of arthritis.
 INVENTOR(S): Yao, Wenging; Decicco, Carl P.
 PATENT ASSIGNEE(S): Du Pont Pharmaceuticals Company, USA
 SOURCE: PCT Int. Appl., 251 pp.
 CODEN: PIXXD2
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 9909000	A2	19990225	WO 1998-US17048	19980818
WO 9909000	A3	19990910		
W:	AU, BR, CA, CN, CZ, DE, HU, IL, JP, KR, LT, LV, MX, NO, NZ, PL, RO, SG, SI, SK, UA, VN, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, PT, SE			
US 6576664	B1	20030610	US 1998-134484	19980814
ZA 9807376	A	20000217	ZA 1998-7376	19980817
CA 2301038	AA	19990225	CA 1998-2301038	19980818
AU 9890214	A1	19990308	AU 1998-90214	19980818
EP 1005448	A2	20000607	EP 1998-942083	19980818
R:	AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, FI			
EE 200000093	A	20001215	EE 2000-200000093	19980818
BR 9815596	A	20010102	BR 1998-15596	19980818
MX 200000956	A	20001020	MX 2000-956	20000127
NO 2000000784	A	20000407	NO 2000-784	20000217
PRIORITY APPLN. INFO.:			US 1997-55944P	P 19970818
			US 1997-68335P	P 19971219
			WO 1998-US17048	W 19980818
OTHER SOURCE(S):		MARPAT 130:209514		
GI				




AB Title compds., e.g., [I; R2, R3, R5 = UXYZUaXaYaZa; U, Ua = null, O, CO,

L4 ANSWER 172 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)
 CO2, NRA, O2CNRA, CO2, S0p, etc.; X, Xa = null, H, alkylene, alkenylene, alkynylene; Y, Ya = null, H, O, NRA, CO, S0p; Z, Za = null, H, (substituted) carbocyclyl, heterocyclyl; Ra = H, alkyl, Ph, PhCH2; R8, R9 = H, (substituted) alkyl, alkenyl, alkaryl, carbocyclyl, heterocyclyl, etc.; R8R9C = atoms to form a heterocyclyl ring; p undefined, were prepd. as inhibitors of aggrecanase and matrix metalloproteinases (no data). Thus, (2R)-isobutyl-3-(tert-butoxycarbonyl)propanoic acid, (1S,2R)-cis-1-amino-2-indanol, TBTU, and (Me2CH)2NEt were stirred at 0° to room temp. to give 87% N1-(2R-hydroxy-1S-indanyl)-2R-isobutyl-3-(tert-butoxycarbonyl)propanamide. This in CH2Cl2/H2O was treated with CF3CO2H to give N-(2R-hydroxy-1S-indanyl)-2R-isobutyl-3-(hydroxycarbonyl)propanamide. The latter in DMF was treated with PhCH2ONH2.HCl, TBTU, and (Me2CH)2NEt at 0° to room temp. to give a product which was hydrogenolyzed in MeOH over Pd/BaSO4 to give N1-(2(R)-hydroxy-1(S)-indanyl)-N4-hydroxy-2(R)-isobutylbutanediamide.

IT 220682-81-5P
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USSS (Uses)
 (preparation of hydroxyindanylbutanediarnides and related compds. as inhibitors of aggrecanase and matrix metalloproteinases for the treatment of arthritis)

RN 220682-81-5 CAPLUS
 CN Butanediamide, N1-[(1S,2R)-2,3-dihydro-2-hydroxy-1H-inden-1-yl]-N4-hydroxy-2-[(3-hydroxyphenyl)methyl]-3-[(2-quinolinylcarbonyl)amino]-, (2R,3S)-(9CI) (CA INDEX NAME)

Absolute stereochemistry.



L4 ANSWER 173 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
 ACCESSION NUMBER: 1999:113708 CAPLUS
 DOCUMENT NUMBER: 130:153982
 TITLE: Preparation of N-sulfonyl phenylalanine dipeptide derivatives and analogs as inhibitors of leukocyte adhesion mediated by VLA-4
 INVENTOR(S): Dappen, Michael S.; Dressen, Darren B.; Grant, Francine S.; Pleiss, Michael A.; Robinson, Cynthia Y.; Sarantakis, Dimitrios; Thorsett, Eugene D.
 PATENT ASSIGNEE(S): Athena Neurosciences, Inc.; USA; American Home Products Corporation
 SOURCE: PCT Int. Appl., 190 pp.
 CODEN: PIXXD2
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 9906433	A1	19990211	WO 1998-US15952	19980731
W:	AL, AM, AT, AU, AZ, BA, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GR, GM, HR, HU, ID, IL, IS, JP, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, RW: GH, GM, KE, LS, MW, SD, SZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG			
AU 9886786	A1	19990222	AU 1998-86786	19980731
EP 1001973	A1	20000524	EP 1998-938207	19980731
R:	AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO			
BR 9811569	A	20000919	BR 1998-11569	19980731
JP 2001512136	T2	20010821	JP 2000-505188	19980731
US 6559127	B1	20030506	US 1998-127533	19980731
NO 2000000451	A	20000323	NO 2000-451	20000128
US 2003166575	A1	20030904	US 2002-266889	20021007
PRIORITY APPLN. INFO.:			US 1997-112010P	P 19970731
			US 1997-904416	A1 19970731
			US 1998-127533	A3 19980731
			WO 1998-US15952	W 19980731
OTHER SOURCE(S):		MARPAT 130:153982		


AB Disclosed are title compds. R1SO2NR2CHR3QCHR5COR6 [R1 = (un)substituted alkyl, (un)substituted aryl, (un)substituted cycloalkyl, (un)substituted heterocyclyl; R2 = H, any group R1; R1R2 may form (un)substituted heterocyclic ring; R3 = H, any group R1; R2R3 may form (un)substituted unsatd. heterocyclic ring; R5 = CH2X1; X1 = H, OH, optionally substituted acylamino, alkyl, aryloxy, aryl, aryloxyaryl, CO2H, carboxyalkyl, carboxyheteroaryl, etc.; Q = C(X)NR7; R7 = H, alkyl; X = O, S; R6 = NH2, (un)substituted alkoxy, (un)substituted cycloalkoxy, succinimidyloxy, adamantylamino, β-cholest-5-en-3-yloxy, NHOY, NH(CH2)pCO2Y, OCH2NR9R10; Y = H, (un)substituted alkyl, (un)substituted aryl; p = 1-8; R9 = (un)substituted CO-aryl; R10 = H, CH2CO2R11, NHSO2; R11 = alkyl; Z = (un)substituted alkyl, (un)substituted cycloalkyl, (un)substituted aryl, (un)substituted heteroaryl, (un)substituted heterocyclyl; and pharmaceutically acceptable salts thereof, with proviso(s) which bind VLA-4 (also referred to as integrin α4β1 and CD49/CD29). Certain of these compds. also inhibit leukocyte adhesion and, in particular, leukocyte adhesion mediated by VLA-4. Such compds. are useful in the treatment of inflammatory diseases in a mammalian patient, e.g., human.

L4 ANSWER 173 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)
 wherein the disease may be, for example, asthma, Alzheimer's disease, atherosclerosis, AIDS dementia, diabetes, inflammatory bowel disease, rheumatoid arthritis, tissue transplantation, tumor metastasis and myocardial ischemia. The compds. can also be administered for the treatment of inflammatory brain diseases such as multiple sclerosis. Thus, reaction of Ts-Gly-OH (Ts = tosyl) with oxalyl chloride in CH₂Cl₂, followed by peptide coupling with L-phenylalanine benzyl ester tosylate and catalytic hydrogenolysis, gave desired title compd. Ts-Gly-Phe-OH. All prepd. compds. have IC₅₀ ≤ 15 μM in a VLA-4 binding assay.

IT 220185-85-3P
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)
 (preparation of N-sulfonyl phenylalanine dipeptide derivs. and analogs as inhibitors of leukocyte adhesion mediated by VLA-4)

RN 220185-85-3 CAPLUS
 CN L-Phenylalanine, N-[(3S)-1,2,3,4-tetrahydro-2-[(4-methylphenyl)sulfonyl]-3-isoquinolinyl]carbonyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



REFERENCE COUNT: 7 THERE ARE 7 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 174 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
 ACCESSION NUMBER: 1999:104519 CAPLUS
 DOCUMENT NUMBER: 130:153971
 TITLE: Preparation of tryptophan ureas as neurokinin antagonists
 INVENTOR(S): Shah, Shrenik K.; Qi, Hongbo; Maccoss, Malcolm
 PATENT ASSIGNEE(S): Merck and Co., Inc., USA
 SOURCE: U.S., 14 pp.
 CODEN: USXXAM
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 5869489	A	19990209	US 1997-814387	19970311
PRIORITY APPLN. INFO.:			US 1997-814387	19970311
OTHER SOURCE(S):		MARPAT 130:153971		

GI

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *


AB Disclosed are substituted azacycles I [ring G = spirocycle Q1 or Q2, piperazine Q3, piperidine Q4; X = CH₂, NSO₂Me, NAc; R = Ph, 2-MeOC₆H₄, 2-MeC₆H₄, CH₂Ph; R1 = Ph, R11 = NOME [sic] (NHAc intended); R1 = H, R11 = CH₂Ph, 1,2,3,4-tetrahydroquinazolin-2-on-1-yl; R2 = OCH₂Ph wherein the Ph is optionally substituted with 1-3 substituents halo, Me, or CF₃; N(R3)-C1-4 alkylphenyl, wherein the C1-4 alkyl may be linear or branched, the Ph is optionally substituted with 1-3 substituents halo, Me, MeO, or CF₃; R3 = H, Me, Et] as tachykinin receptor antagonists useful in the treatment of inflammatory diseases, pain or migraine, and asthma. In particular compds. I are neurokinin antagonists. Thus, amidation of 1.967 g Boc-Trp-OH (Boc = Me₃CO₂C) with 0.87 mL MeNHCH₂Ph gave 2.56 g of the corresponding amide, which underwent deprotection with CF₃CO₂H, condensation with carbonyldiimidazole, and urea formation with spiro[1H-indene-1,4'-piperidine] hydrochloride to give title compound II (L-743,516). II and related Trp derive. showed IC₅₀ values of >1000 to 1 nM for human neurokinin 1 (NK1) antagonist activity.

IT 199110-44-6P
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)
 (preparation of tryptophan ureas as neurokinin antagonists)

RN 199110-44-6 CAPLUS
 CN Spiro[3H-indole-3,4'-piperidine]-1'-carboxamide, 1,2-dihydro-N-[(1S)-1-[(methyl(phenylmethyl)amino)carbonyl]-3-phenylpropyl]-1-(methylsulfonyl)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

L4 ANSWER 174 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)



REFERENCE COUNT: 2 THERE ARE 2 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 175 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
 ACCESSION NUMBER: 1999:96518 CAPLUS
 DOCUMENT NUMBER: 130:153978
 TITLE: Solid-phase synthesis of peptide CGRP-antagonists for use as medicaments
 INVENTOR(S): Beck-Sicking, Annette; Rist, Beate; Entzeroth, Michael
 PATENT ASSIGNEE(S): Dr. Karl Thomae G.m.b.H., Germany
 SOURCE: Ger. Offen., 20 pp.
 CODEN: GWXXBX
 DOCUMENT TYPE: Patent
 LANGUAGE: German
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
DE 19732944	A1	19990204	DE 1997-19732944	19970731
PRIORITY APPLN. INFO.:			DE 1997-19732944	19970731
OTHER SOURCE(S):		MARPAT 130:153978		


AB Variations on the r-CGRP-alpha-27-37 partial sequence H-F27-V28-P29-T30-N31-V32-G33-S34-E35-A36-F37-NH₂ (see text for specifications) were prepared using solid-phase peptide synthesis techniques, for use in acute and prophylactic treatment of headache, non-insulin-dependent diabetes mellitus, cardiovascular disease, skin disease, inflammatory disease, allergic rhinitis, asthma, clotting disorders, and morphine tolerance (no data).


IT 220198-73-2P
 RL: SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)
 (preparation of via solid-phase synthesis as CGRP-antagonists for use as medicaments)

RN 220198-73-2 CAPLUS
 CN L-Phenylalaninamide, L-phenylalanyl-L-valyl-L-prolyl-L-threonyl-L-α-aspartyl-L-valylglycyl-1,2,3,4-tetrahydro-3-isoquinolinecarbonyl-L-phenylalanyl-L-alanyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

PAGE 1-A





L4 ANSWER 176 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
 ACCESSION NUMBER: 1999:89647 CAPLUS
 DOCUMENT NUMBER: 130:196946
 TITLE: Conformational preferences of peptides containing reverse-turn mimetic bicyclic lactams. Inverse γ -turns versus type-II' β -turns. Insights into β -hairpin stability
 AUTHOR(S): Belvisi, Laura; Gennari, Cesare; Mielgo, Antonia; Potenza, Donatella; Scolastico, Carlo
 CORPORATE SOURCE: Dipartimento Chimica Organica Industriale, Univ. Studi Milano, Milan, Italy
 SOURCE: European Journal of Organic Chemistry (1999), (2), 389-400
 CODEN: EJOCHF; ISSN: 1434-193X
 PUBLISHER: Wiley-VCH Verlag GmbH
 DOCUMENT TYPE: Journal
 LANGUAGE: English

AB The conformational preferences of constrained peptides containing reverse-turn mimetic bicyclic lactams were investigated by NMR and IR. The exptl. results were complemented by computer modeling studies and show that the constrained peptides form an inverse γ -turn or a type-II' β -turn through intramol. H-bonding, depending on the nature of the reverse-turn mimic. In N-acetylated tetrapeptide mimics incorporating the two different bicyclic lactams, H(5) is available for either a γ -turn (7-membered ring with the CO group of the bicyclic lactam) or a β -turn (10-membered ring with the CO group of residue 2). Peptides incorporating a (5,7)-bicyclic lactam predominantly induce the γ -turn conformation, while those incorporating a (5,6)-bicyclic lactam can promote either a γ -turn or a β -turn conformation, with the β -turn usually being preferred and with varying degrees of β -hairpin formation.


IT 220563-60-0

RL: PRP (Properties)
 (conformational anal.)

RN 220563-60-0 CAPLUS


CN L-Phenylalanine, N-acetylglycyl-(3S,6S,9aS)-6-aminooctahydro-5-oxo-1H-pyrrolo[1,2-a]azepine-3-carbonyl-, methyl ester (9CI) (CA INDEX NAME)

Absolute stereochemistry. Rotation (-).



REFERENCE COUNT: 68 THERE ARE 68 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 177 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
 ACCESSION NUMBER: 1999:89646 CAPLUS
 DOCUMENT NUMBER: 130:196946
 TITLE: Solid-phase synthesis of peptides containing reverse-turn mimetic bicyclic lactams
 AUTHOR(S): Gennari, Cesare; Mielgo, Antonia; Potenza, Donatella; Scolastico, Carlo; Piarulli, Umberto; Manzoni, Leonardo
 CORPORATE SOURCE: Dipartimento Chimica Organica Industriale, Univ. Studi Milano, Milan, Italy
 SOURCE: European Journal of Organic Chemistry (1999), (2), 379-388
 CODEN: EJOCHF; ISSN: 1434-193X
 PUBLISHER: Wiley-VCH Verlag GmbH
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 GI



AB The solid-phase synthesis and characterization of a series of peptides containing reverse-turn mimetic bicyclic lactams I ($n = 3$, $R = H$; $n = 2$, $R = PhCH_2$) is reported. The bicyclic lactams possess high structural similarity to the 2 central residues of a β -turn. Amino acid conjugates of these bicyclic lactams were synthesized on solid supports following a 9-fluorenylmethoxycarbonyl (Fmoc) protection strategy on Wang-Merrifield resin. Coupling between amino acids was accomplished by diisopropylcarbodiimide (DIC)/hydroxyazabenzotriazole (HOAt). Coupling between amino acids and the mimics was performed with the potent Carpino's reagent, O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate (HATU). The final compds. were cleaved from the resin and obtained as N-acetylated Me esters or benzyl amides.


IT 220563-60-0P

RL: SPN (Synthetic preparation); PREP (Preparation)
 (solid-phase synthesis of peptides containing reverse-turn mimetic bicyclic lactams)

RN 220563-60-0 CAPLUS

CN L-Phenylalanine, N-acetylglycyl-(3S,6S,9aS)-6-aminooctahydro-5-oxo-1H-pyrrolo[1,2-a]azepine-3-carbonyl-, methyl ester (9CI) (CA INDEX NAME)


Absolute stereochemistry. Rotation (-).



L4 ANSWER 177 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)


REFERENCE COUNT: 90 THERE ARE 90 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 178 OF 261 CAPLUS COPYRIGHT 2004 ACS ON STN
 ACCESSION NUMBER: 1999:84975 CAPLUS
 DOCUMENT NUMBER: 130:237857
 TITLE: Bicyclic Tripeptide Mimetics with Reverse Turn Inducing Properties
 AUTHOR(S): Johannesson, Petra; Lindeberg, Gunnar; Tong, Weimin; Gogoll, Adolf; Karlen, Anders; Hallberg, Anders
 CORPORATE SOURCE: Department of Organic Pharmaceutical Chemistry, Uppsala University, Uppsala, SE-751 23, Swed.
 SOURCE: Journal of Medicinal Chemistry (1999), 42(4), 601-608
 CODEN: JMCMAJ; ISSN: 0022-2623
 PUBLISHER: American Chemical Society
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 GI




L4 ANSWER 178 OF 261 CAPLUS COPYRIGHT 2004 ACS ON STN (Continued)

PAGE 1-A



PAGE 1-B



REFERENCE COUNT: 14 THERE ARE 14 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

AB Analogs of the hypertensive octapeptide angiotensin II, comprising novel constrained 5,8-bicyclic and 5,9-bicyclic tripeptide units I and II (R = amino acid side chain) adopting nonclassical β -turn geometries, as deduced from their conformational anal., have been synthesized. Spontaneous bicyclization upon acid-catalyzed deprotection of a model peptide, encompassing a protected α -formyl-L-amino acid in position 5 and Cys residues in positions 3 and 7, revealed a strong preference for bicyclization toward the C-terminus. The bicyclic thiazolidine related angiotensin II analogs synthesized exhibited no affinity for the angiotensin II AT1 receptor.

IT 221235-39-8P
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); PRP (Properties); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation)
 (preparation of bicyclic tripeptide mimetics with reverse turn inducing properties)

RN 221235-39-8 CAPLUS
 CN L-Phenylalanine, L- α -aspartyl-L-arginyl-L-valyl-L-tyrosyl-(α S)- α -[(3S,5R)-3-amino-5-mercapto-2-oxo-1-pyrrolidinyl]-1H-imidazole-4-propanoyl-L-homocysteiny-, cyclic (5+6)-thioether (9CI) (CA INDEX NAME)

Absolute stereochemistry.


L4 ANSWER 179 OF 261 CAPLUS COPYRIGHT 2004 ACS ON STN
 ACCESSION NUMBER: 1999:37202 CAPLUS
 DOCUMENT NUMBER: 130:231893
 TITLE: A uniform molecular model of δ opioid agonist and antagonist pharmacophore conformations
 AUTHOR(S): Brandt, Wolfgang
 CORPORATE SOURCE: Institute of Biochemistry, Martin-Luther-Universität Halle-Wittenberg, Halle, D-06099, Germany
 SOURCE: Journal of Computer-Aided Molecular Design (1998), 12(6), 615-621
 CODEN: JCADEQ; ISSN: 0920-654X
 PUBLISHER: Kluwer Academic Publishers
 DOCUMENT TYPE: Journal
 LANGUAGE: English

AB On the basis of a model of the pharmacophore conformations of agonist of the δ -opioid receptor the corresponding δ -antagonist conformations were determined by means of force field calcs. The results explain the unusual behavior of several cyclic β -casomorphin analogs on the mol. level. Thus, for instance, the model helps to understand why Tyr-c[D-Orn-2-Nal-D-Pro-Gly] is a mixed μ -agonist and δ -antagonist. Furthermore, the model is consistent with low energy conformations of other δ -antagonists such as Tyr-Tic-Phe, Tyr-Tic-Phe-Phe, naltrindole and BMTX. The occupation of a special spatial area by bulky groups close to the protonated N-terminus of opioid peptides is assumed to be highly critical for the switch from agonist to antagonist behavior.

IT 221333-35-3
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); PRP (Properties); BIOL (Biological study)
 (uniform mol. model of δ -opioid agonist and antagonist pharmacophore conformations)

RN 221333-35-3 CAPLUS
 CN Dermorphin, 2-[(1,2,3,4-tetrahydro-3-isoquinolinecarboxylic acid)- (CA INDEX NAME)

Absolute stereochemistry.



REFERENCE COUNT: 27 THERE ARE 27 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 180 OF 261 CAPLUS COPYRIGHT 2004 ACS ON STN
 ACCESSION NUMBER: 1999:4410 CAPLUS
 DOCUMENT NUMBER: 130:139641
 TITLE: Design, synthesis, structure and properties of an α -helix cap template derived from N-[(2S)-2-chloropropionyl]-[(2S)-Pro-(2R)-Ala-(2S,4S)-4-thioPro-OMe] which initiates α -helical structures
 AUTHOR(S): Gani, David; Lewis, Arwel; Rutherford, Trevor; Wilkie, John; Stirling, Iain; Jenn, Thierry; Ryan, Martin D.
 CORPORATE SOURCE: School of Chemistry and Centre for Biomolecular Science, The University, St. Andrews, Fife, KY16 9ST, UK
 SOURCE: Tetrahedron (1998), 54(52), 15793-15819
 CODEN: TETRAH; ISSN: 0040-4020
 PUBLISHER: Elsevier Science Ltd.
 DOCUMENT TYPE: Journal
 LANGUAGE: English


AB A strategy based upon removing the requirement for all of the carbonyl dipoles to align at the same time in the transition state leading to the cyclization of N-[(2S)-2-chloropropionyl]-[(2S)-Pro-(2R)-Ala-(2S,4S)-4-thioPro-OMe] to a Zimm-Brugg type α -helix peptide initiator template was successful. Each amide bond of the 12-membered macrocyclic template existed in the trans-rotameric form. Derivs. of the template were prepared by extending the C-terminus and these were characterized by NMR spectroscopy and restrained simulated annealing. In deuteriochloroform solution at low temperature, sep. sets of NMR signals were observed for two rapidly interconverting helical conformational isomers of the thioether macrocycle which possessed an appended trialkylammonium ion. A similar time-averaged conformation was also observed in aqueous solution. At -80° in d2-dichloromethane the rate of conformational exchange was slowed sufficiently to obtain resonance assignments and NOE data sep. for each isomer. In the minor isomer (40%), the four carbonyl oxygen hydrogen-bond acceptors of the template are aligned in an α -helical conformation and in the major conformer the Pro2 carbonyl dipole was anti-aligned with the other three dipoles. Thus, the conformers differ in the orientation of one carbonyl group. Mol. modeling calcs. showed that the minor isomer was stabilized by coulombic interactions between the trialkylammonium salt and the carbonyl group dipole moments.

IT 220061-81-4P
 RL: PRP (Properties); SPN (Synthetic preparation); PREP (Preparation)
 (design, synthesis, structure and properties of α -helix cap template)

RN 220061-81-4 CAPLUS
 CN L-Phenylalaninamide, 1-[(2S)-2-mercapto-1-oxopropyl]-L-prolyl-D-alanyl-(4S)-4-mercapto-L-prolyl-N-methyl-, cyclic (1+3)-thioether (9CI) (CA INDEX NAME)

Absolute stereochemistry.

L4 ANSWER 180 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)



REFERENCE COUNT: 83 THERE ARE 83 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 181 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 1998:799995 CAPLUS
 DOCUMENT NUMBER: 130:52736
 TITLE: Preparation of biarylalkanoic acids as cell adhesion inhibitors
 INVENTOR(S): Durette, Philippe L.; Hagmann, William K.; Maccoss, Malcolm; Mills, Sander G.; Mumford, Richard A.
 PATENT ASSIGNEE(S): Merck & Co., Inc., USA
 SOURCE: PCT Int. Appl., 96 pp.
 CODEN: PIXXD2
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 9853817	A1	19981203	WO 1998-US10951	19980529
W:	AL, AM, AU, AZ, BA, BB, BG, BR, BY, CA, CN, CU, CZ, EE, GE, GW, HU, ID, IL, IS, JP, KG, KR, KZ, LC, LK, LR, LT, LV, MD, MG, MK, MN, MX, NO, NZ, PL, RO, RU, SG, SI, SK, SL, TJ, TM, TR, TT, UA, US, UZ, VN, YU, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM			
RW:	GH, GM, KE, LS, MW, SD, SZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG			
AU 9877031	A1	19981230	AU 1998-77031	19980529
AU 726585	B2	20001109		
EP 1017382	A1	20000712	EP 1998-924988	19980529
R:	AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, PT, IE, FI			
JP 2001517245	T2	20011002	JP 1999-500938	19980529
US 6291511	B1	20010918	US 1999-359015	19990722
PRIORITY APPLN. INFO.:			US 1997-47856P	P 19970529
			GB 1997-14316	A 19970707
			US 1997-66831P	P 19971125
			GB 1998-680	A 19980114
			US 1998-85793	B1 19980528
			WO 1998-US10951	W 19980529

OTHER SOURCE(S): MARPAT 130:52736
 AB Comps. R1YNR2CR3R4CONR5CR6R7X [R1 = (un)substituted alkyl, alkenyl, alkynyl, a cyclic group Cy, Cy-alkyl, Cy-alkenyl, Cy-alkynyl; R2, R3 independently are H or R1; or R2 and R3 together form a ring; R4, R7 independently are H, (un)substituted alkyl, alkenyl, alkynyl, aryl, arylalkyl, heteroaryl, or heteroarylalkyl; or R3 and R4 together form a ring; R5 = H or (un)substituted alkyl or Cy; R6 = diarylalkyl, -alkenyl, or -alkynyl; X = CO₂H, PO₃H₂, PH(O)OH, SO₂H, SO₃H or ester derivs., carbamoyl group, or 5-tetrazolyl; Y = CO, OCO, NHCO or iminocarbonyl group, SO₂, P(O)(OR₁) (R₁ = alkyl, alkenyl, alkynyl, aryl), COCO were prepared as cell adhesion inhibitors. Pharmaceutical compns. are described. Thus, N-(3,5-dichlorobenzenesulfonyl)-L-prolyl-L-4-(4-fluorophenyl)phenylalanine was prepared by coupling of N-(3,5-dichlorobenzenesulfonyl)-L-proline with 4-iodo-L-phenylalanine and reaction with 4-fluorobenzeneboronic acid.
 IT 217325-07-0P
 RL: SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)
 (Preparation of peptidyl biarylalkanoic acids as cell adhesion inhibitors)
 RN 217325-07-0 CAPLUS

L4 ANSWER 181 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)

CN [1,1'-Biphenyl]-4-propanoic acid, α-[[[[(3S)-2-[[[3,4-dimethoxyphenyl)sulfonyl]-1,2,3,4-tetrahydro-3-isoquinolinyl]carbonyl]amino]-, (αS)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




REFERENCE COUNT: 1 THERE ARE 1 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 182 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 1998:799992 CAPLUS
 DOCUMENT NUMBER: 130:52724
 TITLE: Preparation of heterocyclic dipeptide derivatives as cell adhesion inhibitors
 INVENTOR(S): Durette, Philippe L.; Hagmann, William K.; Maccoss, Malcolm; Mills, Sander G.; Mumford, Richard A.; Van Riper, Gail M.; Schmidt, Jack A.; Kevin, Nancy J.
 PATENT ASSIGNEE(S): Merck & Co., Inc., USA
 SOURCE: PCT Int. Appl., 129 pp.
 CODEN: PIXXD2
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 9853814	A1	19981203	WO 1998-US10940	19980529
W:	CA, JP, US			
RW:	AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE			
EP 1001764	A1	20000524	EP 1998-926122	19980529
R:	AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, PT, IE, FI			
JP 2002512625	T2	20020423	JP 1999-500934	19980529
WO 9964395	A1	19991216	WO 1998-US11623	19980611
W:	AL, AM, AU, AZ, BA, BB, BG, BR, BY, CN, CU, CZ, EE, GE, GW, HU, ID, IL, IS, KG, KR, KZ, LC, LK, LR, LT, LV, MD, MG, MK, MN, MX, NO, NZ, PL, RO, RU, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UZ, VN, YU, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM			
RW:	GH, GM, KE, LS, MW, SD, SZ, UG, ZW, BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG			
AU 9880595	A1	19991230	AU 1998-80595	19980611
PRIORITY APPLN. INFO.:			US 1997-48017P	P 19970529
			GB 1997-14314	A 19970707
			US 1997-66525P	P 19971125
			GB 1998-686	A 19980114
			WO 1998-US10940	W 19980529
			WO 1998-US11623	A 19980611

OTHER SOURCE(S): MARPAT 130:52724
 GI




AB Title compds. I [R1 = (un)substituted C1-10 alkyl, C2-10 alkenyl, C2-10 alkynyl, Cy, Cy-C1-10 alkyl, Cy-C2-10 alkenyl, Cy-C2-10 alkynyl; R2, R5 = independently (un)substituted H, C1-10 alkyl, C2-10 alkenyl, C2-10 alkynyl, aryl, aryl-C1-10 alkyl, heteroaryl, heteroaryl-C1-10 alkyl; R3 = H, (un)substituted C1-10 alkyl, Cy, Cy-C1-10 alkyl; R4 = H, any group R1; R3R4 form mono- or bicyclic ring containing 0-2 heteroatoms N, O, S; R4R5 form 3-7 membered mono- or bicyclic ring containing 0-2 heteroatoms N, O, S; R10, R11 = independently = any group R3, (un)substituted C2-10 alkenyl, C2-10

L4 ANSWER 182 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)
alkynyl; R10R11 may form 5-7 membered heterocyclic ring contg. 0-2 addnl. heteroatoms N, O, S; R6-R8 = independently any group R10, OR10, NO2, halo, S(O)mR10, SR10, SO3R10, NR10R11, COR10, CO2R10, O2R10, CN, CONR10R11, CF3, oxo, NR10S(O)mR11, etc.; two of R6-R8 may form 5-7 membered (un)satd. monocyclic ring contg. 0-3 heteroatoms N, O, S; Cy = cycloalkyl, heterocyclyl, aryl, heteroaryl; A, Z = independently C, C-C; B = bond, C, C-C, N, O, S, S(O)m; X = CO2R10, P(O)(OR10)(OR11), P(O)(R10)(OR11), S(O)mOR10, CONR10R11, 5-tetrazolyl; Y = CO, O2C, NR11CO, SO2, P(O)(OR4), COCO, m = 1-2) * are antagonists of VLA-4 and/or α4β7, and are useful for inhibition or prevention of cell adhesion and cell adhesion mediated pathologies. These compds. may be formulated into pharmaceutical compns. and are suitable for use in the treatment of asthma, allergies, inflammation, multiple sclerosis, and other inflammatory and autoimmune disorders. Thus, coupling of L-2-naphthylalanine tert-Bu ester (H-Nal-OTBu) (prepn. given) with Cbz-Pro-OH (Cbz = PhCH2O2C), followed by catalytic deprotection, sulfonation with 3,5-Cl2C6H3SO2Cl, and acidic deesterification gave desired N-sulfonyldipeptide Cl2C6H3SO2-Nal-Pro-OH. Procedures for inhibition of VLA-4 dependent adhesion to a CS-1 conjugate and VCAM-1G fusion protein are given.

IT 217450-09-4P
RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)
(preparation of heterocyclic dipeptide derivs. as cell adhesion inhibitors)
RN 217450-09-4 CAPLUS
CN L-Phenylalanine, N-[[[(3S)-2-[(3,4-dimethoxyphenyl)sulfonyl]-1,2,3,4-tetrahydro-3-isoquinolyl]carbonyl]-β-phenyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



REFERENCE COUNT: 10 THERE ARE 10 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT


L4 ANSWER 183 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
ACCESSION NUMBER: 1998:756613 CAPLUS
DOCUMENT NUMBER: 130:133637
TITLE: Side Chain Methyl Substitution in the δ-Opioid Receptor Antagonist TIPP Has an Important Effect on the Activity Profile
AUTHOR(S): Tourwe, Dirk; Mannekens, Els; Diem, Trang Nguyen Thi; Verheyden, Patricia; Jaspers, Hendrika; Toth, Geza; Peter, Antal; Kertesz, Istvan; Toeroek, Gabriella; Chung, Nga N.; Schiller, Peter W.
CORPORATE SOURCE: Benheid Organische Chemie, Vrije Universiteit Brussel, Brussels, B-1050, Belg.
SOURCE: Journal of Medicinal Chemistry (1998), 41(26), 5167-5176
CODEN: JMCMAR; ISSN: 0022-2623
PUBLISHER: American Chemical Society
DOCUMENT TYPE: Journal
LANGUAGE: English

AB The δ-opioid antagonist H-Tyr-Tic-Phe-Phe-OH (TIPP-OH) or its C-terminal amide analog was systematically modified topol. by substitution of each amino acid residue by all stereoisomers of the corresponding β-Me amino acid. The potency and selectivity (δ- vs μ- and κ-opioid receptor) were evaluated by radioreceptor binding assays. Agonist or antagonist potency were assayed in the mouse as deferenas and in the guinea pig ileum. In the TIPP analogs containing L-β-Me amino acids the influence on δ-receptor affinity and on δ-antagonist potency is limited, the [(2S,3R)-β-MePhe3]TIPP-OH analog being among the most potent δ-antagonists reported. In the D-β-Me amino acid series, the [D-β-MeTic2] analogs are δ-selective antagonists whereas [D-Tic2]TIPP-NH2 is a δ-agonist. NMR studies did not indicate any influence of the β-Me substituent on the conformation of the Tic residue. The [(2R,3S)-β-MePhe3]TIPP-NH2 is a potent δ-agonist, its C-terminal carboxylic acid analog being more δ-selective but displaying partial agonism in both the δ- and μ-biosassay. These results constitute further examples of a profound influence of β-Me substitution on the potency, selectivity, and signal transduction properties of a peptide.

IT 174147-54-7P
RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation)
(side chain Me substitution in the δ-opioid receptor antagonist TIPP has an important effect on the activity profile)
RN 174147-54-7 CAPLUS
CN L-Phenylalaninamide, L-tyrosyl-(3S)-1,2,3,4-tetrahydro-3-isoquinolinecarboxyl-(βR)-β-methyl-L-phenylalanyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

L4 ANSWER 183 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)




REFERENCE COUNT: 60 THERE ARE 60 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 184 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
ACCESSION NUMBER: 1998:745086 CAPLUS
DOCUMENT NUMBER: 130:4091
TITLE: Preparation of backbone-cyclized peptide derivatives as serine protease and thrombin inhibitors
INVENTOR(S): Adang, Anton Egbert Peter
PATENT ASSIGNEE(S): Akzo Nobel N.V., Neth.
SOURCE: PCT Int. Appl., 52 pp.
CODEN: PIXX2
DOCUMENT TYPE: Patent
LANGUAGE: English
FAMILY ACC. NUM. COUNT: 1
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 9850420	A1	19981112	WO 1998-EP2587	19980428
W: AM, AU, BB, BG, BR, CA, CN, CZ, EE, GE, HU, ID, IS, JP, KG, KP, KR, LK, LR, LT, LV, MD, MG, MN, MX, NO, NZ, PL, RO, RU, SG, SI, SK, TR, TT, UA, US, UZ, VN, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, RW: GH, GM, KE, LS, MW, SD, SZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, GN, TD, TG				
AU 9876520	A1	19981127	AU 1998-76520	19980428
AU 729910	B2	20010215		
EP 979240	A1	20000216	EP 1998-924265	19980428
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, FI				
BR 9809342	A	20000704	BR 1998-9342	19980428
NZ 500620	A	20001027	NZ 1998-500620	19980428
JP 2001524117	T2	20011127	JP 1998-547715	19980428
RU 2183642	C2	20020620	RU 1999-125967	19980428
ZA 9803629	A	19981104	ZA 1998-3629	19980429
US 6534495	B1	20030318	US 1999-403856	19991026
NO 9905316	A	19991101	NO 1999-5316	19991101
MX 9910057	A	20000731	MX 1999-10057	19991101
PRIORITY APPLN. INFO.:			EP 1997-201286	A 19970502
			WO 1998-EP2587	W 19980428

OTHER SOURCE(S): MARPAT 130:4091
GI




AB The invention relates peptide derivative R1S02-B-X-Z-C-O-Y [B = bond, amino acid NHC(H)(CH₂pCOOH)CO or ester derivative thereof, Gly, D-1-perhydroisoguanlinecarboxylic acid (D-1-Pig), D-3-Pig, D-1,2,3,4-tetrahydroisoquinoline-1-carboxylic acid (D-1-Tiq), D-3-Tiq, D-aminoetralinocarboxylic acid, aminoindeanocarboxylic acid, L- or D-amino acid containing hydrophobic, basic, or neutral side chain; X = amino acid containing hydrophobic side chain, Gln, Ser, Thr, 2-aminoisobutyric acid, NR2CH2CO, O, Cyclic amino acid options: ornithine, lysine, histocatom N, O or S, (un)substituted with C1-6 alkyl, C1-6 alkoxy, PhCH2O, XO; Z = Lys, 4-aminocyclohexylglycine; Y = (un)substituted NHCl-6 alkylene-Ph, OR4, NR5R6 = W, CH, R1 = R2O2C(CHR2)m, R2NH(CHR2)n, (un)substituted C1-12 alkyl, C2-12 alkenyl, C6-14 aryl, C7-15 aralkyl, C8-16 aralkenyl, C12H-R2 - independently H, C1-12 alkyl, C3-8 cycloalkyl, (un)substituted C6-C8 aryl or C1-15 H, C2-6 alkyl, C2-6 alkoxy, C2-6 alkenyl substituted with OH, C1-6 alkoxy, CO2H, CO2, C1-6 alkyl, COMH2, halo; R4 = H, C2-6 alkyl, CH2Ph; R5, R6 = independently H, C1-6 alkyl, (un)substituted C1-6 alkyl; RS6 = CH2CH2VCH2CH2; V = O, S, SO2; m = 1-3; n = 2-4; p = 1-3]. The compds. of the invention have anticoagulant activity and can be used in treating or preventing thrombin-related diseases. Thus, coupling of the compounds prepared according to the invention with Lys(Boc)-OH, NaOMe, and benzylamine) with backbone-cyclized dipeptide derivative II (prepared in 4 steps from L-α-amino-ε-caprolactam, Me bromoacetate, and benzyleulfonyl chloride), followed by oxidation and deprotection gave desired title compound III. III inhibited factor Xa with IC50 = 0.64 μM.

IT 215791-78-99
AB: BAC (Biological activity or effector except aureliin); BSU (Biological study, unclassified); SPN (Synthetic Preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES; Urease).

141 ANSWER 185 OF 261 CAPLUS: COPYRIGHT 2004 ACS ON STN
 ACCESSION NUMBER: 1998:720328 CAPLUS
 DOCUMENT NUMBER: 130:62593
 TITLE: Equilibrium of the cis-trans isomerization of the
 peptide bond with N-alkyl amino acids measured by 2D
 NMR
 AUTHOR(S): Misicka, Aleksandra; Verheyden, Patricia M. F.; Van
 Binst, Georges
 CORPORATE SOURCE: Department of Chemistry, Warsaw University, Warsaw,
 PL-02-093, Pol.
 SOURCE: Letters in Peptide Science (1998), 5(5-6), 375-377
 CODEN: LPSCEM; ISSN: 0929-5666
 PUBLISHER: Kluwer Academic Publishers
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 AB The conformational cis-trans equilibrium around the peptide bond in model
 tripeptides has been determined by 2D NMR methods (HMQC, ROESY). The study
 was limited to three different N0-substituted amino acids in position 2,
 namely Pro (proline), Tic (1,2,3,4-tetrahydroisoquinoline-3-carboxylic
 acid), and N-Mephe (N-methylphenylalanine). In all cases the amino acid
 in position 1 was tyrosine and in position 3, phenylalanine. The results
 of our studies show that the cis-trans ratio depends mostly on the
 configuration of the amino acids forming the peptide bond undergoing the
 cis-trans isomerization. The amino acid following the sequence (in
 position 3) does not have much influence on the cis-trans isomerization,
 indicating that there is no interaction of the side chains between these
 amino acids. The model peptides with the L-Tyr-L-AA-(L- or D-Phe (where
 AA is N-substituted amino acid) chiralities give 80-100% more of the cis
 form in comparison to the corresponding peptides with the D-Tyr-L-AA-(L-
 or D-Phe chiralities. These results indicate that the incorporation of the
 N-substituted amino acids in small peptides with the same chirality as the
 precedent amino acid involved in the peptide bond undergoing the cis/trans
 isomerization moves the equilibrium to a significant amount of the cis form.
 IT 217958-85-5
 RL: PEP (Physical, engineering or chemical process); PRP (Properties);
 PROC (Process)
 (equilibrium of cis-trans isomerization of peptide bond with N-alkyl amino
 acids measured by 2D NMR)
 RN 217958-85-5 CAPLUS
 CN D-Phenylalanine, L-tyrosyl-(3S)-1,2,3,4-tetrahydro-3-isoquinolinecarbonyl-
 (9S) (CA INDEX NAME)


Absolute stereochemistry.



REFERENCE COUNT: 16 THERE ARE 16 CITED REFERENCES AVAILABLE FOR THIS
RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 184 OF 261 CAPLUS COPYRIGHT 2004 ACS ON STN (Continued)
(prepn. of backbone-cyclized peptide derivs. as serine protease
inhibitors)
RN 215791-78-9 CAPLUS
CN Heptanoic acid, 7-amino-3-[[[(2S)-2-[[[(3R)-2-(ethylsulfonyl)-1,2,3,4-
tetrahydro-3-isquinolinyl]carbonyl]amino]-1-oxo-3-phenylpropyl]amino]-2-
oxo-, 1-methyl-ethyl ester (9CI) (CA INDEX NAME)


Absolute stereochemistry.



REFERENCE COUNT: 2 THERE ARE 2 CITED REFERENCES AVAILABLE FOR THIS
RECORD. ALL CITATIONS AVAILABLE IN THE REFORMAT


LA ANSWER 186 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
ACCESSION NUMBER: 1998:702092 CAPLUS
DOCUMENT NUMBER: 130:61238
TITLE: The relationship between structure and activity among
opioind peptides
AUTHOR(S): Deschamps, Jeffrey R.; George, Clifford;
Flippen-Anderson, Judith L.
CORPORATE SOURCE: Laboratory for the Structure of Matter, Naval Research
Laboratory, Washington, DC, 20375, USA
SOURCE: Letters in Peptide Science (1998), 5(5-6), 337-340
CODEN: LPSCDM; ISSN: 0929-5666
PUBLISHER: Kluwer Academic Publishers
DOCUMENT TYPE: Journal
LANGUAGE: English
AB Since the discovery and isolation of the endogenous opioind peptides Leu-
and Met-enkephalin, structural studies have been focused on deducing the
bioactive conformation of the peptide ligands. There., linear peptides
can have many different backbone conformations, yet early x-ray studies on
enkephalin and its analogs showed only two different backbone
conformations: extended and single β -bend. More recent reports
include a third conformation for Leu-enkephalin and constrained opioind
peptides from two "new" classes (i.e. cyclic and "all-aromatic" peptides).
In this report the relationship between solid-state x-ray structure and
opioind peptide activity is examined. The N-terminal amine nitrogen and the
two aromatic rings have previously been identified as structural features
important to the biol. activity of opioind peptides. From x-ray studies we
find that the distance between the centroids of the aromatic rings, and
between the N-terminal amino nitrogen and the centroid of the
phenylalanine ring, vary over a large range. There is a discernible
relationship, however, between the separation of the two rings and their
orientation that correlates with activity.
IT 219757-05-6
RL: BAC (Biological activity or effector, except adverse); BSU (Biological
study, unclassified); PRP (Properties); BIOL (Biological study)
(relationship between structure and activity among opioind peptides)
RN 219757-05-6 CAPLUS
CN L-Phenylalanine, L-Tyrosyl-(3R)-1,2,3,4-tetrahydro-3-isquinolinecarbonyl-
(9CI) [CA INDEX NAME]

Absolute stereochemistry.



REFERENCE COUNT: 32 THERE ARE 32 CITED REFERENCES AVAILABLE FOR THIS
RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT


L4 ANSWER 187 OF 261 CAPLUS COPYRIGHT 2004 ACS ON STN
 ACCESSION NUMBER: 1998:715619 CAPLUS
 DOCUMENT NUMBER: 130:139618
 TITLE: Design, construction and properties of peptide N-terminal cap templates devised to initiate α -helices. Part 3. Caps derived from N-[(2S)-2-chloropropionyl]- (2S)-Pro-(2R)-Ala-(2S,4S)-4-thioPro-OMe
 AUTHOR(S): Lewis, Arwel; Rutherford, Trevor J.; Wilkie, John; Jenn, Thierry; Gani, David
 CORPORATE SOURCE: School of Chemistry and Centre for Biomolecular Sciences, The University St. Andrews, St. Andrews, Fife, KY16 9ST, UK
 SOURCE: Journal of the Chemical Society, Perkin Transactions 1: Organic and Bio-Organic Chemistry (1998), (22), 3795-3806
 CODEN: JCPRB4; ISSN: 0300-922X
 PUBLISHER: Royal Society of Chemistry
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 GI



AB The construction of a 12-membered macrocyclic template capable of entraining attached peptides in helical conformations from acyclic tripeptide precursors derived from (S)-ClCHMeCO-Pro-Pro-(2S,4S)-4-thioPro-OMe has been severely hampered by the problem of simultaneously aligning carboxamide dipoles in the transition state for cyclization. Previously, the authors provided a detailed conformational anal. of the system and tested two methods for forcing the acyclic precursor into the macrocyclic conformation required for helix initiation. First, the destabilization of unwanted conformations in the transition state for cyclization, second, the stabilization of the favored transition state structure through the introduction of a hydrogen-bonding interaction. Both strategies were unsuccessful. A third strategy based upon removing the requirement for all of the carbonyl dipoles to align in the transition state at the same time was also tested and the results are presented here. The relaxation of the highly restrained Cu-N bond torsion for Pro3 in the acyclic precursor, through its substitution for a D-Ala residue, effectively decouples the motion of the second carboxamide group from the Cu-N bond torsion and allows the second carboxamide group to rotate. This rotation allows a helical conformation to develop in the transition state to the macrocycle without the need to align all of the carboxamide dipoles and results in successful cyclization to give template structures of the all trans (ttt) form. Deriva. of the template were prepared by extending the C-terminus and these were characterized by NMR and restrained

L4 ANSWER 187 OF 261 CAPLUS COPYRIGHT 2004 ACS ON STN (Continued)
 simulated annealing. In CDCl3 soln. at low temp., sep. sets of NMR signals were obsd. for two rapidly interconverting helical conformational isomers of the thioether macrocycle I which possessed an appended trialkylammonium ion. The free energy of activation for the transition (AGc.dbldag.) was 48 kJ mol⁻¹. A similar time-averaged conformation was also obsd. in aq. soln. At -80° in dichloromethane the rate of conformational exchange was slowed sufficiently to obtain resonance assignments and NOE data sep. for each isomer. In the minor isomer (40%), the four carbonyl oxygen hydrogen-bond acceptors of the template are aligned in an α -helical conformation and in the major conformer the Pro2 carbonyl dipole was anti-aligned with the other three dipoles. Thus, the conformers differ in the orientation of one carbonyl group. Mol. modeling calcns. showed that the minor isomer was stabilized by coulombic interactions between the trialkylammonium salt and the carbonyl group dipole moments.
 IT 220061-81-4P
 RL: FRP (Properties); SPN (Synthetic preparation); PREP (Preparation) (design, construction, and properties of proline-containing cyclotetrapeptide N-terminal cap templates devised to initiate α -helices)
 RN 220061-81-4 CAPLUS
 CN L-Phenylalaninamide, 1-[(2S)-2-mercapto-1-oxopropyl]-L-prolyl-D-alanyl-(4S)-4-mercapto-L-prolyl-N-methyl-, cyclic (1-3)-thioether (9CI) (CA INDEX NAME)

Absolute stereochemistry.




REFERENCE COUNT: 21 THERE ARE 21 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 188 OF 261 CAPLUS COPYRIGHT 2004 ACS ON STN
 ACCESSION NUMBER: 1998:708845 CAPLUS
 DOCUMENT NUMBER: 129:116563
 TITLE: Preparation of peptide amides and deipeptide amides as hepatitis C NS3 protease inhibitors
 INVENTOR(S): Hart, Terance; Quibell, Martin
 PATENT ASSIGNEE(S): Peptide Therapeutics Ltd., UK
 SOURCE: PCT Int. Appl., 45 pp.
 CODEN: PIIXD2
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 9846630	A1	19981022	WO 1998-GB1126	19980416
W: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, GW, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, AM, AZ, BY, BG, KZ, MD, RU, TJ, TW				
RW: GH, GM, KE, LS, MW, SD, SZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG				
AU 9870635	A1	19981111	AU 1998-70635	19980416
EP 975662	A1	20000202	EP 1998-917395	19980416
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, FI				
JP 2001521516	T2	20011106	JP 1998-543648	19980416
PRIORITY APPLN. INFO.: GB 1997-7659 A 19970416				
WO 1998-GB1126 W 19980416				

GI




AB Disclosed is a specific pharmacophoric profile which represents the structure for inhibitors of hepatitis C NS3 protease. The results from mapping studies of the enzyme with deipeptide substrates I (Xaa1 = Glu, D-Glu, Gln, Val, 2-aminobutyl, Xaa2 = Nle, Glu, Val, Tyr; Xaa3 = Glu, Val, Ser, Nle, 3-pyridylalanyl, homophenylalanyl, Tyr; Xaa4 = Glu, Leu, Phe, Pro) allow the generation of a particular pharmacophoric binding profile. Peptide amides R2-Glu-Nle-Xaa5-Xaa6-NHR [Xaa5 = D-1,2,3,4-tetrahydroquinoline-3-carboxylic acid (D-Tic), homophenylalanine; Xaa6 = Leu, Phe, Glu; R = CH2CH2CHMe2, CH2CH2Ph, n-Pr] possessing this motif were shown to be inhibitors of hepatitis NS3 protease. The inhibitors have use in the treatment of hepatitis C.

IT 214910-67-5P
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); THU (Therapeutic use);


L4 ANSWER 188 OF 261 CAPLUS COPYRIGHT 2004 ACS ON STN (Continued)
 BIOL (Biological study); PREP (Preparation); USES (Uses)
 (prepn. of peptide amides as hepatitis C NS3 protease inhibitors)
 RN 214910-67-5 CAPLUS
 CN L-Phenylalaninamide, N-benzoyl-L- α -glutamyl-L-norleucyl-(3R)-1,2,3,4-tetrahydro-3-isoquinolinecarboxyl-N-(3-methylbutyl)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



REFERENCE COUNT: 5 THERE ARE 5 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT


L4 ANSWER 189 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
 ACCESSION NUMBER: 1998:680190 CAPLUS
 DOCUMENT NUMBER: 130:34337
 TITLE: On the role of copper and iron in DNA cleavage by ochratoxin A. Structure-activity relationships in metal binding and copper-mediated DNA cleavage
 AUTHOR(S): Ardu, Jason A.; Gillman, Ivan G.; Manderville, Richard A.
 CORPORATE SOURCE: Department of Chemistry, Wake Forest University, Winston-Salem, NC, 27109-7486, USA
 SOURCE: Canadian Journal of Chemistry (1998), 76(6), 907-918
 CODEN: CJCHAG; ISSN: 0008-4042
 PUBLISHER: National Research Council of Canada
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 GI



AB Ochratoxin A (OTA, I; X = Cl) is a fungal carcinogen that facilitates single-strand DNA cleavage and DNA adduction when metabolically activated. To determine if redox-active transition metals induce OTA-mediated DNA damage, we have examined the toxin's ability to bind Cu(II) and Fe(III) in aqueous media and facilitate DNA cleavage in their presence using agarose gel electrophoresis and supercoiled plasmid DNA. Using fluorescence spectroscopy, I was found to bind Cu(II) readily at physiol. pH, while acidic conditions (pH 2.6) were employed to study Fe(III) binding due to the formation of Fe-oxide ppts. at higher pH values. Structure-activity relationships employing synthetic derivs. of I implied that I binds both Cu(II) and Fe(III) by its phenolic oxygen, while the carboxylic acid of its phenylalanine moiety binds Cu(II), but does not appear to play a role in Fe(III) coordination at pH 2.6. In terms of metal-mediated DNA cleavage, no role for I could be detected in Fe-induced DNA strand


L4 ANSWER 189 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)
 scission. With Cu(II), DNA cleavage by the 1:1 copper-bound complex of I could only be initiated by addn. of a suitable reducing agent (sodium ascorbate). However, I was found to facilitate DNA cleavage by the Cu(II) complex of 1,10-phenanthroline (Cu(OP)2); a prototypical Cu-mediated nuclease system that cleaves DNA upon activation by an external reducing agent. Structure-activity relationships employing analogs lacking the chlorine atom, ochratoxin B, I: X = H, and the lactone (II), indicated that the chlorine atom is essential for activity of the OTA in potentiating DNA cleavage by Cu(OP)2. The implications of our findings to the genotoxic properties of I are discussed.
 IT 216967-81-6P
 RL: PRP (Properties); RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)
 RN 216967-81-6 CAPLUS
 CN L-Phenylalanine, N-[(3,4-dihydro-8-hydroxy-3-methyl-1-oxo-1H-2-benzopyran-7-yl)carbonyl]-, methyl ester (9CI) (CA INDEX NAME)

Absolute stereochemistry.



REFERENCE COUNT: 57 THERE ARE 57 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT


L4 ANSWER 190 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
 ACCESSION NUMBER: 1998:606905 CAPLUS
 DOCUMENT NUMBER: 129:290398
 TITLE: L-Tryptophan urea amides as NK1/NK2 dual antagonists
 AUTHOR(S): Qi, Hongbo; Shah, Shrenik K.; Cascieri, Margaret A.; Sadowski, Sharon J.; MacCoss, Malcolm
 CORPORATE SOURCE: Department of Medicinal Chemistry and Department of Molecular Pharmacology and Biochemistry, Rahway, NJ, 07065, USA
 SOURCE: Bioorganic & Medicinal Chemistry Letters (1998), 8(16), 2259-2262
 CODEN: BMCLE8; ISSN: 0960-894X
 PUBLISHER: Elsevier Science Ltd.
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 GI



AB The authors report that a systematic modification of an NK1 receptor selective antagonist resulted in the identification of novel compds. I (X = CH2, NSO2Me) with high affinity for both NK1 and NK2 receptors.
 IT 199110-44-6P
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation)
 (preparation and neurokinin receptor dual antagonist activity of substituted tryptophan amides)
 RN 199110-44-6 CAPLUS
 CN Spiro[3H-indole-3,4'-piperidine]-1'-carboxamide, 1,2-dihydro-N-[(1S)-1-[methyl(phenylmethyl)amino]carbonyl]-3-phenylpropyl]-1-(methylsulfonyl)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.


L4 ANSWER 190 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)



REFERENCE COUNT: 14 THERE ARE 14 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 191 OF 261 CAPLUS COPYRIGHT 2004 ACS ON STN
 ACCESSION NUMBER: 1998:597984 CAPLUS
 DOCUMENT NUMBER: 130:10306
 TITLE: The receptor-bound conformation of H-Tyr-Tic-(Phe-Phe)-OH related δ -opioid antagonists contains all trans peptide bonds
 AUTHOR(S): Wilkes, B. C.; Nguyen, T. M.-D.; Weltrowska, G.; Carpenter, K. A.; Lemieux, C.; Chung, N. N.; Schiller, P. W.
 CORPORATE SOURCE: Laboratory of Chemical Biology and Peptide Research, Clinical Research Institute of Montreal, Montreal, QC, H2W 1R7, Can.
 SOURCE: Peptides 1996, Proceedings of the European Peptide Symposium, 24th, Edinburgh, Sept. 8-13, 1996 (1998). Meeting Date 1996, 911-912. Editor(s): Ramage, Robert; Epton, Roger. Mayflower Scientific: Kingwinford, UK.
 DOCUMENT TYPE: CODEN: 66RCAS
 LANGUAGE: Conference
 AB A mol. mechanics study and energy minimization of two peptide δ -opioid antagonists containing a tetrahydroisoquinoline-3-carboxylic acid (Tic) residue revealed low energy conformers consistent with the previously proposed receptor-bound conformation containing all trans peptide bonds. No conformations with cis peptide bonds were found for either peptide. Assuming that all compts. of Tic-containing peptides have similar conformational requirements for δ -antagonism, their bioactive conformations must contain all trans peptide bonds.
 IT 207342-54-9
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); PRP (Properties); BIOL (Biological study) (receptor-bound conformation of H-Tyr-Tic-(Phe-Phe)-OH related δ -opioid antagonists contains all trans peptide bonds)
 RN 207342-54-9 CAPLUS
 CN L-Phenylalanine, (3S)-2-[(2S)-2-amino-3-(4-hydroxyphenyl)propyl]-1,2,3,4-tetrahydro-3-isoquinolinecarboxyl-L-phenylalanyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




REFERENCE COUNT: 5 THERE ARE 5 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 191 OF 261 CAPLUS COPYRIGHT 2004 ACS ON STN (Continued)

L4 ANSWER 192 OF 261 CAPLUS COPYRIGHT 2004 ACS ON STN
 ACCESSION NUMBER: 1998:543220 CAPLUS
 DOCUMENT NUMBER: 129:175563
 TITLE: 4-Substituted quinoline derivatives and 4-substituted quinoline combinatorial libraries
 INVENTOR(S): Hayes, Thomas K.; Forood, Behrouz; Kiely, John S.
 PATENT ASSIGNEE(S): Trega Biosciences, Inc., USA
 SOURCE: PCT Int. Appl., 124 pp.
 CODEN: PIXXD2
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 9834115	A1	19980806	WO 1997-US22391	19971205
W: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZW, AM, AZ, BY, BG, KZ, MD, RU, TJ, TW				
RW: GH, KE, LS, MW, SD, SZ, UG, ZW, AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG				
AU 9881919	A1	19980825	AU 1998-81919	19971205
EP 977989	A1	20000209	EP 1997-94975	19971205
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, FI				
US 6262269	B1	20010717	US 1998-17785	19980203
US 6388081	B1	20020514	US 1999-376670	19990816
PRIORITY APPLN. INFO.: US 1997-795392 A 19970204				
US 1997-126414 P 19970204				
WO 1997-US22391 W 19971205				
US 1998-17785 A3 19980203				

OTHER SOURCE(S): MARPAT 129:175563
 GI



AB The invention relates to novel 4-substituted quinoline derivs. I, their salts, and combinatorial libraries containing mixts. of two or more such compts. [wherein R1 = bond, (un)substituted alk(en/yn)ylene, cycloalk(en)ylene, phenylene, naphthylene, heterocycle, heteroaryl, amino, CH2CONH, (CH2)par(CH2)q, etc.; p, q = 0-6 but both cannot be 0; Ar =

L4 ANSWER 192 OF 261 CAPLUS COPYRIGHT 2004 ACS ON STN (Continued)
 (un)substituted Ph or heteroaryl; R2, R3, R4 = H, halo, (un)protected OH, cyano, NO2, (un)substituted alk(en/yn)yl, alkoxy, cycloalk(en)yl, heterocyclyl, phenylalkyl, Ph, naphthyl, etc.; R5 = H, (un)substituted alk(en/yn)yl, cycloalk(en)yl, Ph, naphthyl, phenylalkyl, (un)protected CO2H, acyl, heterocyclyl, etc.; R6 = H, (un)substituted Ph, naphthyl, 2-oxopyrrolidin-1-yl and higher homologs, (un)substituted NHCHO; R7 = H, (un)substituted alkyl; Y = CO2H, OH, SH, NHR8, CONHR8, CH2OH, CH2NH2, CH2NHR8; R8 = H, (un)substituted alkyl, or functionalized resin; R9 = H, (un)substituted alkyl, phenylalkyl, acyl, PhSO2, alkylsulfonyl, alkylaminocarbonyl, or PhNHCO, or is absent; dotted lines = optional pi bonds). The invention also relates to the generation of such libraries. In 12 examples, libraries of I ranging in size from 2380 to 39,440 compts. were prepd. as mixed sublibraries. Data for control compts. (samples of individually known intermediates and products, cleaved from simultaneously processed control resins) are given for some examples. Both quinoline and tetrahydroquinoline libraries were prepd. For instance, tea-bags of MBHA resin were each coupled with L- or D-N-BOC-p-nitrophenylalanine, the BOC groups were removed from both, and the amino groups were each acylated with 170 carboxylic acids. The acylated, resin-bound products were mixed and reduced at the nitro group, and the amine product mixts. were condensed with 58 different aldehydes and cyclized with 4-methoxystyrene. Cleavage of the resin-bound products with HF gave mixed sublibraries of I. Individual control samples of products, such as II (R5 = 1-naphthyl, 2,3-difluorophenyl, cyclohexyl, etc.), were obtained by reactions of pure, resin-bound L-N-propanoyl-p-aminophenylalanine control samples with individual aldehydes and 4-methoxystyrene. Potential applications of I (no data) may include use as antibacterials, NMDA antagonists, or analogues.
 IT 211376-13-5P
 RL: SPN (Synthetic preparation); PREP (Preparation) (resin-cleavage control intermediate; preparation of tricyclic tetrahydroquinoline derivs. and combinatorial libraries)
 RN 211376-13-5 CAPLUS
 CN 9H-Xanthene-9-carboxamide, N-[2-amino-1-[(4-nitrophenyl)methyl]-2-oxoethyl]- (9CI) (CA INDEX NAME)




REFERENCE COUNT: 4 THERE ARE 4 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 192 OF 261 CAPLUS COPYRIGHT 2004 ACS ON STN (Continued)

L4 ANSWER 193 OF 261 CAPLUS COPYRIGHT 2004 ACS ON STN
 ACCESSION NUMBER: 1998:543216 CAPLUS
 DOCUMENT NUMBER: 129:175562
 TITLE: Tricyclic tetrahydroquinoline derivatives and tricyclic tetrahydroquinoline combinatorial libraries
 INVENTOR(S): Hayes, Thomas K.; Kiely, John S.
 PATENT ASSIGNEE(S): Trega Biosciences, Inc., USA
 SOURCE: PCT Int. Appl., 119 pp.
 CODEN: PIXXD2
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 9834111	A1	19980806	WO 1997-US22206	19971205
W:	AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM			
RW:	GH, KE, LS, MW, SD, SZ, UG, ZW, AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG			
US 5925527	A	19990720	US 1997-795893	19970204
AU 985528	A1	19980825	AU 1998-55928	19971205
NZ 337046	A	20000128	NZ 1997-337046	19971205
EP 983507	A1	20000308	EP 1997-95280	19971205
R:	AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, FI			
PRIORITY APPLN. INFO.:			US 1997-795893	A 19970204
			WO 1997-US22206	W 19971205
OTHER SOURCE(S):			MARPAT 129:175562	
GI				



AB The invention relates to novel tricyclic tetrahydroquinoline compds. I, their salts, and combinatorial libraries containing mixts. of two or more such compds. [wherein R1 = bond, (un)substituted alk(en/yn)ylene, cycloalk(en)ylene, phenylene, naphthylene, heterocycle, heteroaryl, amino, CH2CONH, (CH2)qAr(CH2)q; p, q = 0-6 but both cannot be 0; Ar = (un)substituted Ph or heteroaryl; R2, R3, R4 = H, halo, (un)protected OH, cyano, NO2, (un)substituted alk(en/yn)yl, alkoxy, cycloalk(en)yl, heterocyclyl, phenylalkyl, Ph, naphthyl, etc.; R5 = H, (un)substituted alk(en/yn)yl, cycloalk(en)yl, Ph, naphthyl, phenylalkyl, (un)protected CO2H, acyl, heterocyclyl, etc.; R6 = H, (un)substituted alkyl,


L4 ANSWER 193 OF 261 CAPLUS COPYRIGHT 2004 ACS ON STN (Continued)
 phenylalkyl, acyl, PhSO2, alkylsulfonyl, alkylaminocarbonyl, PhNHCO; n = 1-3; Y = CO2H, OH, SH, NHR7, CONHR7, CH2OH, CH2NH2, CH2NHR7; R7 = H, (un)substituted alkyl, or functionalized resin; R1 must be present and R5 = Ph when Y = CO2H. The invention also relates to the generation of such libraries. In 2 examples, libraries of 2774 and approx. 17,000 compds. I were prepd. as mixed sublibraries. Data for control compds. (samples of individually known intermediates and products, cleaved from simultaneously processed control resins) are given. For instance, tea-bags of MBHA resin were each coupled with one of 19 aminobenzoic acids, such as 4-aminobenzoic acid. Diagnostic cleavage of each of these resins with HF gave 19 aminobenzamide controls in 34-99% yield. The 19 resins were mixed together and placed in new tea-bags, then condensed with 73 different aldehydes, and finally cyclized with cyclopentadiene. Cleavage of the resin-bound products with HF gave approx. 73 mixts. of 38 compds. (counting sep. enantiomers). Individual control samples of products, such as II (R5 = H, CH2Cl, cyclohexyl, CO2H, (un)substituted Ph, etc.), were typically obtained in 50-100% yield by reactions of pure, resin-bound 4-aminobenzoic acid control samples in sibling tea-bags. Potential applications of I (no data) may include use as antibacterials or analgesics.

IT 211376-13-5P

RL: SPN (Synthetic preparation); PREP (Preparation)
 (resin-cleavage control intermediate; preparation of tricyclic tetrahydroquinoline derivs. and combinatorial libraries)

RN 211376-13-5 CAPLUS

CN 9H-Xanthene-9-carboxamide, N-[2-amino-1-[(4-nitrophenyl)methyl]-2-oxoethyl]- (9CI) (CA INDEX NAME)



REFERENCE COUNT: 2 THERE ARE 2 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 194 OF 261 CAPLUS COPYRIGHT 2004 ACS ON STN
 ACCESSION NUMBER: 1998:507698 CAPLUS
 DOCUMENT NUMBER: 129:245476
 TITLE: Conformationally constrained opioid peptide analogs with novel activity profiles
 AUTHOR(S): Schiller, Peter W.; Schmidt, Ralf; Weltrowska, Grazyna; Berzowska, Irena; Nguyen, Thi M.-D.; Dupuis, Sebastien; Chung, Nga N.; Lemieux, Carole; Wilkes, Brian C.; Carpenter, Katharine A.
 CORPORATE SOURCE: Laboratory of Chemical Biology and Peptide Research, Clinical Research Institute of Montreal, Montreal, QC, H2W 1R7, Can.
 SOURCE: Letters in Peptide Science (1998), 5(2-3), 209-214
 CODEN: LPSCEM; ISSN: 0929-5666
 PUBLISHER: Kluwer Academic Publishers
 DOCUMENT TYPE: Journal
 LANGUAGE: English

AB Novel conformationally constrained opioid peptide analogs, having properties as δ antagonist, mixed μ agonist/ δ antagonist or δ agonist, were developed. TIP(P)-related δ antagonists showed unprecedented δ antagonist potency and δ receptor selectivity, and may have potential for use in analgesia in combination with μ agonists. A definitive model of their δ receptor-bound conformation was developed. Three prototype mixed μ agonist/ δ antagonists were discovered. They represent the only known compds. with this pharmacol. profile and, as expected, one of them was shown to be a potent analgesic and to produce no dependence and less tolerance than morphine. Novel dipeptide derivs. turned out to be potent and selective δ agonists. Because of their low mol. weight and lipophilic character, these compds. may cross the blood-brain barrier and, thus, may have potential as centrally acting analgesics.


IT 207342-55-0

RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); BIOL (Biological study)
 (activity profiles of conformationally constrained opioid peptide analogs)

RN 207342-55-0 CAPLUS

CN L-Phenylalanine, (3S)-2-[(2S)-2-amino-3-(4-hydroxyphenyl)propyl]-1,2,3,4-tetrahydro-3-methyl-3-isoquinolinecarboxyl-L-phenylalanyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



REFERENCE COUNT: 20 THERE ARE 20 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT



L4 ANSWER 194 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)

L4 ANSWER 195 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
 ACCESSION NUMBER: 1998:479021 CAPLUS
 DOCUMENT NUMBER: 129:122868
 TITLE: Preparation of peptidylboronic ester and acid compounds as proteasome inhibitors
 INVENTOR(S): Adams, Julian; Ma, Yu-Ting; Stein, Ross; Baevsky, Matthew; Grenier, Louis; Plamondon, Louis
 PATENT ASSIGNEE(S): Proscript, Inc., USA
 SOURCE: U.S., 37 pp., Cont.-in-part of U.S. Ser. No. 442,581.
 CODEN: USXXAM
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 3
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 5780454	A	19980714	US 1995-549318	19951027
US 6083903	A	20000704	US 1995-442581	19950516
US 6066730	A	20000523	US 1998-85404	19980526
US 6297217	B1	20011002	US 2000-490511	20000125
US 6465433	B1	20021015	US 2001-953540	20010914
US 2002173488	A1	20021121	US 2002-100295	20020318
US 6548668	B2	20030415		
US 6617317	B1	20030909	US 2002-125997	20020419
US 2003199561	A1	20031023	US 2003-392165	20030319
PRIORITY APPLN. INFO.:			US 1994-330525	R2 19941028
			US 1995-442581	A2 19950516
			US 1995-549318	A3 19951027
			US 1998-85404	A3 19980526
			US 2000-490511	A1 20000125
			US 2001-953540	A1 20010914
			US 2002-100295	A1 20020318
			US 2002-125997	A1 20020419

OTHER SOURCE(S): MARPAT 129:122868
 GI

L4 ANSWER 195 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)


L4 ANSWER 195 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)
 RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

AB Disclosed herein is a method for reducing the rate of degradation of proteins in an animal comprising contacting cells of the animal with certain boronic ester and acid compds I [P = aryl-, aralkyl-, heteroaryl-, or heteroarylalkylcarbonyl or -sulfonyl; B1 = N, CH; X1, X2 = CONH, CH(OH)CH2, COCH2; n = 0, 1, 2; R = H, alkyl, R1 or R2 (for n = 0) may form a ring; R1, R2, R3 = H, alkyl, cycloalkyl, aryl, etc.; Z1, Z2 = alkyl, hydroxy, alkoxy, aryloxy; Z1Z2 may form a moiety derived from a dihydroxy compound]. Also disclosed herein are novel boronic ester and acid compds., their synthesis and uses. Thus, peptidylboronic acid II was prepared by coupling pinanediol leucine boronate ester III with N-Boc-β-(1-naphthyl)-L-alanine, followed by deprotection, acylation with 4-morpholinecarbonyl chloride, and cleavage of the pinanediol moiety. II inhibited proteasome 20S with Ki = 0.18 nM.

IT 179324-53-9P
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)
 (preparation of peptidylboronic ester and acid compds. as proteasome inhibitors)

RN 179324-53-9 CAPLUS
 CN Boronic acid, [(1R)-3-methyl-1-[(2S)-1-oxo-3-phenyl-2-[(2-quinolinylcarbonyl)amino]propyl]amino]butyl]- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



REFERENCE COUNT: 19 THERE ARE 19 CITED REFERENCES AVAILABLE FOR THIS

L4 ANSWER 196 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
 ACCESSION NUMBER: 1998:427361 CAPLUS
 DOCUMENT NUMBER: 129:156465
 TITLE: A potent dipeptide inhibitor of dipeptidyl peptidase IV
 AUTHOR(S): Yamada, Masaki; Okagaki, Chieko; Higashijima, Takanori; Tanaka, Sumiko; Ohnuki, Tetsuo; Sugita, Takahisa
 CORPORATE SOURCE: Lead Generation Research Laboratory, Tanabe Seiyaku Co., Ltd., Osaka, 532-8505, Japan
 SOURCE: Bioorganic & Medicinal Chemistry Letters (1998), 8(12), 1537-1540
 CODEN: BMCLEB; ISSN: 0960-894X
 PUBLISHER: Elsevier Science Ltd.
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 AB A series of novel potent inhibitors of dipeptidyl peptidase IV (DPP-IV) has been developed. A brief structure-activity relation of the inhibitors was investigated. The dipeptide TSL-225, tryptophyl-1,2,3,4-tetrahydroisoquinoline-3-carboxylic acid, was identified with the critical structure for the inhibitory activity.
 IT 207228-59-9P
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation)
 (preparation of potent dipeptide inhibitors of dipeptidyl peptidase IV in relation to structure)
 RN 207228-59-9 CAPLUS
 CN L-Phenylalanine, L-tryptophyl-(3S)-1,2,3,4-tetrahydro-3-isoquinolinecarbonyl- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



REFERENCE COUNT: 27 THERE ARE 27 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 197 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
 ACCESSION NUMBER: 1998:293475 CAPLUS
 DOCUMENT NUMBER: 129:4862
 TITLE: Preparation of amino acid-containing tetrahydroisoquinoline derivatives as dipeptidyl peptidase IV inhibitors
 INVENTOR(S): Sugita, Takahisa; Ohnuki, Tetsuo; Yamada, Masaki; Tanaka, Sumiko; Nonaka, Nobuaki; Asai, Yasuyuki
 PATENT ASSIGNEE(S): Tanabe Seiyaku Co., Ltd., Japan; Sugita, Takahisa; Ohnuki, Tetsuo; Yamada, Masaki; Tanaka, Sumiko; Nonaka, Nobuaki; Asai, Yasuyuki
 SOURCE: PCT Int. Appl., 87 pp.
 CODEN: PIXXD2
 DOCUMENT TYPE: Patent
 LANGUAGE: Japanese
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 9818763	A1	19980507	WO 1997-JP3804	19971022
W: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, HU, ID, IL, IS, KE, KG, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, AM, AZ, BY, BG, KZ, MD, RU, TJ, TM				
RW: GH, KE, LS, MW, SD, SE, UG, ZW, AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CO, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG				
JP 10182613	A2	19980707	JP 1997-288322	19971021
AU 9747218	A1	19980522	AU 1997-47218	19971022
PRIORITY APPLN. INFO.:			JP 1996-284328	19961025
			WO 1997-JP3804	19971022
OTHER SOURCE(S):			MARPAT 129:4862	
GI				



AB The title compds. 1 [R1 = amino protecting group, etc.; R2 = optionally protected hydroxy, etc.; R3 - R6 = H, hydroxy, alkoxyl are prepared. In an in vitro test, 21 compds. of this invention showed inhibiting activity against dipeptidyl peptidase IV. (3S)-2-(L-Tryptophyl)-1,2,3,4-tetrahydroisoquinoline-3-carboxylic acid hydrochloride at 30 mg/kg/day s.c. for 18 days gave significant inhibition of exptl. arthritis in rats.
 IT 207228-59-9P

L4 ANSWER 197 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)
 (prepn. of amino acid-contg. tetrahydroisoquinoline deriva. as dipeptidyl peptidase IV inhibitors)
 RN 207228-59-9 CAPLUS
 CN L-Phenylalanine, L-tryptophyl-(3S)-1,2,3,4-tetrahydro-3-isoquinolinecarbonyl- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



REFERENCE COUNT: 5 THERE ARE 5 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 198 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
 ACCESSION NUMBER: 1998:277671 CAPLUS
 DOCUMENT NUMBER: 128:289517
 TITLE: Rapid characterization of combinatorial libraries using electrospray ionization Fourier transform ion cyclotron resonance mass spectrometry
 AUTHOR(S): Fang, A. S.; Vouras, P.; Stacey, C. C.; Kruppa, G. H.; Laukien, F. H.; Wintner, E. A.; Carell, T.; Rebek, J., Jr.
 CORPORATE SOURCE: Department of Chemistry, Barnett Institute, Northeastern University, Boston, MA, 02115, USA
 SOURCE: Combinatorial Chemistry and High Throughput Screening (1998), 1(1), 23-33
 CODEN: CCHSFL; ISSN: 1386-2073
 PUBLISHER: Bentham Science Publishers
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 AB The relatively new field of combinatorial chemical has enabled researchers to create large mixts. of compds. that can be screened for leads in developing potential drug candidates. The new synthetic method has also created a need for better procedures to analyze the complex mixts. that are generated. The immediate goal in most cases is to verify the synthetic procedure and to determine the purity and completeness of the library sample before binding studies are initiated. The authors report here a method to rapidly characterize small-mol. combining a core mol. bearing two acid chloride functionalities with various amino acids to generate libraries of 36, 78 and 120 components. Using electrospray ionization Fourier transform ICR mass spectrometry (ESI-FTICR-MS) the authors were able to identify 70-80% of the library components. All samples were analyzed as mixts. by direct infusion without chromatog. separation. Also, nominally isobaric components could be resolved and identified through exact mass assignments without tandem mass spectrometry. ESI-FTICR-MS is a rapid and convenient tool for the characterization of small-mol. libraries. The method is especially useful for the anal. of larger libraries that contain many nominally isobaric components and impurities.
 IT 178916-07-9
 RL: ANT (Analyte); PRP (Properties); ANST (Analytical study)
 (rapid characterization of combinatorial libraries using electrospray ionization Fourier transform ion cyclotron resonance mass spectrometry)
 RN 178916-07-9 CAPLUS
 CN L-Phenylalanine, N-[(2,7-bis(1,1-dimethylethyl)-5-[(2-methoxy-2-oxoethyl)amino]carbonyl)-9,9-dimethyl-9H-xanthen-4-yl]carbonyl-, 1,1-dimethylethyl ester (9CI) (CA INDEX NAME)

Absolute stereochemistry.




REFERENCE COUNT: 20 THERE ARE 20 CITED REFERENCES AVAILABLE FOR THIS

L4 ANSWER 198 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)
RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT


L4 ANSWER 199 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
ACCESSION NUMBER: 1998:2652 CAPLUS
DOCUMENT NUMBER: 129:4851
TITLE: The receptor-bound conformation of H-Tyr-Tic-(Phe-Phe)-OH-related δ -opioid antagonists contains all trans peptide bonds
AUTHOR(S): Wilkes, Brian C.; Nguyen, Thi M. -D.; Weltrowska, Grazyna; Carpenter, Katharine A.; Lemieux, Carole; Chung, Nga N.; Schiller, Peter W.
CORPORATE SOURCE: Laboratory of Chemical Biology and Peptide Research, Clinical Research Institute of Montreal, Montreal, QC, H2W 1R7, Can.
SOURCE: Journal of Peptide Research (1998), 51(5), 386-394
CODEN: JPERFA; ISSN: 1397-002X
PUBLISHER: Munksgaard International Publishers Ltd.
DOCUMENT TYPE: Journal
LANGUAGE: English
AB Two different models for the receptor-bound conformation of δ -opioid peptide antagonists containing the N-terminal dipeptide segment "Tyr-Tic" (Tic = 1,2,3,4-tetrahydroisoquinoline-3-carboxylic acid) have been proposed. Both models are based on spatial overlap of the Tyr1 and Tic2 aromatic rings and N-terminal amino group with the corresponding aromatic rings and nitrogen atom of the nonpeptide δ -antagonist naltrindole. However, in one model the peptide bond between the Tyr1 and Tic2 residues assumes the trans conformation, whereas in the other it is in the cis conformation. To distinguish between these two models, the authors prepared the two peptides H-Tyrw(CH2NH)Tic-Phe-Phe-OH and H-Tyrw(CH2NH)MeTic-Phe-Phe-OH (MeTic = 3-methyl-1,2,3,4-tetrahydroisoquinoline-3-carboxylic acid) in which a cis peptide bond between the Tyr and Tic (or MeTic) residues is sterically forbidden. Both compds. turned out to be moderately potent δ -opioid antagonists in the mouse vas deferens assay. A mol. mechanics study performed with both peptides resulted in low-energy conformations in which the torsional angle (" ω 1") of the reduced peptide bond between Tyr and Tic (or MeTic) had a value of 180° (trans conformation) and which were in good agreement with the proposed model with all trans peptide bonds. Also, this study confirmed that neither of these two peptides could assume low-energy conformations in which " ω 1" had a value of 0° (cis conformation). Conformers with that same bond in the gauche conformation (" ω 1" = -60°) were also identified, but were higher in energy and showed no spatial overlap with naltrindole. Thus, it is concluded that the receptor-bound conformation of δ -peptide antagonists containing an N-terminal "Tyr-Tic" dipeptide segment must have all trans peptide bonds.
IT 207342-54-9P
RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); PRP (Properties); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation)
(conformation study of the receptor bound "Tyr-Tic" peptide segment of δ -opioid antagonists)
RN 207342-54-9 CAPLUS
CN L-Phenylalanine, (3S)-2-[(2S)-2-amino-3-(4-hydroxyphenyl)propyl]-1,2,3,4-tetrahydro-3-isoquinolinecarboxyl-L-phenylalanyl- (9CI) (CA INDEX NAME)
Absolute stereochemistry.

L4 ANSWER 199 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)



REFERENCE COUNT: 31 THERE ARE 31 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT


L4 ANSWER 200 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
ACCESSION NUMBER: 1998:265646 CAPLUS
DOCUMENT NUMBER: 129:4850
TITLE: Synthesis of cyclic dipeptide templates, their incorporation into peptides and studies on their conformational and biological properties
AUTHOR(S): Asche, Geert; Kunz, Horst; Nar, Herbert; Koppen, Herbert; Briem, Hans; Pook, Karl-Heinz; Schiller, Peter W.; Chung, Nga N.; Lemieux, Carole; Esser, Franz
CORPORATE SOURCE: Departments of Medicinal Chemistry and Analytical Sciences, Boehringer Ingelheim, Ingelheim, Germany
SOURCE: Journal of Peptide Research (1998), 51(5), 323-336
CODEN: JPERFA; ISSN: 1397-002X
PUBLISHER: Munksgaard International Publishers Ltd.
DOCUMENT TYPE: Journal
LANGUAGE: English
GI



AB This study investigated the diastereoselective synthesis of three dipeptide templates I-III [R = Cl3CCH2O2C, PhCH2O2C (Chz); R1 = OH], which may be regarded as conformationally restricted analogs of H-Gly-Xaa-OH, in which Xaa constitutes an aromatic amino acid. Bond formation between α -C of Gly and the aromatic moiety was achieved by proton-catalyzed intramol. electrophilic aromatic substitution. The absolute configuration of the dipeptide templates was determined by single-crystal x-ray crystallog. or by NOE measurements. A protective group strategy was elaborated to allow their incorporation into peptide sequences by liquid phase as well as by solid-phase peptide synthesis. The templates were used to generate enkephalin analog II (R = H-Tyr-Gly, R1 = Leu-NH2), modified neurokinin antagonist III (R = N-cyclohexylcarbonylglycyl, R1 = Phe-NMe2) and dermorphin derivs. I and II (R = H-Tyr-D-Ala, Phe, R1 = Pro-Ser-NH2). Mol. dynamic simulations of enkephalin analog II (R = H-Tyr-Gly, R1 = Leu-NH2) and neurokinin antagonist III (R = N-cyclohexylcarbonylglycyl, R1 = Phe-NMe2) revealed the preference for a turn-like motif for the enkephalin analog. The biol. activity, as investigated by resp. receptor binding and functional assays, was strongly diminished with all four derivs., indicating that their receptor-relevant mol. geometries lie outside the examined conformational space.
IT 207443-93-4P
RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation)
(preparation, conformation, and receptor-binding of conformationally constrained aromatic dipeptide template-containing peptides)
RN 207443-93-4 CAPLUS
CN Azepino[4,5-b]indole-2-carboxamide, 5-[[[(cyclohexylcarbonyl)amino]acetyl]

L4 ANSWER 200 OF 261 CAPLUS COPYRIGHT 2004 ACS ON STN (Continued)
 AMINO-N-[(1S)-2-(dimethylamino)-2-oxo-1-(phenylmethyl)ethyl]-1,2,3,4,5,6-hexahydro-4-oxo-, (2R,5R)- (9CI) (CA INDEX NAME)

Absolute stereochemistry. Rotation (-).




REFERENCE COUNT: 23 THERE ARE 23 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 201 OF 261 CAPLUS COPYRIGHT 2004 ACS ON STN
 ACCESSION NUMBER: 1998:233609 CAPLUS
 DOCUMENT NUMBER: 128:295021
 TITLE: Solid-phase synthesis of substituted glutamic acid derivatives via Michael addition reactions
 AUTHOR(S): Dominguez, Esteban; O'donnell, Martin J.; Scott, William L.
 CORPORATE SOURCE: Centro de Investigacion Lilly, Madrid, 28130, Spain
 SOURCE: Tetrahedron Letters (1998), 39(15), 2167-2170
 CODEN: TELEAY; ISSN: 0040-4039
 PUBLISHER: Elsevier Science Ltd.
 DOCUMENT TYPE: Journal
 LANGUAGE: English

AB The conjugate addition of Michael acceptors to the resin-bound benzophenone imine of glycine, Ph2C:NCH2CO2-Resin (I), leads to a variety of racemic unnatural glutamic acid deriva. This new approach expands the scope of unnatural amino acid and peptide synthesis. For example, RCONHCH(CO2H)CH(C6H4NO2-4)CH2CO2EC (R = 2-quinolinyl) was synthesized in 88% yield from I, EtO2CCH:CHC6H4NO2-4 and quinaldic acid.

IT 206070-97-5P
 RL: SPN (Synthetic preparation); PREP (Preparation)
 (solid-phase synthesis of substituted glutamic acids via Michael addition reactions)
 RN 206070-97-5 CAPLUS
 CN Glutamic acid, 3-(4-nitrophenyl)-N-(2-quinolinylcarbonyl)-, 5-ethyl ester (9CI) (CA INDEX NAME)




REFERENCE COUNT: 12 THERE ARE 12 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 202 OF 261 CAPLUS COPYRIGHT 2004 ACS ON STN
 ACCESSION NUMBER: 1998:197358 CAPLUS
 DOCUMENT NUMBER: 128:257695
 TITLE: Preparation of modified amino acids and their use as calcitonin gene-related peptide antagonists in pharmaceutical compositions
 INVENTOR(S): Rudolf, Klaus; Eberlein, Wolfgang; Engel, Wolfgang; Pieper, Helmut; Doods, Henri; Hallermayer, Gerhard; Entzeroth, Michael; Wienen, Wolfgang
 PATENT ASSIGNEE(S): Karl Thomae G.m.b.H., Germany
 SOURCE: FCT Int. Appl., 461 pp.
 CODEN: PIXXD2
 DOCUMENT TYPE: Patent
 LANGUAGE: German
 FAMILY ACC. NUM. COUNT: 2
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 9811128	A1	19980319	WO 1997-EP4862	19970908
W: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM				
RW: GH, KE, LS, MW, SD, SZ, UG, ZW, AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG				
DE 19636623	A1	19980312	DE 1996-19636623	19960910
DE 19720011	A1	19981119	DE 1997-19720011	19970514
AU 9741196	A1	19980402	AU 1997-41196	19970908
AU 721035	B2	20000622		
EP 927192	A1	19990707	EP 1997-938928	19970908
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO				
BR 9712023	A	19990831	BR 1997-12023	19970908
JP 2000505100	T2	20000425	JP 1998-513227	19970908
JP 3483893	B2	20040106		
NO 9901130	A	19990505	NO 1999-1130	19990309
KR 2000044040	A	20000715	KR 1999-702008	19990310
US 6344449	B1	20020205	US 1999-254281	19991012
US 2001036946	A1	20011101	US 2001-789391	20010221
US 2003069231	A1	20030410	US 2002-119875	20020410
PRIORITY APPLN. INFO.:			DE 1996-19636623 A	19960910
			DE 1997-19720011 A	19970514
			WO 1997-EP4862 W	19970908
			US 1999-254281 A1	19991012
			US 2001-789391 A1	20010221
OTHER SOURCE(S):		MARPAT 128:257695		
GI				

L4 ANSWER 202 OF 261 CAPLUS COPYRIGHT 2004 ACS ON STN (Continued)



AB The invention concerns modified amino acids of general formula I [A = bond, CX; Z = CH2, NR1; R1 = H, alkyl, phenyl-alkyl; X = O, H, N; n = 1-2; m = 0-1; R = (substituted)alkyl; R2 = Ph, (substituted) (hetero) (bi)cycle; R3 = H, (substituted)alkyl, Ph, pyridinyl; R4 = H, (substituted)alkyl, R3R4 = (hetero)cycle; R5 = H, alkyl, alkoxy-carbonyl, PhCH2], pharmaceuticals containing these compounds, their use and the method for their production, as well as their use for the production and purification of antibodies and


as marked compounds in RIA and ELISA assays and as diagnostic or analytic auxiliary agents in neurotransmitter research. Thus, 3,5-dibromo-N2-[4-(1,3-dihydro-2(2H)-oxo-benzimidazol-1-yl)-1-piperidinyl]carbonyl-D-tyrosine was reacted with 1-(4-pyridinyl)-piperazine, to give II (224). Title compounds show human calcitonin gene related peptide (CGRP) antagonist activity; in in-vitro binding studies with SK-N-MC-cells, 1 had IC50 510000 nM, and in the same system, had CGRP-antagonist activity at doses from 10-11 to 10-6 M.

IT 204698-94-2P
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)
 (preparation of amino acids and their use as calcitonin gene-related peptide antagonists in pharmaceutical compns.)

RN 204698-94-2 CAPLUS
 CN 3H-3-Benzazepine-3-carboxamide, N-[2-[[5-amino-1-[[4-(4-pyridinyl)-1-piperazinyl]carbonyl]pentyl]amino]-1-[[3,5-dibromo-4-hydroxyphenyl]methyl]-2-oxoethyl]-1,2,4,5-tetrahydro-, [R-(R*,S*)]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

L4 ANSWER 202 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)




REFERENCE COUNT: 10 THERE ARE 10 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 203 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
 1998:186625 CAPLUS
 DE 1998-186625
 DOCUMENT NUMBER: 128:230701
 TITLE: Preparation of varied amino acids as calcitonin gene-related peptide antagonists in pharmaceutical compositions
 INVENTOR(S): Rudolf, Klaus; Eberlein, Wolfgang; Engel, Wolfhard; Pieper, Helmut; Doods, Henri; Hallermayer, Gerhard; Entzeroth, Michael; Wienen, Wolfgang
 PATENT ASSIGNEE(S): Karl Thomas G.m.b.H., Germany
 SOURCE: Ger. Offen. 142 pp.
 CODEN: GWXXBX
 DOCUMENT TYPE: Patent
 LANGUAGE: German
 FAMILY ACC. NUM. COUNT: 2
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
DE 19636623	A1	19980312	DE 1996-19636623	19960910
WO 9811128	A1	19980319	WO 1997-EP4862	19970908
W: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZM, ZW, AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG				
AU 9741196	A1	19980402	AU 1997-41196	19970908
AU 721035	B2	20000622		
EP 927192	A1	19990707	EP 1997-938928	19970908
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO				
BR 9712023	A	19990831	BR 1997-12023	19970908
CN 1230196	A	19990929	CN 1997-197772	19970908
CN 1129605	B	20031203		
JP 2000505100	T2	20000425	JP 1998-513227	19970908
JP 3483893	B2	20040106		
JP 2003300959	A2	20031021	JP 2003-21750	19970908
ZA 9708083	A	19991217	ZA 1997-8083	19970909
TW 477752	B	20020301	TW 1997-86113120	19970910
TW 498076	B	20020811	TW 2000-89125839	19970910
NO 9901130	A	19990505	NO 1999-1130	19980309
US 6344449	B1	20020205	US 1999-254281	19991012
PRIORITY APPLN. INFO.: DE 1996-19636623 A 19960910 DE 1997-19720011 A 19970514 JP 1998-513227 A3 19970908 WO 1997-EP4862 W 19970908				
OTHER SOURCE(S): MARPAT 128:230701 GI				

L4 ANSWER 203 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)




AB Title compds. RCOZCR1R2C((X)ANR3R4 [(I); R = (substituted) alkyl; R1 = H, alkyl, PhCH2; R2 = (CO)m(CH2)nR5; m = 0, 1; n = 1, 2; R5 = Ph, heterocycle; X = O, (H,H); Z = CH2, NR6; R6 = H, alkyl, phenyl-alkyl; A = bond, proline; R3 = H, substituted alkyl, Ph, pyridinyl; R4 = H, substituted alkyl; NR3R4 = (substituted) heterocycle], useful as calcitonin gene-related peptide (CGRP) antagonists, were prepared. Thus, 3,5-dibromo-N2-[4-((1,3-dihydro-2(2H)-oxo-benzimidazol-1-yl)-1-piperidinyl)carbonyl-D-tyrosine was reacted with 1-(4-pyridinyl)-piperazine, to give II (22%). In in-vitro binding studies with human CGRP-receptors, I had IC50 \leq 10000 nM; in CGRP-antagonist in vitro tests, I was effective at doses from 10⁻¹¹ to 10⁻⁵ M.

IT 204698-94-29
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)
 (preparation of amino acids and their use as calcitonin gene-related peptide antagonists in pharmaceutical compns.)

RN 204698-94-2 CAPLUS
 CN 3H-3-Benzazepine-3-carboxamide, N-[2-[[[5-amino-1-[[4-(4-pyridinyl)-1-piperazinyl]carbonyl]pentyl]amino]-1-[(3,5-dibromo-4-hydroxyphenyl)methyl]-2-oxoethyl]-1,2,4,5-tetrahydro-, (R-(R*,S*))]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.


L4 ANSWER 203 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)



L4 ANSWER 204 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
 ACCESSION NUMBER: 1998:100848 CAPLUS
 DOCUMENT NUMBER: 128:243960
 TITLE: 8-Hydroxy-7-substituted quinolines as anti-viral agents
 INVENTOR(S): Vaillancourt, Valerie A.; Romines, Karen R.; Romero, Arthur G.; Tucker, John A.; Strohhach, Joseph W.; Bezencon, Olivier; Thaisrivongs, Suvit; et al.
 PATENT ASSIGNEE(S): Pharmacia & Upjohn Co., USA
 SOURCE: PCT Int. Appl., 280 pp.
 CODEN: PIXXD2
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 9811073	A1	19980319	WO 1997-US15310	19970905
W:	AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM			
RW:	GH, KE, LS, MW, SD, SZ, UG, ZW, AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG			
AU 9741721	A1	19980402	AU 1997-41721	19970905
EP 927164	A1	19990707	EP 1997-939690	19970905
R:	AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO			
US 6310211	B1	20011030	US 1997-924683	19970905
JP 2002505660	T2	20020219	JP 1998-513685	19970905
US 6211376	B1	20010403	US 1999-425789	19991022
US 6252080	B1	20010626	US 1999-425564	19991022
US 6500842	B1	20021231	US 2001-14780	20011023
PRIORITY APPLN. INFO.:			US 1996-258709 P	19960910
			US 1997-50720P P	19970625
			US 1997-924683 A3	19970905
			WO 1997-US15310 W	19970905

OTHER SOURCE(S): MARPAT 128:243960
 GI



AB The present invention provides for 8-hydroxy-7-substituted quinoline compds. I (R = alkyl, alkylamino, alkoxyalkyl, etc.; R1 = H, F, Cl, Br,


L4 ANSWER 205 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
 ACCESSION NUMBER: 1998:172195 CAPLUS
 DOCUMENT NUMBER: 128:240972
 TITLE: Ochratoxin A acts as a photoactivatable DNA cleaving agent
 AUTHOR(S): Gillman, Ivan G.; Yezek, Jennifer M.; Manderville, Richard A.
 CORPORATE SOURCE: Department of Chemistry, Wake Forest University, Winston-Salem, NC, 27109-7486, USA
 SOURCE: Chemical Communications (Cambridge) (1998), (6), 647-648
 CODEN: CHCOFS; ISSN: 1359-7345
 PUBLISHER: Royal Society of Chemistry
 DOCUMENT TYPE: Journal
 LANGUAGE: English

AB The ability of ochratoxin A to photoinduce DNA cleavage is described; in the presence of DNA the photoreaction yields the non-chlorinated derivative, ochratoxin B, while a hydroquinone derivative is produced under anaerobic conditions.

IT 205034-32-8P
 RL: BPN (Biosynthetic preparation); BIOL (Biological study); PREP (Preparation)
 (Ochratoxin A acts as a photoactivatable DNA cleaving agent)

RN 205034-32-8 CAPLUS
 CN L-Phenylalanine, N-[[[(3R)-3,4-dihydro-5,8-dihydroxy-3-methyl-1-oxo-1H-2-benzopyran-7-yl]carbonyl]- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



REFERENCE COUNT: 23 THERE ARE 23 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 204 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)
 CF3, etc.; R2 = H, alkyl, OH, arylalkenyl, etc.; R3 = H, OH, CF3, Cl-C3alkyl) are prepd. as anti-viral agents. Specifically, these compds. have anti-viral activity against the herpes virus, cytomegalovirus (CMV). Many of these compds. are also active against other herpes viruses, such as the varicella zoster virus, the Epstein-Barr virus, the herpes simplex virus and the human herpes virus type 8 (HHV-8).
 IT 205038-81-9P
 RL: BAC (Biological activity or effector, except adverse); ESU (Biological study, unclassified); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)
 (Preparation of 8-hydroxy-7-substituted quinolines as anti-viral agents)
 RN 205038-81-9 CAPLUS
 CN L-Tyrosine, N-[(8-hydroxy-7-quinolinyl)carbonyl]-, 1,1-dimethylethyl ester (9CI) (CA INDEX NAME)

Absolute stereochemistry.



REFERENCE COUNT: 2 THERE ARE 2 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 206 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
 ACCESSION NUMBER: 1998:147347 CAPLUS
 DOCUMENT NUMBER: 128:217634
 TITLE: Preparation of protease-resistant B1-bradykinin receptor antagonists for treatment of inflammatory conditions
 INVENTOR(S): Regoli, Domenico; Plante, Gerard E.; Cobeil, Fernand; Neugebauer, Witold A.; Zuccollo, Adriana; Catanzaro, Orlando L.
 PATENT ASSIGNEE(S): Universite De Sherbrooke, Can.
 SOURCE: PCT Int. Appl., 41 pp.
 CODEN: PIXXD2
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 9807746	A1	19980226	WO 1997-CA582	19970814
W:	AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM			
RW:	GH, KE, LS, MW, SD, SZ, UG, ZW, AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, CA, CN, ML, MR, NE, SN, TD, TG			
AU 9739358	A1	19980306	AU 1997-39358	19970814
PRIORITY APPLN. INFO.:			US 1996-23971P P	19960819
			WO 1997-CA582 W	19970814

OTHER SOURCE(S): MARPAT 128:217634


AB The present invention relates to novel B1-bradykinin (B1-BK) receptor antagonists A-Arg-Pro-B-Gly-Phe-Ser-C-D [I; A = H-D-Arg, Ac-Lys, H-D-Lys, H-Sar, Ac-Tyr-e-Ahx-Lys, H-Sar-Tyr-e-Ahx-Lys, H-Sar-Tyr(12-3,5)-e-Ahx-Lys; B = Pro, Hyp; C = Pro, D-1,2,3,5-tetrahydroisoquinoline-3-carbonyl (D-Tic), O-3-(2-naphthyl)alanyl (D-β-Nal); D = Leu-OH, Ile-OH; e-Ahx = NH(CH2)5CO; with the proviso that C = Pro when A does not contain e-Ahx] which have a good affinity and selectivity therefore, some of which being at least partially resistant to enzymic degradation. The synthesis of the B1 receptor is induced during inflammation. Symptoms associated with inflammation (elevated hydrostatic pressure and plasma leakage or extravasation) have been observed in diabetic animal models [streptozotocin-induced diabetes (STZ)] as well as in spontaneously hypertensive rats (SHR). The present inventors confirm the presence of B1-BK receptors in these two models. B1-BK antagonists abolished the vasoconstriction induced by B1-BK in SHR and STZ, and reduced the glycemia of diabetic animals to normal levels. The present B1-antagonists are useful for treating any condition wherein B1-receptor is expressed, particularly during inflammation, and more particularly wherein B1-receptor expression results in diabetic vasculopathy, other diabetic symptoms associated with an insulinitis and a post-capillary resistance building as a consequence of the presence of a B1-receptor.

IT 185052-06-6P


RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); PRP (Preparation); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)
 (Preparation of protease-resistant B1-bradykinin receptor antagonists for treatment of inflammatory conditions)

L4 ANSWER 206 OF 261 CAPLUS COPYRIGHT 2004 ACS ON STN (Continued)
 RN 185052-06-6 CAPLUS
 CN L-Phenylalanine, D-arginyl-L-arginyl-L-prolyl-(4R)-4-hydroxy-L-prolylglycyl-L-phenylalanyl-L-eryl-(3R)-1,2,3,4-tetrahydro-3-isoquinolinecarbonyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



PAGE 1-A



PAGE 1-B

REFERENCE COUNT: 7 THERE ARE 7 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT


L4 ANSWER 207 OF 261 CAPLUS COPYRIGHT 2004 ACS ON STN
 ACCESSION NUMBER: 1998:118 CAPLUS
 DOCUMENT NUMBER: 128:110919
 TITLE: From Micromolar to Nanomolar Affinity: A Systematic Approach To Identify the Binding Site of CGRP at the Human Calcitonin Gene-Related Peptide 1 Receptor
 AUTHOR(S): Rist, Beate; Entzeroth, Michael; Beck-Sickinger, Annette G.
 CORPORATE SOURCE: Department of Pharmacy, ETH Zuerich, Zurich, Switz.
 SOURCE: Journal of Medicinal Chemistry (1998), 41(1), 117-123
 CODEN: JMCMAH; ISSN: 0022-2623
 PUBLISHER: American Chemical Society
 DOCUMENT TYPE: Journal
 LANGUAGE: English

AB CGRP Y0-28-37 is known as a selective CGRP1 receptor antagonist. To elucidate the essential requirements for its receptor interaction, the authors performed a variety of systematic approaches by modifying the C-terminal segments CGRP Y0-28-37 and CGRP 27-37. N-Terminal and C-terminal segments have been synthesized, as well as chimeras which combine segments of CGRP, adrenomedullin, and amylin. Furthermore, the authors carried out an Ala scan, a Phe scan, a D-amino acid scan and a Pro scan of CGRP 27-37. Addnl., single amino acids were replaced by those with similar biophys. properties. Receptor binding studies of all analogs were performed at human neuroblastoma cells SK-N-MC, which selectively express the hCGRP1 receptor. On the basis of the obtained results, the authors synthesized a series of ligands with multiple amino acid replacements to optimize the exchange at each position. This approach yielded to a series of high affinity ligands, including [D31,P34,F35] CGRP 27-37 which exhibits a 100-fold increased affinity compared to the unmodified segment. So far, this is the smallest CGRP analog that shows affinity in the nanomolar range.


IT 201613-69-6
 RL: BPR (Biological process); BSU (Biological study, unclassified); PRP (Properties); BIOL (Biological study); PROC (Process)
 (from micromolar to nanomolar affinity: a systematic approach to identify the binding site of CGRP at the human calcitonin gene-related peptide 1 receptor)
 RN 201613-69-6 CAPLUS
 CN L-Phenylalaninamide, L-phenylalanyl-L-valyl-L-prolyl-L-threonyl-L-aspartyl-L-valylglycyl-(3S)-1,2,3,4-tetrahydro-3-isoquinolinecarbonyl-L-phenylalanyl-L-alanyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

L4 ANSWER 207 OF 261 CAPLUS COPYRIGHT 2004 ACS ON STN (Continued)



PAGE 1-A




PAGE 1-B

REFERENCE COUNT: 45 THERE ARE 45 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 208 OF 261 CAPLUS COPYRIGHT 2004 ACS ON STN
 ACCESSION NUMBER: 1997:798601 CAPLUS
 DOCUMENT NUMBER: 128:13436
 TITLE: Preparation of tryptophan urea derivatives as tachykinin receptor antagonists
 INVENTOR(S): Maccoss, Malcolm; Oi, Hongbo; Shah, Shrenik K.
 PATENT ASSIGNEE(S): Merck and Co., Inc., USA
 SOURCE: Brit. UK Pat. Appl., 47 pp.
 CODEN: BAXXDU
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
GB 2311523	A1	19971001	GB 1997-5861	19970321
PRIORITY APPLN. INFO.:			US 1996-14003P	P 19960325
			GB 1996-11786	A 19960606

OTHER SOURCE(S): MARPAT 128:13436
 GI




AB Substituted title azacycles I [Z = N, R = CH2Ph, Ph, 2-MeOC6H4, 2-MeC6H4, R1 = absent; Z = C, R = Ph, R1 = NHOMe; R = CH2Ph, 2-oxo-1,2,3,4-tetrahydroquinazolin-1-yl, R1 = H; R2R1 = spiro-fused 1-indanyl, 3-indenyl, 1-methylsulfonyl-2,3-dihydroindol-3-yl, 1-acetyl-2,3-dihydroindol-3-yl; R2 = OCH2Ph wherein the Ph is substituted with 0-3 groups halo, Me, or CF3; or R2 = NR3-C1-4-alkylphenyl wherein the C1-4-alkyl may be linear or branched and the Ph may be substituted with 0-3 groups halo, Me, OMe, CF3; R3 = H, Me, Et] and pharmaceutically acceptable salts thereof are tachykinin receptor antagonists useful in the treatment of inflammatory diseases, pain or migraine, and asthma. In particular, compds. I are neurokinin antagonists. Thus, amidation of 1.967 g Boc-Trp-OH (Boc = MeO2C) with 0.87 mL MeNHCH2Ph gave 2.56 g of the corresponding amide, which underwent deprotection with CF3CO2H, condensation with carbonyldiimidazole, and urea formation with spiro[1H-indene-1,4'-piperidine] hydrochloride to give title compound II (L-743,516). I and related Trp deriva. showed IC50 values of >1000 to 1 nM for human neurokinin 1 (NK1) antagonist activity.


IT 199110-44-6P

L4 ANSWER 208 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)
 (prepn. of tryptophan urea derivs. as tachykinin receptor antagonists)
 RN 199110-44-6 CAPLUS
 CN Spiro[3H-indole-3,4'-piperidine]-1'-carboxamide, 1,2-dihydro-N-[(1S)-1-[(methyl(phenylmethyl)amino)carbonyl]-3-phenylpropyl]-1-(methylsulfonyl)-(9CI) (CA INDEX NAME)

Absolute stereochemistry.



L4 ANSWER 209 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
 ACCESSION NUMBER: 1997:794681 CAPLUS
 DOCUMENT NUMBER: 128:75653
 TITLE: Tandem UPS: sequential mono- and dialkylation of resin-bound glycine via automated synthesis
 AUTHOR(S): Griffith, David L.; O'Donnell, Martin J.; Pottorf, Richard S.; Scott, William L.; Porco, John A., Jr.
 CORPORATE SOURCE: Argonaut Technologies, San Carlos, CA, 94070, USA
 SOURCE: Tetrahedron Letters (1997), 38(51), 8821-8824
 CODEN: TELEAY; ISSN: 0040-4039
 PUBLISHER: Elsevier Science Ltd.
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 AB A method has been developed for the synthesis of racemic α,α -disubstituted amino acids by a tandem alkylation process ("Tandem UPS") on solid-support. Consecutive alkylations of Wang resin-bound benzophenone imines of glycine afforded unnatural, disubstituted amino acid derivs. Automated chemical synthesis was used to efficiently optimize conditions for both formation and hydrolysis of resin-bound disubstituted benzophenone imines and to generate a matrix of disubstituted amino acid derivs.
 IT 200573-87-1P
 RL: SPN (Synthetic preparation); PREP (Preparation)
 (sequential mono- and dialkylation of resin-bound glycine via automated synthesis)
 RN 200573-87-1 CAPLUS
 CN Phenylalanine, N-(2-quinolinylcarbonyl)-, mono(trifluoroacetate) (9CI) (CA INDEX NAME)
 CM 1
 CRN 197392-59-9
 CMF C19 H16 N2 O3




CM 2
 CRN 76-05-1
 CMF C2 H F3 O2



L4 ANSWER 209 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)
 REFERENCE COUNT: 16 THERE ARE 16 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT


L4 ANSWER 210 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
 ACCESSION NUMBER: 1997:707354 CAPLUS
 DOCUMENT NUMBER: 127:307645
 TITLE: Solid-phase synthesis of unnatural amino acids using unactivated alkyl halides
 AUTHOR(S): O'Donnell, Martin J.; Lugar, Charles W.; Pottorf, Richard S.; Zhou, Changyou; Scott, William L.; Cwi, Cynthia L.
 CORPORATE SOURCE: Department of Chemistry, Indiana University-Purdue University at Indianapolis, Indianapolis, IN, 46202, USA
 SOURCE: Tetrahedron Letters (1997), 38(41), 7163-7166
 CODEN: TELEAY; ISSN: 0040-4039
 PUBLISHER: Elsevier
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 AB Conditions were developed for the efficient alkylation of the resin-bound benzophenone imine of glycine with a variety of unreactive alkyl halides. Alkylations were accomplished at room temperature in NMP using the phosphazene-type base BEMP.
 IT 197392-59-9P
 RL: SPN (Synthetic preparation); PREP (Preparation)
 (solid-phase synthesis of unnatural amino acids using unactivated alkyl halides)
 RN 197392-59-9 CAPLUS
 CN Phenylalanine, N-(2-quinolinylcarbonyl)- (9CI) (CA INDEX NAME)



L4 ANSWER 211 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)
 ACCESSION NUMBER: 1997:674193 CAPLUS
 DOCUMENT NUMBER: 127:355226
 TITLE: In vitro and in vivo characterization of NK3 receptors in the rabbit eye by use of selective non-peptide NK3 receptor antagonists
 AUTHOR(S): Medhurst, Andrew D.; Hay, Douglas W. P.; Parsons, Andrew A.; Martin, Lenox D.; Griswold, Don E.
 CORPORATE SOURCE: Department of Neurosciences Research, SmithKline Beecham Pharmaceuticals, Essex, CM19 5AW, UK
 SOURCE: British Journal of Pharmacology (1997), 122(3), 469-476
 PUBLISHER: Stockton
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 AB 1 Inhibition of NK3 receptor agonist-induced contraction in the rabbit isolated iris sphincter muscle was used to assess the in vitro functional activity of three 2-phenyl-4-quinolinecarboxamides, members of a novel class of potent and selective non-peptide NK3 receptor antagonists. In addition, an in vivo correlate of this in vitro response, namely NK3 receptor agonist-induced miosis in conscious rabbits, was characterized with some of these antagonists. 2 In vitro senktide (succinyl-[Asp⁹, MePhe⁸]-substance P (6-11) and [MePhe⁷]-neurokinin B([MePhe⁷]-NKB) were potent contractile agents in the rabbit iris sphincter muscle but exhibited quite different profiles. Senktide produced monophasic log concentration-effect curves with a mean pD₂=9.03±0.06 and mean nH=1.2±0.02 (n=14). In contrast, [MePhe⁷]-NKB produced shallow log concentration-effect curves which often appeared biphasic (nH=0.54±0.04, n=8), preventing the accurate determination of pD₂ values. 3 The contractile responses to the NK3 receptor agonist senktide were antagonized in a surmountable and concentration-dependent manner by SB 223412 [(1-)--(S)-N-(α-ethylbenzyl)-3-hydroxy-2-phenylquinoline-4-carboxamide; 3-30 nM, pA₂=8.4, slope=1.8±0.3, n=4], SB 222200 [(1-)--(S)-N-(α-ethylbenzyl)-3-methyl-2-phenylquinoline-4-carboxamide; 30-300 nM, pA₂=7.9, slope=1.4±0.06, n=4] and SB 218795 [(1-)--(R)-N-(α-methoxycarbonylbenzyl)-2-phenylquinoline-4-carboxamide; 0.3 and 3 μM apparent pK_B=7.4±0.06, n=6]. 4 Contractile responses to the NK3 receptor agonist [MePhe⁷]-NKB in the rabbit iris sphincter muscle were unaffected by SB 218795 (0.3 and 3 μM, n=8). In contrast, SB 223412 (30 and 300 μM, n=4) and SB 222200 (0.3 and 3 μM, n=4) inhibited responses to low concns. (<1 nM), to a greater extent than higher concns. (>1 nM) of [MePhe⁷]-NKB. Furthermore, log concentration-effect curves to [MePhe⁷]-NKB became steeper and monophasic in the presence of each antagonist. 5 SB 218795 (3 μM, n=4) had no effect on contractions induced by transmural nerve stimulation (2 Hz) or substance P, exemplifying the selectivity of this class of antagonist for functional NK3 receptors over NK1 receptors in the rabbit. 6 In vivo, senktide (1, 10 and 25 μg i.v., i.e. 1.2, 11.9 and 29.7 nmol, resp.) induced concentration-dependent bilateral miosis in conscious rabbits (maximum pupillary constriction=4.25±0.25 mm; basal pupillary diameter 7.75±0.48 mm; n=4). The onset of miosis was within 2-5 min of application of senktide and responses lasted up to 30 min. Responses to two i.v. administrations of 25 μg senktide given 30 min apart revealed no evidence of tachyphylaxis. Topical administration of atropine (1%) to the eye enhanced pupillary responses to 25 μg senktide. This was probably due to the mydriatic effect of atropine since it significantly

L4 ANSWER 211 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)
 increased baseline pupillary diam. from 7.0±0.4 mm to 9.0±0.7 mm (n=4), thereby increasing the max. capacity for miosis. Senktide-induced miosis was inhibited by SB 222200 (1 and 2 mg kg⁻¹, i.v., i.e. 2.63 and 5.26 μmol kg⁻¹; max. inhibition 100%; n=3-4), SB 223412 (0.5 and 1 mg kg⁻¹, i.v., i.e. 1.31 and 2.61 μmol kg⁻¹; max. inhibition 100%; n=3), SB 218795 (0.5 and 1 mg kg⁻¹, i.v., i.e. 1.26 and 2.52 μmol kg⁻¹; max. inhibition 78%; n=3), and the structurally distinct NK3 receptor antagonist SR 142801 [(S)-N-(1-{3-(1-benzoyl-3-(3,4-dichlorophenyl)piperidin-3-yl)propyl}-4-phenylpiperidin-4-yl)-N-methylacetamide; 1.5 mg kg⁻¹, i.v., i.e. 2.47 μmol kg⁻¹, max. inhibition 92%; n=3]. Optical administration of senktide (25 μg; 29.7 nmol) to the eye induced unilateral miosis in the treated eye only. At this dose there was no significant difference (P<0.05) between pupillary constriction obtained by topical or i.v. senktide, and topically administered atropine had no significant effect on responses to topical senktide (n=4). 8 [MePhe⁷]-NKB (125, 250 and 500 μg, i.v., i.e. 98.31, 196.62 and 393.24 nmol, resp.) also induced bilateral miosis in conscious rabbits (max. pupillary constriction=4.13±0.30 mm; n=4), but in contrast to in vitro studies this agonist was approx. 100 fold less potent than senktide. [MePhe⁷]-NKB-induced miosis was inhibited by SB 222200 (5 mg kg⁻¹, i.v., i.e. 13.14 μmol kg⁻¹; max. inhibition 69%; n=3). 9 In summary, SB 223412, SB 222200 and SB 218795 are potent and selective antagonists of NK3 receptor-mediated contraction in the rabbit isolated iris sphincter muscle. In addn., NK3 receptor agonist-induced miosis in conscious rabbits is a good in vivo correlate of the in vitro rabbit iris sphincter muscle prep. and appears to be a useful model for characterizing the pharmacodynamic profile and efficacy of structurally distinct NK3 receptor antagonists, such as SB 222200, SB 223412, SB 218795 and SR 142801.
 IT 174635-53-1, SB 218795
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); BUU (Biological use, unclassified); THU (Therapeutic use); BIOL (Biological study); USES (Uses)
 (characterization of NK3 receptors in the rabbit eye by use of selective non-peptide NK3 receptor antagonists)
 RN 174635-53-1 CAPLUS
 CN Benzeneacetic acid, α-[(2-phenyl-4-quinolinyl)carbonyl]amino]-, methyl ester, (αR)- (9CI) (CA INDEX NAME)

Absolute stereochemistry. Rotation (-).




L4 ANSWER 212 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
 ACCESSION NUMBER: 1997:594721 CAPLUS
 DOCUMENT NUMBER: 127:278064
 TITLE: Substituted 4-hydroxyphenylalkanoic acid derivatives with agonist activity to PPAR-gamma
 INVENTOR(S): Willson, Timothy Mark; Mook, Robert Anthony, Jr.; Kaldor, Istvan; Henke, Brad Richard; Deaton, David Norman; Collins, Jon Loren; Cobb, Jeffrey Edmond; et al.
 PATENT ASSIGNEE(S): Glaxo Group Ltd., UK
 SOURCE: PCT Int. Appl., 157 pp.
 CODEN: P1HX22
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 9731907	A1	19970904	WO 1997-EP916	19970226
W: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM				
RW: GH, KE, LS, MW, SD, SZ, UG, AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG				
CA 2247443	AA	19970904	CA 1997-2247443	19970226
AU 9720935	A1	19970916	AU 1997-20935	19970226
AU 717699	B2	20000330		
ZA 9701645	A	19971210	ZA 1997-1645	19970226
EP 888317	A1	19990107	EP 1997-906130	19970226
EP 888317	B1	20010912		
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI				
CN 1218460	A	19990602	CN 1997-193988	19970226
CN 1093124	B	20021023		
BR 9707786	A	19990727	BR 1997-7786	19970226
JP 2000507216	T2	20000613	JP 1997-530586	19970226
JP 3255930	B2	20020212		
NZ 331381	A	20000623	NZ 1997-331381	19970226
IL 125796	A1	20010614	IL 1997-125796	19970226
AT 205495	A	20010915	AT 1997-906130	19970226
ES 2163125	T3	20020116	ES 1997-906130	19970226
PT 888317	T	20020328	PT 1997-97906130	19970226
SK 282753	B6	20021203	SK 1998-1163	19970226
HR 970110	B1	20030630	HR 1997-970110	19970226
TW 391958	B	20000601	TW 1997-86102826	19970307
US 6294580	B1	20010925	US 1998-125750	19980825
NO 9803940	A	19981027	NO 1998-3940	19980827
HK 1015369	A1	20020215	HK 1999-100498	19990205
PRIORITY APPLN. INFO.:			GB 1996-4242	A 19960228
			WO 1997-EP916	W 19970226




OTHER SOURCE(S): MARPAT 127:278064
 AB Compds. 4-(A-B-O)C6H4-Q-CHZCO2R1 [A = (un)substituted Ph, heterocyclyl, fused bicyclic ring; B = alkylene, heterocyclyl; Q = alkylene; R1 = H, alkyl; Z = alkylenophenyl, NR3R4 (R3 = H, alkyl; R4 = YXOTR5, YCH(OH)TR5 with Y = bond, alkylene, alkenylene, cycloalkylene, etc. and T = bond, O, etc. and R5 = alkyl, cycloalkyl, (un)substituted Ph)] were prepared and their agonist activity to PPAR-gamma determined. E.g., O-benzyl L-tyrosine,

L4 ANSWER 212 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)
 dicyclohexylamine, and 1-benzoylacetone were refluxed in MeOH to give 3-(4-benzoyloxyphenyl)-2(S)-(1-methyl-3-oxo-3-phenylpropenylamino)propionic acid dicyclohexylamine salt.
 IT 196808-22-79
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)
 (preparation of (hydroxyphenyl)alkanoic acids with agonist activity to PPAR-gamma)
 RN 196808-22-7 CAPLUS
 CN L-Tyrosine, N-[(4-oxo-4H-1-benzopyran-3-yl)carbonyl]-O-(phenylmethyl)-, methyl ester (9CI) (CA INDEX NAME)

Absolute stereochemistry.



L4 ANSWER 213 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
 ACCESSION NUMBER: 1997:520078 CAPLUS
 DOCUMENT NUMBER: 127:205826
 TITLE: Design, synthesis and pharmacological test of a quinoline based, nonpeptidic analog of neurotensin(8-13)
 AUTHOR(S): Hong, Feng; Pang, Yuan-Ping; Cusack, Bernadette; Richelson, Elliott
 CORPORATE SOURCE: Neurochemistry Research, Mayo Foundation for Medical Education and Research, Jacksonville, FL, 32224, USA
 SOURCE: Journal of the Chemical Society, Perkin Transactions 1: Organic and Bio-Organic Chemistry (1997), (14), 2083-2088
 CODEN: JCPRB4; ISSN: 0300-922X
 PUBLISHER: Royal Society of Chemistry
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 GI




AB Based on the multiple template approach to developing nonpeptidic mimetics of neuropeptides, the design, synthesis and pharmacol. testing of a quinoline based analog of neurotensin(8-13) I (Ad = 1-adamantyl) are reported. The newly synthesized quinoline analog is found to be less active in binding to the neurotensin receptors than previously reported mimetics II and III, which are partial nonpeptidic analogs of neurotensin(8-13). The correct structures of II and III are reported,

L4 ANSWER 213 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)
 here, since this study led to the discovery of a mistake in the previously reported literature procedure for alkylation at position-3 of Et indole-2-carboxylate.

IT 194673-23-9P
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation)
 (synthesis and pharmacol. testing of a quinoline based, nonpeptidic analog of neurotensin(8-13))
 RN 194673-23-9 CAPLUS
 CN 2-Quinolinecarboxamide, 4,8-bis[(5-aminopentyl)oxyl]-N-[1-[(4-hydroxyphenyl)methyl]-2-oxo-2-(tricyclo[3.3.1.1.3,7]dec-1-ylamino)ethyl]-, (S)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



REFERENCE COUNT: 13 THERE ARE 13 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 214 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
 ACCESSION NUMBER: 1997:475133 CAPLUS
 DOCUMENT NUMBER: 127:162123
 TITLE: Peptides having bradykinin antagonist action
 INVENTOR(S): Henke, Stephan; Anagnostopoulos, Hristo; Breipohl, Gerhard; Knolle, Jochen; Stechl, Jens; Scholten, Bernward; et al.
 PATENT ASSIGNEE(S): Hoechst A.-G., Germany
 SOURCE: U.S., 26 pp., Cont. of U.S. Ser. No. 236,018.
 CODEN: USXXAM
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 4
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 5648333	A	19970715	US 1995-487442	19950607
DD 284030	A5	19901031	DD 1989-331416	19890802
ZA 8906068	A	19910130	ZA 1989-6068	19890809
DE 3926822	A1	19910221	DE 1989-3926822	19890814
DE 4013270	A1	19911031	DE 1990-4013270	19900426
RU 2083586	C1	19970710	RU 1992-5052703	19921012
LT 3375	B	19950825	LT 1993-717	19930625
PRIORITY APPLN. INFO.:			DE 1988-3839581	A 19881124
			DE 1989-3916291	A 19890519
			DE 1989-3926225	A 19890603
			US 1989-374162	B2 19890630
			DE 1989-3926822	A 19890814
			DE 1990-4013270	A 19900426
			US 1990-565270	B2 19900810
			US 1991-690297	B1 19910424
			US 1991-746149	B1 19910814
			US 1992-837090	B2 19920218
			US 1992-841766	B1 19920302
			US 1992-969523	B2 19921030
			US 1992-982052	B2 19921125
			US 1993-12849	B1 19930203
			US 1994-236018	A1 19940502


OTHER SOURCE(S): MARPAT 127:162123
 AB Peptides A-B-C-E-F-K-P-G-M-F [A = H, alkyl, alkanoyl, cycloalkyl, aryl, etc.; B = basic amino acid which may be substituted in side chain; C = G'-G'-Gly or G'-NH(CH2)nCO, where G' = heterocyclylcarbonyl and n = 2-8; E = aromatic amino acid radical; F, M = bond or amino acid which may be substituted in side chain; K = bond or NH(CH2)xCO, where x = 1-4; P = D-Tic (Tic = 1,2,3,4-tetrahydroisoquinolin-3-ylcarbonyl); G = bond or G'] were prepared as bradykinin antagonists. Thus, H-D-Arg-Arg-Hyp-Pro-Gly-Phe-Ser-D-Tic-Phe-Arg-OH was prepared by the solid phase method and assayed for bradykinin antagonist activity (IC50 = 4.6 x 10-6 M).

IT 193618-41-6P
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)
 (peptides having bradykinin antagonist action)
 RN 193618-41-6 CAPLUS
 CN L-Arginine, D-arginyl-L-arginyl-(4R)-4-hydroxy-L-prolyl-L-prolylglycyl-L-phenylalanyl-L-glutamyl-(3R)-1,2,3,4-tetrahydro-3-isoquinolinecarbonyl-L-phenylalanyl- (9CI) (CA INDEX NAME)


Absolute stereochemistry.

L4 ANSWER 214 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)

PAGE 1-A



PAGE 1-B




L4 ANSWER 215 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
 ACCESSION NUMBER: 1997:465088 CAPLUS
 DOCUMENT NUMBER: 127:95204
 TITLE: Preparation of quinoline-4-carboxamides and their use as neurokinin-3 and neurokinin-2 receptor antagonists
 INVENTOR(S): Giardina, Giuseppe Arnaldo Maria; Grugni, Mario; Raveglia, Luca Francesco; Farina, Carlo
 PATENT ASSIGNEE(S): Smithkline Beecham S.P.A., Italy; Giardina, Giuseppe Arnaldo Maria; Grugni, Mario; Raveglia, Luca Francesco; Farina, Carlo
 SOURCE: PCT Int. Appl., 79 pp.
 DOCUMENT TYPE: CODEN: PIXXD2
 LANGUAGE: Patent
 FAMILY ACC. NUM. COUNT: 1 English
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 9719926	A1	19970605	WO 1996-EP5207	19961122
W:	AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, US, UZ, VN, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM			
RW:	KE, LS, MW, SD, SZ, UG, AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG			
IT 1307330	B1	20011030	IT 1996-M11688	19960802
CA 2230328	AA	19970605	CA 1996-2238328	19961122
AU 9710318	A1	19970619	AU 1997-10318	19961122
ZA 9609811	A	19960522	ZA 1996-9811	19961122
CN 1207729	A	19990210	CN 1996-199747	19961122
BR 9611757	A	19990406	BR 1996-11757	19961122
EP 1019377	A1	20000719	EP 1996-941025	19961122
R:	AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, FI, RO			
JP 200051325	T2	20001010	JP 1997-520158	19961122
TW 409123	B	20001021	TW 1996-85114501	19961123
NO 9802333	A	19980722	NO 1998-2333	19980522
US 2000268827	A1	20020606	US 2001-994402	20011126
PRIORITY APPL. INFO.:			IT 1995-M12462	A 19951124
			IT 1996-M11688	A 19960802
			WO 1996-EP5207	W 19961122
			US 1998-77262	B1 19980806
			US 2000-515336	B1 20000605

OTHER SOURCE(S): MARPAT 127:95204
 GI


L4 ANSWER 215 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)




AB The title compds. [I; A = (un)substituted aryl, C5-7 cycloalkdienyl, (un)substituted single or fused ring aromatic heterocyclyl; R = (un)substituted C1-6 alkyl, C3-7 cycloalkyl, C3-7 cycloalkylalkyl, (un)substituted Ph, an optionally substituted five-membered heteroarom. ring, etc.; R1 = hydrogen or up to four substituents selected from C1-6 alkyl, C1-6 alkenyl, aryl, C1-6 alkoxy, hydroxy, halogen, nitro, cyano, carboxy, carboxamido, sulfonamido, C1-6 alkoxy, trifluoromethyl, alkoxy, phthalimido, (un)substituted amino, etc.; R2 = hydrogen, C1-6 alkyl, hydroxy, halogen, cyano, (un)substituted amino, etc.; R3 = C1-6 alkyl, C3-7 cycloalkyl, C4-7 cycloalkylalkyl, (un)substituted aryl, (un)substituted single or fused ring aromatic heterocyclyl; R4 = hydrogen, C1-6 alkyl, useful as neurokinin 3 and neurokinin 2 receptor antagonists, are prepared. Thus, (S)-N-(α -ethylbenzyl)-3-(2-aminoethoxy)-2-phenylquinoline-4-carboxamide was reacted with α,α' -dibromo-o-xylene and sulfated with HCl, producing (S)-N-(α -ethylbenzyl)-3-(2-(2-isindolyl)ethoxy)-2-phenylquinoline-4-carboxamide dihydrochloride (m.p. 95°; decomposition) which demonstrated a binding affinity in human neurokinin-3 receptors (expressed in CHO cell lines) against [125I]-[Me-Ph7]-neurokinin B of 1.2 nM.

IT 191796-50-6P
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)
 (preparation of quinoline-4-carboxamides and their use as neurokinin-3 and neurokinin-2 receptor antagonists)

RN 191796-50-6 CAPLUS
 CN Benzenecarboxylic acid, α -methyl- α -[[(2-phenyl-4-quinolinyl)carbonylamino]-, methyl ester (9CI) (CA INDEX NAME)




L4 ANSWER 216 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)
 (prepn. of quinoline-deriv. NK3 receptor antagonists)
 RN 174635-51-9 CAPLUS
 CN Benzenecarboxylic acid, α -[[(2-phenyl-4-quinolinyl)carbonylamino]-, methyl ester (9CI) (CA INDEX NAME)



L4 ANSWER 216 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
 ACCESSION NUMBER: 1997:461591 CAPLUS
 DOCUMENT NUMBER: 127:95205
 TITLE: Preparation of quinoline-derivative NK3 receptor antagonists
 INVENTOR(S): Giardina, Giuseppe Arnaldo Maria; Farina, Carlo; Grugni, Mario; Raveglia, Luca Francesco
 PATENT ASSIGNEE(S): Smithkline Beecham S.P.A., Italy; Giardina, Giuseppe Arnaldo Maria; Farina, Carlo; Grugni, Mario; Raveglia, Luca Francesco
 SOURCE: PCT Int. Appl., 101 pp.
 DOCUMENT TYPE: CODEN: PIXXD2
 LANGUAGE: Patent
 FAMILY ACC. NUM. COUNT: 1 English
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 9719927	A1	19970605	WO 1996-EP5209	19961122
W:	JP, US			
RW:	AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE			
EP 874827	A1	19981104	EP 1996-939926	19961122
EP 874827	B1	20030521		
R:	BE, CH, DE, ES, FR, GB, IT, LI, NL			
JP 2000512614	T2	20000926	JP 1997-520159	19961122
US 2003195204	A1	20031016	US 2002-140452	20020507
PRIORITY APPL. INFO.:			GB 1995-24104	A 19951124
			WO 1996-EP5209	W 19961122
			US 1998-77156	B1 19980521

OTHER SOURCE(S): MARPAT 127:95205
 GI



AB The title compds. [I; A = (un)substituted Ph, (un)substituted naphthyl, C5-7 cycloalkdienyl, (un)substituted heterocyclyl; R = C1-8 alkyl, C3-7 cycloalkyl, C4-7 cycloalkylalkyl, (un)substituted Ph, phenylalkyl, etc.; R1, R2 = H, C1-6 alkyl, or together form a (CH2)n, etc.; n = 3-5; R3, R4 = H, C1-6 alkyl, C1-6 alkenyl, aryl, C1-6 alkoxy, hydroxy, halogen, nitro, cyano, carboxy, carboxamido, sulfonamido, C1-6 alkoxy, trifluoromethyl, acyloxy, phthalimido, (un)substituted amino, etc.; R5 = C1-6 alkyl, C3-7 cycloalkyl, etc.; X = O, S, NC, tpbond, N, etc.; which are NK3 receptor antagonists (no data), are prepared. Thus, α -methylbenzylamine was amidated with 2-phenylquinoline-4-carbonyl chloride, producing N-(α -methylbenzyl)-2-phenyl-4-quinolinecarboxamide, m.p. 156-157°.

IT 174635-51-9P
 RL: SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)

L4 ANSWER 217 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
 ACCESSION NUMBER: 1997:348323 CAPLUS
 DOCUMENT NUMBER: 127:81757
 TITLE: The solid phase synthesis of α,α -disubstituted unnatural amino acids and peptides (di-UPS)
 AUTHOR(S): Scott, William L.; Zhou, Changyou; Fang, Zhiqiang; O'Donnell, Martin J.
 CORPORATE SOURCE: Res. Technologies Proteins, Lilly Res. Labs., Indianapolis, IN, 46285, USA
 SOURCE: Tetrahedron Letters (1997), 38(21), 3695-3698
 CODEN: TELEAY; ISSN: 0040-4039
 PUBLISHER: Elsevier
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 OTHER SOURCE(S): CASREACT 127:81757


AB This paper reports a new, mild procedure (di-UPS) for the solid phase synthesis of racemic α,α -disubstituted amino acids and epimeric α,α -disubstituted terminal amino acid residues in a resin-bound peptide. The synthetic route is compatible with most protected amino acid side chains and can be used in a continuing solid phase synthesis. Di-UPS should find a wide applicability in the design and solid phase synthesis of hybrid amino acids and peptides and the construction of basis units for combinational chemical

IT 191675-94-2P

RL: SPN (Synthetic preparation); PREP (Preparation)
 (solid phase synthesis of disubstituted unnatural amino acids and peptides)

RN 191675-94-2 CAPLUS

CN Phenylalanine, α -methyl-N-(2-quinolinylcarbonyl)- (9CI) (CA INDEX NAME)



REFERENCE COUNT: 43 THERE ARE 43 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 218 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
 ACCESSION NUMBER: 1997:320920 CAPLUS
 DOCUMENT NUMBER: 126:338400
 TITLE: Discovery of a Novel Class of Selective Non-Peptide Antagonists for the Human Neurokinin-3 Receptor. 1. Identification of the 4-Quinolonecarboxamide Framework
 AUTHOR(S): Giardina, Giuseppe A. M.; Sarau, Henry M.; Farina, Carlo; Medhurst, Andrew D.; Grugni, Mario; Raveglia, Luca F.; Schmidt, Dulcie B.; Rigolio, Roberto; Luttmann, Mark; Vecchiotti, Vittorio; Hay, Douglas W. P.
 CORPORATE SOURCE: Department of Chemistry, SmithKline Beecham S.p.A., Baranzate, 20021, Italy
 SOURCE: Journal of Medicinal Chemistry (1997), 40(12), 1794-1807
 CODEN: JMCMAR; ISSN: 0022-2623
 PUBLISHER: American Chemical Society
 DOCUMENT TYPE: Journal
 LANGUAGE: English

AB A novel class of potent and selective non-peptide neurokinin-3 (NK-3) receptor antagonists, featuring the 4-quinolonecarboxamide framework, was designed based upon chemical diverse NK-1 receptor antagonists. The novel compounds, prompted by chemical modifications of the prototype, were characterized by binding anal. using a membrane preparation of chinese hamster ovary (CHO) cells expressing the human neurokinin-3 receptors (hNK-3-CHO), and clear structure-activity relationships (SARs) were established. From SARs, (R)-N-[(4-methoxycarbonyl)benzyl]-2-phenylquinoline-4-carboxamide (I, SB 218795, hNK-3-CHO binding $K_i = 13$ nM) emerged as one of the most potent compounds of this novel class. Selectivity studies vs. the other neurokinin receptors (hNK-2-CHO and hNK-1-CHO) revealed that 65 is about 90-fold selective for hNK-3 vs. hNK-2 receptors (hNK-2-CHO binding $K_i = 1221$ nM) and over 7000-fold selective vs. hNK-1 receptors (hNK-1-CHO binding $K_i = >100$ μ M). In vitro functional studies in rabbit isolated iris sphincter muscle preparation demonstrated that I is a competitive antagonist of the contractile response induced by the potent and selective NK-3 receptor agonist senktide with a $K_B = 43$ nM. Overall, the data indicate that I is a potent and selective hNK-3 receptor antagonist and a useful lead for further chemical optimization.

IT 174635-51-9P


RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation)

(preparation of quinolonecarboxamide-containing nonpeptide neurokinin-3 receptor antagonists)


RN 174635-51-9 CAPLUS

CN Benzeneacetic acid, α -[[(2-phenyl-4-quinolinyl)carbonyl]amino]-, methyl ester (9CI) (CA INDEX NAME)

L4 ANSWER 218 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)



L4 ANSWER 219 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
 ACCESSION NUMBER: 1997:316770 CAPLUS
 DOCUMENT NUMBER: 127:47593
 TITLE: Susceptibility of Pseudomonas aeruginosa of various pyocin types to the newly synthesized ampicillin derivatives, N-(6,7-difluoroquinolonyl)ampicillin
 AUTHOR(S): Chen, C. H.; Tsou, T. L.; Chiang, H. Y.; Lee, S. H.; Lee, F.; Lee, J. H.; Wang, T. M.; Liu, Y. T.
 CORPORATE SOURCE: Section of Bacteriology, Division of Clinical Pathology, National Defence Medical Center, Tri-Service General Hospital, Taipei, Taiwan
 SOURCE: Journal of Antimicrobial Chemotherapy (1997), 39(3), 325-330
 CODEN: JACHDX; ISSN: 0305-7453
 PUBLISHER: Oxford University Press
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 GI



AB Six hundred and thirty-two isolates of Pseudomonas aeruginosa of 17 pyocin types were collected in 1993 in Taiwan. Types 1, 10, 3, 35 and 12 were the most common pyocin types identified in Taiwan with isolation frequencies of 47.3%, 24.4%, 7.6%, 3.6% and 2.2%, resp. Several pyocin subtypes were determined. All pyocin types (one isolate of each tested) were resistant to ampicillin and nalidixic acid, but sensitive to fluoroquinolone antibiotics, such as norfloxacin and enoxacin, indicating that cross-resistance to quinolone antibiotics of nalidixic acid and fluoroquinolone derivs. has not developed. A new ampicillin derivative of 6,7-difluoroquinolonic acid, N-(6,7-difluoroquinolonyl)-ampicillin (AU-1, I), was synthesized by coupling ampicillin with 6,7-difluoroquinolonic acid (FP-3). I was much more active than either ampicillin or FP-3 alone against all pyocin types of P. aeruginosa and induced filamentation in most growing cells.

IT 190902-80-8P


RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); PRP (Properties); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)

(susceptibility of Pseudomonas aeruginosa of various pyocin types to (difluoroquinolonyl)ampicillin)

RN 190902-80-8 CAPLUS


CN 4-Thia-1-azabicyclo[3.2.0]heptane-2-carboxylic acid, 6-[[[(2R)-[(6,7-difluoro-1,4-dihydro-4-oxo-3-quinolinyl)carbonyl]amino]phenylacetyl]amino]-3,3-dimethyl-7-oxo-, (2S,5R,6R)- (9CI) (CA INDEX NAME)

L4 ANSWER 219 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)
Absolute stereochemistry.




L4 ANSWER 220 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
ACCESSION NUMBER: 1997:290042 CAPLUS
DOCUMENT NUMBER: 126:293586
TITLE: Ferrocenyl Amino Acids: A Synthetic and Structural Study
AUTHOR(S): Kraatz, Heinz-Bernhard; Lusztyk, Janusz; Enright, Gary D.
CORPORATE SOURCE: Supramolecular Chemistry and Biology Group Steacie Institute of Molecular Science, National Research Council of Canada, Ottawa, ON, K1A 0R6, Can.
SOURCE: Inorganic Chemistry (1997), 36(11), 2400-2405
CODEN: INOCAL; ISSN: 0020-1669
PUBLISHER: American Chemical Society
DOCUMENT TYPE: Journal
LANGUAGE: English
OTHER SOURCE(S): CASREACT 126:293586
AB Ester-protected amino acids were coupled to ferrocenecarboxylic acid using the DCC/HOBt protocol to give ferrocenyl N-amino acids (amino acid = Glu(OBz)2 (2a), Gly(OEt) (2b), Pro(OBz) (2c), Cys(SBz)OMe (2d), Ala(OBz) (2e), Tyr(OBz) (2f), Phe(OBz) (2g)). All products were fully characterized. The intermediate hydroxybenzotriazole active ester FcCOOBt (3) was isolated and fully characterized. The solid state structures of 2a, 2d, and 3 were determined by single-crystal x-ray diffraction. 2A: monoclinic space group P21 with a 11.8142(5), b 9.7560(5), c 22.9456(10) Å, β 90.246(5)°, Z = 2, R = 0.046. 2D: orthorhombic space group P212121 with a 9.957(2), b 11.680(2), c 36.452(2) Å, Z = 4, R = 0.065. The solid state structures of 2a and 2d show extensive C...H-N-H bonding. 3: Triclinic space group P-hivin.1 with a 7.0391(5), b 10.7922(7), c 11.1690(7) Å, α 108.071(5), β 107.957(5), γ 103.896(5)°, Z = 2, R = 0.030. The long ester bond distance of 1.427(2) Å provides a rationale for its inherent reactivity toward primary and secondary amines. Some of the ester-protected ferrocenyl amino acids were also studied by cyclic voltammetry.
IT 189116-56-1P
RL: SPN (Synthetic preparation); PREP (Preparation) (preparation of)
RN 189116-56-1 CAPLUS
CN Ferrocene, ([1-[(4-hydroxyphenyl)methyl]-2-oxo-2-(phenylmethoxy)ethyl]amino]carbonyl)-, (S)- (9CI) (CA INDEX NAME)

L4 ANSWER 220 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)




PAGE 1-A



PAGE 2-A

L4 ANSWER 221 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
ACCESSION NUMBER: 1997:287115 CAPLUS
DOCUMENT NUMBER: 127:30759
TITLE: Potency comparison of peptidomimetic inhibitors against HIV-1 and HIV-2 proteinases: design of equipotent lead compounds
AUTHOR(S): Weber, Jan; Majer, Pavel; Litera, Jaroslav; Urban, Jan; Soucek, Milan; Vondrasek, Jiri; Konvalinka, Jan; Novek, Petr; Sedlacek, Juraj; Strop, Petr; Krausslich, Hans-Georg; Pichova, Iva
CORPORATE SOURCE: Dep. Biochem., Inst. Org. Chem. Biochem., Acad. Sci. Czech Republic, Prague, 166 10, Czech Rep.
SOURCE: Archives of Biochemistry and Biophysics (1997), 341(1), 62-69
CODEN: ABBIA4; ISSN: 0003-9861
PUBLISHER: Academic
DOCUMENT TYPE: Journal
LANGUAGE: English
AB HIV-1 and HIV-2 proteinases (PR) are responsible for the processing of viral polyproteins, a step that is crucial for the formation of infectious virus particles. PR represents one of the most important targets for antiviral chemotherapy. Inhibitors of HIV-1 PR usually exhibit a 10- to 100-fold weaker affinity for HIV-2 PR. To design subnanomolar inhibitors for both HIV-1 and HIV-2 PRs, we prepared a series of compds. varying in the type of scissile bond replacement as well as in the P1, P1', and P2' side chains. While inhibitors containing reduced amide, hydroxyethylamine and statine isosteres had Ki values in the range of 10-10-10-9 M against HIV-1 PR, their activities against HIV-2 PR were several orders of magnitude lower. Glutamic acid was identified to be the optimal P2' residue for both PRs. HIV-2 PR was shown to be more sensitive to P2' Glu-OH replacement. Using this data set we were able to design and prepare hydroxyethylene isostere containing inhibitors that were equipotent against both PRs.
IT 190667-94-8
RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); PRP (Properties); BIOL (Biological study) (potency comparison of peptidomimetic inhibitors against HIV-1 and HIV-2 proteinases)
RN 190667-94-8 CAPLUS
CN L-Phenylalaninamide, (3S)-2-[(1,1-dimethylethoxy)carbonyl]-1,2,3,4-tetrahydro-3-isoquinolinecarbonyl-L-phenylalanyl-L-α-glutamyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



L4 ANSWER 222 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
 ACCESSION NUMBER: 1997:283758 CAPLUS
 DOCUMENT NUMBER: 126:264364
 TITLE: Acylated oligopeptide derivatives having cell signal inhibiting activity
 INVENTOR(S): Garcia-Echeverria, Carlos; Gay, Brigitte; Furet, Pascal; Rahuel, Joseph; Caravatti, Giorgio; Pretz, Heinz; Schoepfer, Joseph
 PATENT ASSIGNEE(S): Ciba-Geigy A.-G., Switz.
 SOURCE: PCT Int. Appl., 257 pp.
 CODEN: PIXXD2
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 9708193	A1	19970306	WO 1996-EP3473	19960806
W:	AL, AU, BB, BG, BR, CA, CN, CU, CZ, EE, GE, HU, IL, IS, JP, KP, KR, LK, LR, LT, LV, MG, MK, MN, MX, NO, NZ, PL, RO, SG, SI, SK, TR, TT, UA, US, UZ, VN, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM			
RW:	KE, LS, MW, SD, SZ, UG, AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG			
AU 9667425	A1	19970319	AU 1996-67425	19960806
EP 846127	A1	19980610	EP 1996-927694	19960806
R:	AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, FI			
ZA 9606967	A	19970217	ZA 1996-6967	19960816
PRIORITY APPLN. INFO.:			GB 1995-17060	A 19950817
			WO 1996-EP3473	W 19960806


OTHER SOURCE(S): MARPAT 126:264364
 AB Peptides X-PTI (AA)-Y (AA = natural or unnatural amino acid residue, n = 0-15, PTI = tyrosine or preferably phosphotyrosine or phosphotyrosine mimic, X = arylcarbonyl, cycloalkylcarbonyl, tricycloalkylcarbonyl, arylsulfonyl, etc., Y = OH, C-terminal protecting group, amino group) or their salts were prepared for the treatment of diseases that respond to inhibition of the interaction of a protein comprising an SH2 domain and a protein tyrosine. Thus, 3-aminobenzoyloxycarbonyl-Tyr(PO3H2)-Ile-Asn-Gln-NH2 trifluoroacetate salt was prepared by the solid phase method and had an IC50 value of 0.1 μ M in a test system using the phosphorylated "tail" EGFR-MBP fusion protein as ligand. Formulations containing acylated oligopeptides are described.

IT 188749-93-1P
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)
 (preparation of acylated oligopeptide derivs. having cell signal inhibiting activity)
 RN 188749-93-1 CAPLUS
 CN L-Glutamamide, O-phosphono-N-(6-quinolinylcarbonyl)-L-tyrosyl-L-isoleucyl-L-asparaginyl- (9CI) (CA INDEX NAME)


Absolute stereochemistry.

L4 ANSWER 223 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
 ACCESSION NUMBER: 1997:277558 CAPLUS
 DOCUMENT NUMBER: 126:289228
 TITLE: Reduction of ochratoxin A toxicity by heat-induced epimerization. In vitro effects of ochratoxins on embryonic chick meningeal and other cell cultures
 AUTHOR(S): Bruinink, A.; Rasonyi, T.; Sidler, C.
 CORPORATE SOURCE: Inst. Toxicol., Swiss Fed. Inst. Technol., Univ. Zuerich, Schwerzenbach, CH-8603, Switz.
 SOURCE: Toxicology (1997), 118(2,3), 205-210
 CODEN: TXCYAC; ISSN: 0300-483X
 PUBLISHER: Elsevier
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 AB The present study was designed to determine the toxic potential of three structurally related ochratoxins: ochratoxin A (OTA), ochratoxin B (OTB) and the heat-induced 3S-epimer of OTA (3S-OTA) recently discovered in roasted coffee and human serum. The toxicity was determined using serum-free cell cultures of embryonic chick meningeal fibroblasts, taking the effects on mitochondrial and lysosomal activity and culture protein content as an index for toxicity. OTA, OTB and 3S-OTA were toxic. However, the concentration necessary to induce comparable effects were nearly 19- and 10-fold higher for OTB and 3S-OTA, resp., than those for OTA. In a next step and sensitivity of serum-free cell cultures of embryonic chick neural retina and brain were compared in relation to meningeal cell cultures. In the present study, no indications for differences in sensitivity could be detected. Furthermore, our study suggest that the OTA-induced toxic effects are not due to the inhibition by OTA of phenylalanine-tRNA synthetase.
 IT 189152-21-4
 RL: ADV (Adverse effect, including toxicity); BIOL (Biological study) (ochratoxin A toxicity in relation to structure)
 RN 189152-21-4 CAPLUS
 CN L-Phenylalanine, N-[[[3S]-5-chloro-3,4-dihydro-8-hydroxy-3-methyl-1-oxo-1H-2-benzopyran-7-yl]carbonyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



L4 ANSWER 222 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)




L4 ANSWER 224 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
 ACCESSION NUMBER: 1997:207658 CAPLUS
 DOCUMENT NUMBER: 126:199840
 TITLE: Preparation of peptide derivatives as cell adhesion inhibitors
 INVENTOR(S): Lin, Ko-Chung; Adams, Steven P.; Castro, Alfredo C.; Zimmerman, Craig N.; Cuervo, Julio Hernan; Lee, Wen-Cherng; Hammond, Charles E.; Carter, Mary Beth; Almqvist, Ronald G.; Ensinger, Carol Lee
 PATENT ASSIGNEE(S): Biogen, Inc., USA; Lin, Ko-Chung; Adams, Steven, P.; Castro, Alfredo, C.; Zimmerman, Craig, N.; Cuervo, Julio, Hernan; Lee, Wen-Cherng; Hammond, Charles, E.; Carter, Mary, Beth; et al.
 SOURCE: PCT Int. Appl., 117 pp.
 CODEN: PIXXD2
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 9703094	A1	19970130	WO 1996-US11570	19960711
W:	AL, AM, AT, AU, AZ, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG			
RW:	KE, LS, MW, SD, SZ, UG, AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA			
US 6248713	B1	20010619	US 1995-498237	19950711
CA 2226868	AA	19970130	CA 1996-2226868	19960711
AU 9664894	A1	19970210	AU 1996-64894	19960711
AU 716276	B2	20000224		
EP 842196	A1	19980520	EP 1996-924444	19960711
R:	AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI			
CN 1193325	A	19980915	CN 1996-196380	19960711
BR 9609782	A	19990309	BR 1996-9782	19960711
JP 11511124	T2	19990928	JP 1996-505989	19960711
NZ 312950	A	20000128	NZ 1996-312950	19960711
EE 3694	B1	20020415	EE 1997-362	19960711
EE 200200384	A	20021015	EE 2002-200200384	19960711
FI 9800033	A	19980305	FI 1998-33	19980109
NO 9800097	A	19980311	NO 1998-97	19980109
BG 63876	B1	20030430	BG 1998-102241	19980210
US 6239108	B1	20010529	US 1998-983391	19980810
US 6596687	B1	20030722	US 2000-482296	20000112
AU 758886	B2	20030403	AU 2000-36445	20000525
PRIORITY APPLN. INFO.:			US 1995-498237	A 19950711
			AU 1996-64894	A3 19960711
			WO 1996-US11570	W 19960711


OTHER SOURCE(S): MARPAT 126:199840
 AB The present invention relates to novel peptide derivs. that are useful for inhibition and prevention of cell adhesion and cell adhesion-mediated pathologies. This invention also relates to pharmaceutical formulations comprising these compds. and methods of using them for inhibition and prevention of cell adhesion and cell adhesion-mediated pathologies. The compds. and pharmaceutical composition of this invention can be used as therapeutic or prophylactic agents. They are particularly well-suited for treatment of many inflammatory and autoimmune diseases. Thus, coupling of 4-(2-Mec6H4NHCONH)C6H4CH2CO2H (preparation given) with protected peptide

L4 ANSWER 224 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)
 H-Leu-Asp(OCH₂Ph)-Val-OCH₂Ph (prepn. given), followed by catalytic hydrogenolysis, gave cell adhesion inhibitor peptide 4-(2-MeC₆H₄NHCOH)C₆H₄CH₂CO-Leu-Asp-Val-OH (I). All 408 prepd. peptide derivs., including I, inhibited VLA4-dependent adhesion to a bovine serum albumin conjugate with H-Cys-Tyr-Asp-Glu-Leu-Pro-Gln-Leu-Val-Thr-Leu-Pro-His-Pro-Arg-Leu-His-Gly-Pro-Glu-Ile-Leu-Asp-Val-Pro-Ser-Thr-OH, with IC₅₀ values of <1 mM.
 IT 187734-70-9P
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)
 (preparation of peptide derivs. as cell adhesion inhibitors)
 RN 187734-70-9 CAPLUS
 CN L-Valine, N-(3-isoquinolinylcarbonyl)-L-phenylalanyl-L-α-aspartyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




L4 ANSWER 225 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
 ACCESSION NUMBER: 1997:153973 CAPLUS
 DOCUMENT NUMBER: 126:215990
 TITLE: Transformation of the mycotoxin ochratoxin A in plants. 1. Isolation and identification of metabolites formed in cell suspension cultures of wheat and maize
 Ruhland, Monika; Engelhardt, Gabriele; Schaefer, Wolfram; Wallnoefer, Peter R.
 CORPORATE SOURCE: Bayerische Landesanstalt für Ernährung, Abteilung Ernährung, München, 80638, Germany
 SOURCE: Natural Toxins (1996), 4(6), 254-260
 CODEN: NATOEE; ISSN: 1056-9014
 PUBLISHER: Wiley-Liss
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 AB The metabolism of the mycotoxin ochratoxin A in plant cells was investigated by using cell suspension cultures of wheat and maize. A number of metabolites were detected by HPLC-chromatog. with fluorescence detection. The main metabolites were ochratoxin α, ochratoxin A Me ester, two isomers of hydroxyochratoxin A, and the glucosides and Me esters of both hydroxyochratoxin A isomers. The compds. were isolated by TLC and preparative HPLC and identified by mass spectrometry and specific enzymic reactions.
 IT 188348-37-0
 RL: BSU (Biological study, unclassified); MFM (Metabolic formation); BIOL (Biological study); FORM (Formation, nonpreparative)
 (from transformation of ochratoxin A by suspension cultures of wheat and maize)
 RN 188348-37-0 CAPLUS
 CN L-Phenylalanine, N-[[[(3R,4S)-5-chloro-3,4-dihydro-4,8-dihydroxy-3-methyl-1-oxo-1H-2-benzopyran-7-yl]carbonyl]-, methyl ester (9CI) (CA INDEX NAME)
 Absolute stereochemistry.




REFERENCE COUNT: 23 THERE ARE 23 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 226 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
 ACCESSION NUMBER: 1997:114532 CAPLUS
 DOCUMENT NUMBER: 126:225198
 TITLE: Combinatorial synthesis of heterocycles: solid phase synthesis of 2-arylquinoline-4-carboxylic acid derivatives
 Gopalsamy, Ariamala; Pallai, Peter V.
 CORPORATE SOURCE: Department of Rational Drug Design, Procept, Inc., Cambridge, MA, 02139, USA
 SOURCE: Tetrahedron Letters (1997), 38(6), 907-910
 CODEN: TELEAY; ISSN: 0040-4039
 PUBLISHER: Elsevier
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 OTHER SOURCE(S): CASREACT 126:225198
 GI



AB The Doebner quinoline synthesis has been adapted to solid phase. Acylation of an amino acid coupled to the Rink polystyrene resin with pyruvyl chloride afforded the immobilized amide. Further reaction of the amide with the preformed Schiff's base R1C₆H₄N:CHC₆H₄R₂ (R₁ = 3-MeO, R₂ = 4-NO₂, 4-cyano; R₁ = H, R₂ = 4-NO₂) or aldehyde R₂C₆H₄CHO and aniline R1C₆H₄NH₂ gave, after trifluoroacetic acid cleavage, 2-arylquinoline-4-carboxylic acid amides I (R₃ = H, CH₂Ph, Me) in good yields.
 IT 188367-36-4P
 RL: SPN (Synthetic preparation); PREP (Preparation)
 (solid phase synthesis of arylquinolinecarboxylic acid derivs.)
 RN 188367-36-4 CAPLUS
 CN 4-Quinolinecarboxamide, N-[2-amino-2-oxo-1-(phenylmethyl)ethyl]-5-(or 7)-methoxy-2-(4-nitrophenyl)-, (S)- (9CI) (CA INDEX NAME)



D1-O-Me

L4 ANSWER 226 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)
 REFERENCE COUNT: 24 THERE ARE 24 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT


L4 ANSWER 227 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
 ACCESSION NUMBER: 1997:11367 CAPLUS
 DOCUMENT NUMBER: 126:122464
 TITLE: Novel cathepsin Y and methods and compositions for inhibition thereof
 INVENTOR(S): Tung, Jay S.; Sinha, Sukanto; Mcconlogue, Lisa; Tatauno, Gwen; Anderson, John; Semko, Christopher M. F.; Chrysler, Susanna
 PATENT ASSIGNEE(S): Athena Neurosciences, Inc., USA
 SOURCE: PCT Int. Appl., 83 pp.
 CODEN: PIXXD2
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 9639194	A1	19961212	WO 1996-US6211	19960426
W: CA, JP				
RW: AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE				
US 5783434	A	19980721	US 1995-467607	19950606
US 5849711	A	19981215	US 1995-469362	19950606
CA 2221684	AA	19961212	CA 1996-2221684	19960426
EP 831920	A1	19980401	EP 1996-913917	19960426
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, FI				
JP 11506923	T2	19990622	JP 1996-500507	19960426
US 5858982	A	19990112	US 1997-850392	19970502
PRIORITY APPLN. INFO.: US 1995-467607 19950606				
US 1995-469362 19950606				
WO 1996-US6211 19960426				

OTHER SOURCE(S): MARPAT 126:122464
 AB Methods for inhibiting the secretion of β -amyloid peptide (BAP) from cells comprise administering to the cells certain compds. which inhibit the activity of an approx. 31 kD protease involved in BAP secretion. The 31 kD protease has been designated Cathepsin Y. Screening methods for BAP inhibitors rely on determining the activity of test compds. in the presence of Cathepsin Y and a suitable peptide substrate. This invention is also directed to a nucleic acid sequence that encodes Cathepsin Y and the expression and isolation of Cathepsin Y.
 IT 186030-84-2P
 RL: PNU (Preparation, unclassified); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)
 (novel cathepsin Y and methods and compns. for inhibition thereof)
 RN 186030-84-2 CAPLUS
 CN 9H-Xanthene-9-carboxamide, N-[2-(methoxymethylamino)-2-oxo-1-(phenylmethyl)ethyl]-, (S)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

L4 ANSWER 227 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)






L4 ANSWER 228 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
 ACCESSION NUMBER: 1997:11361 CAPLUS
 DOCUMENT NUMBER: 126:117068
 TITLE: Peptides and compounds that bind to the interleukin 1 (IL-1) receptor
 INVENTOR(S): Barrett, Ronald W.; Yanofsky, Stephen D.; Baldwin, David; Jacobs, Jeff W.; Bovy, Philippe R.; Leahy, Ellen M.; Pottorf, Richard S.; Dharanipragada, Ramalinga; Tomlinson, Ronald C.
 PATENT ASSIGNEE(S): Affymax Technologies N.V., UK; Barrett, Ronald W.; Yanofsky, Stephen D.; Baldwin, David; Jacobs, Jeff W.; Bovy, Philippe R.; Leahy, Ellen M.; Pottorf, Richard S.; Dharanipragada, Ramalinga; Tomlinson, Ronald C.
 SOURCE: PCT Int. Appl., 73 pp.
 CODEN: PIXXD2
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 6
 PATENT INFORMATION:


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 9639165	A1	19961212	WO 1996-US9835	19960605
W: AL, AM, AT, AU, AZ, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG				
RW: KE, LS, MW, SD, SZ, UG, AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA				
US 5861476	A	19990119	US 1995-464538	19950605
AU 9663820	A1	19961224	AU 1996-63820	19960605
EP 833654	A1	19980408	EP 1996-923258	19960605
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, FI				

PRIORITY APPLN. INFO.: US 1995-464538 19950605
 US 1994-190788 19940202
 US 1995-383474 19950201
 WO 1996-US9835 19960605
 AB Peptides that bind to the interleukin-1 type I receptor (IL-1Rt) can be used to assay the amount of IL-1R, or an IL-1R agonist or antagonist that is useful for treatment of interleukin 1-mediated inflammatory responses or diseases to infection, tissue injury, rheumatoid arthritis, osteoarthritis, psoriasis, inflammatory bowel disease, encephalitis, glomerulonephritis and respiratory distress syndrome. Also provided are peptides which bind to the IL-1Rt, which are 11 to 40 amino acids in length.
 IT 186251-93-4
 RL: ARU (Analytical role, unclassified); THU (Therapeutic use); ANST (Analytical study); BIOL (Biological study); USES (Uses)
 (peptides and compds. that bind to the interleukin 1 receptor)
 RN 186251-93-4 CAPLUS
 CN L-Tyrosine, N-acetyl-L-phenylalanyl-L-a-glutamyl-L-tryptophyl-L-threonyl-L-prolyl-D-alanyl-(3S)-2,3,4,9-tetrahydro-1H-pyrido[3,4-b]indole-3-carbonyl-L-tyrosyl-L-glutamyl-L-(2S)-2-azetidinecarbonyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.


L4 ANSWER 228 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)





ACCESSION NUMBER: 1997:64014 CAPLUS
 DOCUMENT NUMBER: 126:180811
 TITLE: Inhibition of the cardiac sarcolemma Na⁺/Ca²⁺ exchanger by conformationally constrained small cyclic peptides
 AUTHOR(S): Khananashvili, Daniel; Mester, Brenda; Saltoun, Miriam; Wang, Xiaolan; Shaulov, Gilat; Baazov, David
 CORPORATE SOURCE: Department of physiology and Pharmacology, Sackler School of Medicine, Tel-Aviv University, Ramat-Aviv, 69978, Israel
 SOURCE: Molecular Pharmacology (1997), 51(1), 126-131
 CODEN: MOPMA3; ISSN: 0026-895X
 PUBLISHER: Williams & Wilkins
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 AB Pos. charged cyclic peptides (three to seven amino acids) have been tested for their inhibitory effects on Na⁺/Ca²⁺ exchange in the cardiac sarcolemma vesicles. The lead structure of Phe-Arg-Cys-Arg-Cys-Phe-CONH₂ (FRCRCFa) has been systematically modified for identification of important pharmacophores. In cyclic peptides (intramol. S-S bond), the carboxyl terminal is locked with amide (CONH₂), and pos. charge is retained by one or two arginines, ornithines, or lysines. Thirty-five different cyclic peptides show IC₅₀ values in the range of 2-800 μM, suggesting that some specific structure-activity relationships may determine the inhibitory effects. Shortening of the FRCRCFa length to four amino acids decreases the inhibitory potency by 10-80-fold. The substitution of Arg2 or Arg4 in FRCRCFa with lysine or ornithine decreases the inhibitory potency by 5-12-fold, suggesting that both arginines are beneficial for inhibition. The substitution of Phe1 in FRCRCFa by 1,2,3,4-tetrahydroisoquinoline-3-carboxylic acid produces a potent inhibitor (IC₅₀ = 2-4 μM). The N-myristoylated FRCRCFa exhibits an inhibitory potency (IC₅₀ = 8-10 μM) similar to that of the parent FRCRCFa peptide, thereby arousing a new possibility for the development of a cell-permeable blocker of the Na⁺/Ca²⁺ exchanger. D-Arg4 or D-Cys5 substitutions in FRCRCFa do not alter the inhibitory effect, whereas the L-to-D substitutions of other amino acids in FRCRCFa reduce the inhibitory potency by 4-5-fold. Thus, the L-to-D substitutions of Arg4 and/or Cys5 have a potential to increase the peptide stability to proteolytic degradation. The insertion of proline outside of the ring of FRCRCFa diminishes the inhibitory potency by 3-6-fold, whereas proline introduction into the ring decreases the inhibitory potency by 16-20-fold. The replacement of Cys3 and Cys5 in FRCRCFa with β,β-dimethylcysteine has no significant effect on the inhibitory potency, suggesting that the S-S bond is not exposed to the interface of the peptide/receptor interaction. In conclusion, the current data support a proposal that the conformationally constrained Arg-Cys-Arg-Cys structure is obligatory for inhibition of Na⁺/Ca²⁺ exchange, whereas hydrophobic addns. at the carboxyl and amino ends have limited effects in increasing the inhibitory potency.
 IT 187536-11-4P
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); PRP (Properties); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation)
 (inhibition of cardiac sarcolemma Na⁺/Ca²⁺ exchanger by conformationally constrained small cyclic peptides in relation to structure)
 RN 187536-11-4 CAPLUS


L4 ANSWER 229 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)
 CN L-Phenylalaninamide, L-phenylalanyl-L-arginyl-L-cysteinyl-L-prolyl-L-arginyl-L-cysteinyl-, cyclic (3+5)-disulfide (9CI) (CA INDEX NAME)




REFERENCE COUNT: 20 THERE ARE 20 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

ACCESSION NUMBER: 1997:16200 CAPLUS
 DOCUMENT NUMBER: 126:99466
 TITLE: Induction of histamine release from rat mast cells by bradykinin analogs
 AUTHOR(S): Vietinghoff, Gabriele; Paegelow, Inge; Reissmann, Sigmund
 CORPORATE SOURCE: Dep. Pharmacol. Toxicol., Univ. Rostock, Rostock, D-18055, Germany
 SOURCE: Peptides (Tarrytown, New York) (1996), 17(8), 1467-1470
 CODEN: PPTDD5; ISSN: 0196-9781
 PUBLISHER: Elsevier
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 AB Independently of their agonistic or antagonistic activity on different isolated tissue preps., the kinin analogs investigated induce histamine release on rat peritoneal mast cells. The effectivity of most compds. is 10 to 100 times higher than that of bradykinin. Beside the pos. charged amino acids, the elongation at the N-terminus with hydrophobic amino acids and the replacement of amino acids in the bradykinin sequence (especially at position 7) with aromatic residues is important for a high histamine-releasing activity.
 IT 183664-42-8
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); PRP (Properties); BIOL (Biological study)
 (induction of histamine release from rat mast cells by bradykinin analogs)
 RN 183664-42-8 CAPLUS
 CN Bradykinin, 7-[(3R)-1,2,3,4-tetrahydro-3-isoquinolinecarboxylic acid]- (9CI) (CA INDEX NAME)



Absolute stereochemistry.






REFERENCE COUNT: 17 THERE ARE 17 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT


L4 ANSWER 231 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
 ACCESSION NUMBER: 1996:696042 CAPLUS
 DOCUMENT NUMBER: 126:26940
 TITLE: The use of the message-address concept in the design of potential antagonists based on dynorphin A
 AUTHOR(S): Kulkarni, S. N.; Choi, H.; Murray, T. F.; Delander, G. E.; Aldrich, J. V.
 CORPORATE SOURCE: College Pharmacy, Oregon State University, Corvallis, OR, 97331, USA
 SOURCE: Peptides: Chemistry, Structure and Biology, Proceedings of the American Peptide Symposium, 14th, Columbus, Ohio, June 18-23, 1995 (1996), Meeting Date 1995, 655-656. Editor(s): Kaumaya, Pravin T. P.; Hodges, Robert S. Mayflower Scientific: Kingswinford, UK.
 CODEN: 63NTAF
 DOCUMENT TYPE: Conference
 LANGUAGE: English
 AB The authors used the message-address concept to design dynorphin A (Dyn A) analogs as potential opioid receptor antagonists by combining the message sequence from novel peptides reported to have δ and μ antagonist activity with the C-terminal address sequence of Dyn A (1-11). The affinities of the peptides for κ , δ , and μ receptors were evaluated in radioligand binding assays using [3 H]DPDPE and [3 H]DAMGO and cloned opioid receptors stably expressed in CHO cells. Opioid activity was evaluated in the elec. stimulated guinea pig ileum. Results indicated that incorporation of a modified message sequence into Dyn A analogs can affect opioid activity and that C-terminal address sequence of Dyn A can be used to significantly enhance κ opioid receptor affinity.
 IT 184641-83-6P
 RL: BAC (Biological activity or effector, except adverse); BPR (Biological process); BSU (Biological study, unclassified); PRP (Properties); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation); PROC (Process)
 (opioid antagonist activity of peptides designed using message-address concept based on dynorphin A)
 RN 184641-83-6 CAPLUS
 CN 1-11-Dynorphin A (swine), 2-[(3S)-1,2,3,4-tetrahydro-3-isoquinolinecarboxylic acid]-3-L-phenylalanine-10-D-proline- (9CI) (CA INDEX NAME)
 Absolute stereochemistry.




L4 ANSWER 232 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
 ACCESSION NUMBER: 1996:695907 CAPLUS
 DOCUMENT NUMBER: 126:26919
 TITLE: Conformational re-addressing of peptides towards interactions with other specific receptors
 AUTHOR(S): Mikhaylovich, G. V.; Kolodziej, S. A.; Zhang, W. -J.; Nock, B.; Bernad, N.; Martinez, J.; Marshall, G. R.
 CORPORATE SOURCE: Center Molecular Design, Washington University, St. Louis, MO, 63130, USA
 SOURCE: Peptides: Chemistry, Structure and Biology, Proceedings of the American Peptide Symposium, 14th, Columbus, Ohio, June 18-23, 1995 (1996), Meeting Date 1995, 346-347. Editor(s): Kaumaya, Pravin T. P.; Hodges, Robert S. Mayflower Scientific: Kingswinford, UK.
 CODEN: 63NTAF
 DOCUMENT TYPE: Conference
 LANGUAGE: English
 AB To induce potent and selective peptide-receptor interactions, "message" functional groups of a ligand should be spatially arranged to satisfy a specific 3D "address" of receptor. In this way, almost any peptide containing corresponding "message" elements could be modified to bind a receptor with known 3D "address". To demonstrate this, conformationally constrained analogs were designed starting from the sequences of cholecystokinin and angiotensin fragments (CCK-8, Asp-Tyr-Met-Gly-Trp-Met-Asp-Phe-NH₂ and AT 4-8, Tyr-Val-His-Pro-Phe). The aim was to target δ -opioid receptor ("message" elements are shown in bold).
 IT 184637-94-3
 RL: BPR (Biological process); BSU (Biological study, unclassified); PRP (Properties); BIOL (Biological study); PROC (Process)
 (conformational re-addressing of peptides towards interactions with δ -opioid receptors)
 RN 184637-94-3 CAPLUS
 CN D-Phenylalanine, L-tyrosyl-D-cysteinyl-D-alanyl-(4S)-4-mercapto-L-prolyl-, cyclic (2+4)-disulfide (9CI) (CA INDEX NAME)
 Absolute stereochemistry.



L4 ANSWER 233 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
 1996:686665 CAPLUS
 DOCUMENT NUMBER: 126:54997
 TITLE: Structure-activity studies of B1 receptor-related peptides: antagonists
 AUTHOR(S): Gobeil, Fernand; Neugebauer, Withold; Filteau, Catherine; Jukic, Daniela; Allogho, Susanne Nsa; Pheng, Leng Hong; Nguyen-Le, Xuan Khai; Blouin, Daniel; Regoli, Domenico
 CORPORATE SOURCE: Dept. of Pharmacology, Univ. de Sherbrooke Medical School, Sherbrooke, QC, J1H 5N4, Can.
 SOURCE: Hypertension (Dallas) (1996), 28(5), 833-839
 CODEN: HPRTDN; ISSN: 0194-911X
 PUBLISHER: American Heart Association
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 AB We tested several peptides related to des-Arg9-bradykinin as stimulants or inhibitors of B1 (rabbit aorta, human umbilical vein) and B2 (rabbit jugular vein, guinea pig ileum, human umbilical vein) receptors. We also incubated the compds. with purified angiotensin-converting enzyme from rabbit lung to test their resistance to degradation. We evaluated apparent affinities (in terms of the affinity constant pA2) of compds. and their potential residual agonistic activities (αE). Bradykinin and des-Arg9-bradykinin were used as agonists for the B2 and B1 receptors, resp. Degradation of peptides by the angiotensin-converting enzyme was prevented in the presence of a D-residue in position 7 of des-Arg9-bradykinin. Replacement of Pro7 with D-Tic combined with Leu, Ile, Ala, or D-Tic in position 8 led to weak B1 receptor antagonists, some of which had strong residual agonistic activities on the B2 receptor preprns. The use of D-Pnal in position 7, combined with Ile in position 8 and Ac-Lys at the N-terminal (e.g., Ac-Lys[D-Pnal7,Ile8]des-Arg9-bradykinin) gave the most active B1 receptor antagonist (pA2 of 8.5 on rabbit aorta and human umbilical vein), which is also partially resistant to enzymic degradation. Extension of the N-terminal end by Sar-Tyr-αNhx (used for labeling purposes) and even cold-labeling to Tyr with iodine were compatible with high, selective, and specific antagonism of the B1 receptors. We compared some compds. with some already known B1 receptor antagonists to underline the novelty of new peptidic compds.
 IT 185052-06-6
 RL: BAC (Biological activity or effector, except adverse); BPR (Biological process); BSU (Biological study, unclassified); PRP (Properties); BIOL (Biological study); PROC (Process)
 (structure-activity studies of B1 receptor antagonist peptides)
 RN 185052-06-6 CAPLUS
 CN L-Phenylalanine, D-arginyl-L-arginyl-L-prolyl-(4R)-4-hydroxy-L-prolylglycyl-L-phenylalanyl-L-seryl-(3R)-1,2,3,4-tetrahydro-3-isoquinolinecarbonyl- (9CI) (CA INDEX NAME)
 Absolute stereochemistry.




PAGE 1-B



L4 ANSWER 234 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
 1996:681493 CAPLUS
 DOCUMENT NUMBER: 126:42242
 TITLE: Development of Potent Thrombin Receptor Antagonist Peptides
 AUTHOR(S): Bernatowicz, Michael S.; Klimas, Clifford E.; Hartl, Karen S.; Peluso, Marianne; Allegretto, Nick J.; Seiler, Steven M.
 CORPORATE SOURCE: Bristol-Myers Squibb Pharmaceutical Research Institute, Princeton, NJ, 08543, USA
 SOURCE: Journal of Medicinal Chemistry (1996), 39(25), 4879-4887
 CODEN: JMCMDR; ISSN: 0022-2623
 PUBLISHER: American Chemical Society
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 AB A peptide-based structure-activity study is reported leading to the discovery of novel potent thrombin receptor antagonists. Systematic substitution of nonproteogenic amino acids for the 2nd and 3rd residues of the human thrombin receptor tethered ligand sequence (SFLLR) led to a series of agonists with enhanced potency. The most potent pentapeptide agonist identified was Ser-p-fluorophenyl-p-guanidinophenyl-Leu-Arg-NH2 (I) (EC50 approx 0.04 μM for stimulation of human platelet aggregation, approx.10-fold more potent than the natural pentapeptide). Systematic substitution of the NH2-terminal Ser in I with neutral hydrophobic NH2-acyl groups led to partial agonists and eventually antagonists with unprecedented potency (>1000-fold increase over the previously reported antagonist 3-mercaptopropionyl-Phe-Cha-Cha-Arg-Lys-Pro-An-Asp-Lys-NH2). In the series of NH2-acyl tetrapeptide antagonists, N-trans-cinnamoyl-p-fluorophenyl-p-guanidinophenyl-Leu-Arg-NH2 (II) was identified as the tightest binding (IC50 approx 8 nM) and most potent with an IC50 approx 0.20 μM for inhibition of SFLLRNP-NH2-stimulated platelet aggregation. Systematic single substitutions in (II) indicated that, in addition to the NH2-terminal acyl group, the side chains at the 2nd and 3rd positions were also responsible for important and specific receptor interactions. The p-fluorophenyl and p-guanidinophenyl residues in the 2nd and 3rd positions of II were observed to be optimal in both the agonist and antagonist series. In the case of antagonists, however, an appropriately positioned pos. charged group (i.e., protonated base) at the 3rd residue was required. In contrast, such a substitution was not required for potent agonist activity. An even more potent antagonist resulted when II was extended at the C-terminus by a single Arg residue giving rise to analog RMS-200261 (III) which had an IC50 approx 20 nM for inhibition of SFLLRNP-NH2-stimulated platelet aggregation. When the C-terminal Arg of III was replaced by an Orn(N5-propionyl) residue, the resulting antagonist (RMS-20061) was suitable for use in radioligand binding assays (Kd = 10-30 nM). Antagonist activity observed for selected compds. was verified through secondary assays in that these analogs prevented SFLLRNP-NH2-stimulated GTPase activity in platelet membranes and Ca2+ mobilization in cultured human smooth muscle cells and mouse fibroblasts. Furthermore, this inhibition occurred at concns. that had no effect on thrombin catalytic activity, indicating a specific activity attributable to receptor binding and not enzyme inhibition.
 IT 185028-23-3P
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); PRP (Properties); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation)
 (development of potent thrombin receptor agonist and antagonist peptides)
 RN 185028-23-3 CAPLUS
 CN L-Argininamide, 4-fluoro-N-[(2-oxo-2H-1-benzopyran-3-yl)carbonyl]-L-

L4 ANSWER 234 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)
phenylalanyl-4-[(aminoiminomethyl)amino]-L-phenylalanyl-L-leucyl- (9CI)
(CA INDEX NAME)

Absolute stereochemistry.




L4 ANSWER 235 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 1996:639677 CAPLUS
DOCUMENT NUMBER: 125:295746
TITLE: N-coumarinyl- or N-quinolinonyl peptide p-nitroanilides as intramolecularly quenched fluorogenic substrates
AUTHOR(S): Kokotos, G.; Charitos, C.; Tzougraki, C.
CORPORATE SOURCE: Department Chemistry, University Athens, Athens, GR-15771, Greece
SOURCE: Peptides 1994, Proceedings of the European Peptide Symposium, 23rd, Braga, Port., Sept. 4-10, 1994 (1995), Meeting Date 1994, 891-892. Editor(s): Maia, Hernani L. S. ESCOM: Leiden, Neth.
CODEN: 63MBAO
DOCUMENT TYPE: Conference
LANGUAGE: English

AB Intramolecularly quenched fluorogenic substrates for proteases, i.e. a peptide chain bearing a fluorophore on one end and a quencher on the other, have been used for the determination of various proteases. The synthesis of four model substrates and studies on the quenching of fluorescence of aminocoumarin or aminoquinolinone-type fluorophores by the p-nitroanilide group are now reported for the assay of neutral endopeptidase-24.11.

IT 182944-07-6
RL: ARG (Analytical reagent use); BPR (Biological process); BSU (Biological study, unclassified); ANST (Analytical study); BIOL (Biological study); PROC (Process); USES (Uses)
RN 182944-07-6 CAPLUS
CN 2H-1-Benzopyran-3-carboxamide, N-[(1S)-2-[(4-nitrophenyl)amino]-2-oxo-1-(phenylmethyl)ethyl]-2-oxo- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




L4 ANSWER 236 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
ACCESSION NUMBER: 1996:639551 CAPLUS
DOCUMENT NUMBER: 126:1294
TITLE: New linear and cyclic bradykinin agonists and antagonists
AUTHOR(S): Reissmann, S.; Pineda, L. P.; Seyfarth, L.; Greiner, G.; Schoelkens, B.; Vietinghoff, G.; Paegelow, I.
CORPORATE SOURCE: Institute Biochem. & Biophys, Friedrich-Schiller-University, Jena, D-07743, Germany
SOURCE: Peptides 1994, Proceedings of the European Peptide Symposium, 23rd, Braga, Port., Sept. 4-10, 1994 (1995), Meeting Date 1994, 619-620. Editor(s): Maia, Hernani L. S. ESCOM: Leiden, Neth.
CODEN: 63MBAO
DOCUMENT TYPE: Conference
LANGUAGE: English

AB Despite the high potency of the bradykinin antagonist HOE-140 with the combination of D-Tic and Oic, unexpectedly D-Tic alone provides only a weak agonistic activity ([D-Tic]BK RUT: 7.20%) and Oic alone leads to a poor antagonist ([Oic]BK RUT: pA2 5.52). The combination of D-Tic with hydrophobic amino acids other than Oic is unable to give potent antagonists. The hydrophobic and constrained amino acid Oic at position 8 destroys the activity of potent agonists but enhances the antagonistic potencies of analogs with D-amino acids at position 7. Therefore, the authors conclude that in antagonists D-Tic can be replaced by other aromatic and non-aromatic amino acids, but Oic at position 8 is necessary for enhancement of the antagonistic activity. Beside the well known type of bradykinin antagonists with the key replacement at position 7 the authors could obtain two other types with amino acid replacements at positions 5 and 2.

IT 183664-42-8
RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); PRP (Properties); BIOL (Biological study)
RN 183664-42-8 CAPLUS
CN Bradykinin, 7-[(3R)-1,2,3,4-tetrahydro-3-isoquinolinecarboxylic acid]- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



PAGE 1-A


L4 ANSWER 236 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)

PAGE 1-B




L4 ANSWER 237 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
 ACCESSION NUMBER: 1996:494173 CAPLUS
 DOCUMENT NUMBER: 125:143330
 TITLE: Peptide compounds for prevention and/or treatment of nitric oxide (NO)-mediated diseases
 INVENTOR(S): Itoh, Yoshikuni; Iwamoto, Toshiro; Yatabe, Takumi; Hamashima, Hitoshi; Inoue, Takayuki; Hashimoto, Seiji; Oku, Teruo
 PATENT ASSIGNEE(S): Fujisawa Pharmaceutical Co., Ltd., Japan
 SOURCE: PCT Int. Appl., 739 pp.
 CODEN: PIXXD2
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 9616981	A2	19960606	WO 1995-JP2428	19951129
WO 9616981	A3	19960906		
W:	AU, CA, CN, FI, HU, JP, KR, MX, NO, NZ, RU, UA, US			
RW:	AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, CA, GN, ML, MR, NE, SN, TD, TG			
AU 9539937	A1	19960619	AU 1995-39937	19951129
EP 796270	A2	19970924	EP 1995-938602	19951129
R:	AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LI, LU, NL, PT, SE			
ZA 9510201	A	19960625	ZA 1995-10201	19951130
US 5932737	A	19990803	US 1997-849076	19970530
PRIORITY APPLN. INFO.:			GB 1994-24408	A 19941202
			GB 1995-4891	A 19950310
			GB 1995-10042	A 19950518
			WO 1995-JP2428	W 19951129
OTHER SOURCE(S):		MARPAT 125:143330		
GI				



AB Peptides WA1NR8CH(A2T)CONR9CH(A3R3)R4 [W = alkyl, (un)substituted aryl or fluorenyl, etc.; A1 = alkylene, NHCO, CO, CS, SO2; A2 = alkylene; T = H, aryl, heterocyclyl, OH, etc.; R8 = H, alkyl; R9 may link with A2T to form CH2C6H4CH2-O (O); A3 = bond, alkylene, R3 = H, aryl, OH, etc.; R9 = H, alkyl or may link with A3R3 to form Q; R4 = CO2H, protected carboxy, carboxamido, etc. or CH(A3R3)R4 = N-alkyl-2-oxoquinoline moiety] or their pharmaceutically acceptable salts were prepared for use as medicaments. Thus, dipeptide I was prepared by acylation of aspartylphenylalaninamide derivative with 2-benzofurancarboxylic acid. I and six other peptides showed

L4 ANSWER 237 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)
 100% inhibition of NO prodn. in tests of murine macrophage cells.
 IT 179875-74-2P
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)
 (preparation of peptides for prevention and/or treatment of nitric oxide-mediated diseases)
 RN 179875-74-2 CAPLUS
 CN 2(1H)-Isoquinolinecarboxylic acid, 3,4-dihydro-3-[[[2-[methyl(phenylmethyl)aminol-2-oxo-1-(phenylmethyl)ethyl]amino]carbonyl]-, 1,1-dimethylethyl ester, [S-(R*,R*)]- (9CI) (CA INDEX NAME)
 Absolute stereochemistry.




L4 ANSWER 238 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
 ACCESSION NUMBER: 1996:466915 CAPLUS
 DOCUMENT NUMBER: 125:143315
 TITLE: Boronic ester and acid compounds, synthesis and uses
 INVENTOR(S): Adams, Julian; Ma, Yu-Ting; Stein, Ross; Baevsky, Matthew; Grenier, Louis; Flamondon, Louis
 PATENT ASSIGNEE(S): Proscript, Inc., USA
 SOURCE: PCT Int. Appl., 144 pp.
 CODEN: PIXXD2
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 3
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 9613266	A1	19960509	WO 1995-US14117	19951027
W:	AL, AM, AT, AU, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK			
RW:	KE, LS, MW, SD, SZ, UG, AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG			
US 6083903	A	20000704	US 1995-442581	19950516
AU 9641398	A1	19960523	AU 1996-41398	19951027
AU 710564	B2	19990923		
EP 788360	A1	19970813	EP 1995-939670	19951027
EP 788360	B1	20030528		
R:	AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LI, LU, MC, NL, PT, SE			
JP 10510245	T2	19981006	JP 1995-514834	19951027
AT 241631	E	20030615	AT 1995-939670	19951027
FI 9701746	A	19970606	FI 1997-1746	19970423
NO 9701929	A	19970612	NO 1997-1929	19970425
PRIORITY APPLN. INFO.:			US 1994-330525	A 19941028
			US 1995-442581	A 19950516
			WO 1995-US14117	W 19951027
OTHER SOURCE(S):		MARPAT 125:143315		


AB Peptidyl boronic acids and esters PNR[B(R1X1)ACHR2X2CHR3BZ1Z2] (P = aryl-, aralkyl-, heteroaryl-, or heteroarylalkylcarbonyl or -sulfonyl; B1 = N, CH; X1, X2 = CONH, CH(OH)CH2, COCH2; A = 0, 1, 2; R = H, alkyl; RR1 or RR2 (for A = 0) may form a ring; R1, R2, R3 = H, alkyl, cycloalkyl, aryl, etc.; Z1, Z2 = alkyl, hydroxy, alkoxy, aryloxy; Z1Z2 may form a moiety derived from a dihydroxy compound) and their pharmaceutically acceptable salts were prepared. The rate of degradation of proteins of an animal can be reduced by contacting cells of the animal with these boronic compds. Thus, N-(4-morpholinecarbonyl)-β-(1-naphthyl)-L-alanine-L-leucine boronic acid was prepared by coupling (1S,2S,3R,5S)-pinanediol leucine boronate trifluoroacetate salt with N-Boc-β-(1-naphthyl)-L-alanine, followed by deprotection, acylation with 4-morpholinecarbonyl chloride, and cleavage of the pinanediol moiety.

IT 179324-53-9
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); BIOL (Biological study)
 (synthesis of peptidyl boronic acids and esters as proteolytic enzyme inhibitors)
 RN 179324-53-9 CAPLUS
 CN Boronic acid, [(1R)-3-methyl-1-[(2S)-1-oxo-3-phenyl-2-[(2-quinolinylcarbonyl)amino]propyl]amino]butyl]- (9CI) (CA INDEX NAME)
 Absolute stereochemistry.

L4 ANSWER 238 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)




L4 ANSWER 239 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
 ACCESSION NUMBER: 1996:457338 CAPLUS
 DOCUMENT NUMBER: 125:248351
 TITLE: Ordered conformations in bis(amino acid) derivatives of 1,1'-ferrocenedicarboxylic acid
 AUTHOR(S): Herrick, Richard S.; Jarret, Ronald M.; Curran, Timothy P.; Dragoli, Dean R.; Flaherty, Maryellen B.; Lindyberg, Susan E.; Slate, Rebecca A.; Thornton, Lisa C.
 CORPORATE SOURCE: Dep. Chem., Coll. Holy Cross, Worcester, MA, 01610, USA
 SOURCE: Tetrahedron Letters (1996), 37(30), 5289-5292
 CODEN: TETLAV; ISSN: 0040-4039
 PUBLISHER: Elsevier
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 GI




AB Two bis(amino acid) derivs. I (R = Me2CH, PhCH2) of 1,1'-ferrocenedicarboxylic acid were characterized by IR, 1H NMR and 13C NMR spectroscopy. Each was found to adopt an ordered, intramolecularly H bonded conformation in CHCl3.
 IT 181589-78-6P
 RL: PRP (Properties); SPN (Synthetic preparation); PREP (Preparation) (ordered conformations in bis(amino acid) derivs. of ferrocenedicarboxylic acid)
 RN 181589-78-6 CAPLUS
 CN Ferrocene, 1,1'-bis[[[(1S)-2-methoxy-2-oxo-1-(phenylmethyl)ethyl]amino]carbonyl]- (9CI) (CA INDEX NAME)

L4 ANSWER 239 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)



L4 ANSWER 240 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
 ACCESSION NUMBER: 1996:443908 CAPLUS
 DOCUMENT NUMBER: 125:115147
 TITLE: Preparation of peptide aldehyde derivatives as cysteine protease inhibitors
 INVENTOR(S): Sohma, Takashi; Fujisawa, Yukio; Yasuma, Tsuneo; Mizoguchi, Junji
 PATENT ASSIGNEE(S): Takeda Chemical Industries, Ltd., Japan
 SOURCE: PCT Int. Appl., 85 pp.
 CODEN: PIXXD2
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 9610014	A1	19960404	WO 1995-JP1933	19950925
W: AM, AU, BB, BG, BR, BY, CA, CN, CZ, EE, FI, GE, HU, IS, KG, KR, KZ, LK, LR, LT, LV, MD, MG, MK, MN, MX, NO, NZ, PL, RO, RU, SG, SI, SK, TJ, TM, TT, UA, US, UZ, VN				
RW: KE, MW, SD, SZ, UG, AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG				
CA 2196182	AA	19960404	CA 1995-2196182	19950925
AU 9535341	A1	19960419	AU 1995-35341	19950925
JP 08151355	A2	19960611	JP 1995-245957	19950925
EP 783489	A1	19970716	EP 1995-932228	19950925
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LI, LU, NL, PT, SE				
PRIORITY APPL. INFO.: JP 1994-231839 19940927				
WO 1995-JP1933 19950925				
OTHER SOURCE(S): MARPAT 125:115147				
GI				




AB The present invention relates to acylaminoaldehyde compds. of formula R4-Q-NHCH(R1)-X-CHO [Q = one or two amino acid residual groups which may be substituted; R1 = hydrogen atom or an optionally substituted hydrocarbon or heterocyclic group; R4 = an optionally esterified carboxyl group or an acyl group; X = a optionally substituted straight-chain or branched divalent hydrocarbon group having a chain length of 1 to 4 atoms as the linear moiety], or salts thereof, which have strong cysteine protease inhibitory activities and are useful as prophylactic and therapeutic agent of various diseases, including bone diseases, caused by abnormal exasperation of cystine protease, are prepared. Thus, 2.4 g N-tert-butoxycarbonyl-L-phenylalanyl-L-tryptophanal and 1.76 g (formylmethylene)triphenylphosphorane were dissolved in 10 mL THF and 30 mL toluene and stirred for 15 h to give the title compound (I; R = Boc-Phe).

L4 ANSWER 240 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)
 The latter compd. and I (R = PhCH2O2C-Leu-Leu) (II) in vitro showed IC50 of 3.5 + 10-8 and 9.7 + 10-9 M, resp., against cathepsin L and that of 2.4 + 10-6 and 9.7 + 10-7 M, resp., against cathepsin B, resp. In a bone resorption inhibitory assay, they in vitro inhibited by 83 and 51%, resp., the Ca release from fetal rat's forearm bones. A gelatin capsule formulation contg. II was described.
 IT 178910-81-1P
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses) (preparation of peptide aldehyde derivs. as cysteine protease inhibitors and bone resorption inhibitors for treating bone diseases)
 RN 178910-81-1 CAPLUS
 CN 2-Quinolincarboxamide, N-[1-[(4-hydroxyphenyl)methyl]-2-[[1-(1H-indol-3-ylmethyl)-4-oxo-2-butenyl]amino]-2-oxoethyl]-, [S-(R*,R*)]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.
 Double bond geometry unknown.




L4 ANSWER 241 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
 ACCESSION NUMBER: 1996:393848 CAPLUS
 DOCUMENT NUMBER: 125:25636
 TITLE: 2-Phenyl-4-quinolinecarboxamides: A Novel Class of Potent and Selective Non-Peptide Competitive Antagonists for the Human Neurokinin-3 Receptor
 AUTHOR(S): Giardina, Giuseppe A. M.; Sarau, Henry M.; Farina, Carlo; Medhurst, Andrew D.; Grugni, Mario; Foley, James J.; Raveglia, Luca F.; Schmidt, Dulcie B.; Rigolio, Roberto; et al.
 CORPORATE SOURCE: Department of Chemistry, SmithKline Beecham S.p.A., Baranzate, 20021, Italy
 SOURCE: Journal of Medicinal Chemistry (1996), 39(12), 2281-2284
 CODEN: JMCMAR; ISSN: 0022-2623
 PUBLISHER: American Chemical Society
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 AB A novel class of potent and selective, non-peptide NK-3 receptor antagonists, based on the 2-phenylquinoline framework, has been identified and characterized by binding anal. using membrane preparation of CHO cells expressing the human neurokinin receptors (hNKs-CHO). Functional activity was determined by inhibition of senktide-induced contraction of the rabbit isolated iris sphincter muscle preparation. An extensive structure-activity study led to the identification of (S)-(-)-N-(α -ethylbenzyl)-3-hydroxy-2-phenylquinoline-4-carboxamide (SB 223412) as the most potent (K_i = 1.0 nM in hNK-3-CHO binding; K_D = 5.4 nM for antagonism of senktide-induced contraction in rabbit iris sphincter muscle) and selective (hNK-2/hNK-3 K_i ratio of 144 and hNK-1/hNK-3 K_i ratio > 100,000) hNK-3 receptor antagonist of this class. In addition, NK3-induced Ca^{2+} mobilization studies in hNK-3-HEK 293 cells indicated that SB 223412 is a reversible, competitive antagonist. Compds. from this novel class will be extremely useful in the functional characterization of hNK-3 receptors and elucidation of potential therapeutic indications for selective hNK-3 receptor antagonists.
 IT 174635-51-9P
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation)
 (preparation and structure-activity of human neurokinin 3 receptor antagonists phenylquinolinecarboxamides)
 RN 174635-51-9 CAPLUS
 CN Benzeneacetic acid, α -[[(2-phenyl-4-quinolinyl)carbonyl]amino]-, methyl ester (9CI) (CA INDEX NAME)




L4 ANSWER 241 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)

L4 ANSWER 243 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
 ACCESSION NUMBER: 1996:366713 CAPLUS
 DOCUMENT NUMBER: 125:76666
 TITLE: Modification of the C-terminal dipeptide of angiotensin II yielded a novel series of analogs with II (AT2) receptor selectivity
 AUTHOR(S): Cody, Wayne L.; He, John X.; Lunney, Elizabeth A.; Humblet, Christine C.; Lu, Gina H.; Panek, Robert L.; Dudley, David T.
 CORPORATE SOURCE: Department Chemistry, Parke-Davis Pharmaceutical Research, Ann Arbor, MI, 48105, USA
 SOURCE: Protein and Peptide Letters (1996), 3(2), 107-112
 CODEN: PPELEN; ISSN: 0929-8665
 PUBLISHER: Bentham Science Publishers BV
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 AB Angiotensin II is a potent pressor agent acting through two distinct G-protein coupled receptors. The type I (AT1) receptor is responsible for the pressor activity while the function of the type II (AT2) receptor remains unclear. Specific modifications of the C-terminal dipeptide (-Pro7-Phe8) of angiotensin II with constrained aromatic (Tic) and hydrophobic (oic) amino acids have led to analogs with negligible affinity for the AT1 receptor, but nanomolar affinity for the AT2 receptor. These compds. may provide useful tools to help delineate the physiol. role of the AT2 receptor.
 IT 178403-48-0
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); BIOL (Biological study)
 (modification of C-terminal dipeptide of angiotensin II yielded series of analogs AT2 receptor selectivity)
 RN 178403-48-0 CAPLUS
 CN Angiotensin II, 1-(N-methylglycine)-5-L-valine-7-(L-1,2,3,4-tetrahydro-3-isoquinolinecarboxylic acid)- (9CI) (CA INDEX NAME)
 Absolute stereochemistry.



L4 ANSWER 242 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
 ACCESSION NUMBER: 1996:367060 CAPLUS
 DOCUMENT NUMBER: 125:81189
 TITLE: Application of capillary electrophoresis-electrospray ionization mass spectrometry in the determination of molecular diversity
 AUTHOR(S): Dunayevskiy, Yuriy M.; Vouros, Paul; Wintner, Edward A.; Shipp, Gerald W.; Carell, Thomas; Rebek, Julius, Jr.
 CORPORATE SOURCE: Dep. Chem., Northeastern Univ., Boston, MA, 02115, USA
 SOURCE: Proceedings of the National Academy of Sciences of the United States of America (1996), 93(12), 6152-6157
 CODEN: PNASAC; ISSN: 0027-8424
 PUBLISHER: National Academy of Sciences
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 AB By capillary electrophoresis coupled online to electrospray ionization MS, a library of theor. 171 distributed xanthene derivs. was analyzed. The method allowed the purity and makeup of the library to be determined: 160 of the expected compds. were found to be present, and 12 side-products were also detected in the mixture. Due to the ability of capillary electrophoresis to sep. analytes on the basis of charge, most of the xanthene derivs. could be resolved by simple capillary electrophoresis-MS procedures even though 124 of the 171 theor. compds. were isobaric with \pm 1 mol. in the mixture. Any remaining unresolved peaks were resolved by MS/MS expts. The method shows promise for the anal. of small combinatorial libraries with <1000 components.
 IT 178916-07-9P
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)
 (application of capillary electrophoresis-electrospray ionization mass spectrometry in determination of mol. diversity of xanthene derivs.)
 RN 178916-07-9 CAPLUS
 CN L-Phenylalanine, N-[[2,7-bis(1,1-dimethylethyl)-5-[[[(2-methoxy-2-oxoethyl)amino]carbonyl]-9,9-dimethyl-9H-xanthen-4-yl]carbonyl]-, 1,1-dimethylethyl ester (9CI) (CA INDEX NAME)


Absolute stereochemistry.



PAGE 1-A




L4 ANSWER 244 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
 ACCESSION NUMBER: 1996:353996 CAPLUS
 DOCUMENT NUMBER: 125:1605
 TITLE: Novel Cyclic Analogs of Angiotensin II with Cyclization between Positions 5 and 7: Conformational and Biological Implications
 AUTHOR(S): Zhang, Wei-Jun; Nikiforovich, Gregory V.; Perodin, Jacqueline; Richard, Darren E.; Escher, Emanuel; Marshall, Garland R.
 CORPORATE SOURCE: Department of Molecular Biology and Pharmacology, Washington University, St. Louis, MO, 63130, USA
 SOURCE: Journal of Medicinal Chemistry (1996), 39(14), 2738-2744
 CODEN: JMCNAR; ISSN: 0022-2623
 PUBLISHER: American Chemical Society
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 AB To study the conformational features of mol. recognition of angiotensin II (Asp1-Arg2-Val3-Tyr4-Val5/Ile5-His6-Pro7-Phe8, AII), the synthesis and biol. testing of several cyclic analogs of AII cyclized between positions 5 and 7 have been performed. The synthesized analogs were Sar1-Arg2-Val3-Tyr4-cyclo(Cys5-His6-Pen7)-Phe8 (3), Sar1-Arg2-Val3-Tyr4-cyclo(Asp5-His6-Apt7)-Phe8 (4), Sar1-Arg2-Val3-Tyr4-cyclo(Glu5-His6-Apt7)-Phe8 (5), Sar1-Arg2-Val3-Tyr4-cyclo(Cys5-His6-Mpt7)-Phe8 (6), Sar1-Arg2-Val3-Tyr4-cyclo(Cys5-His6-Mpc7)-Phe8 (7), Sar1-Arg2-Val3-Tyr4-cyclo(Hcy5-His6-Mpc7)-Phe8 (8), and Sar1-Arg2-Val3-Tyr4-cyclo(Hcy5-His6-Mpc7)-Phe8 (9), where Apt stands for 4-amino-trans-proline, and Mpc and Mpc for 4-mercapto-trans- and -cis-prolines, resp. Compound (9) showed good affinity at AT-1 receptors, namely a $K_D = 20$ nM. In functional assays, it showed the characteristics of a weak partial agonist with a relative affinity of 0.26% of that for AII and an intrinsic efficacy, αE , of 0.42. Mol. modeling suggested a possible explanation for this finding: the relatively strong binding and the weak partial agonistic activity of compound 9 are due to interaction with AT-1 receptor of only two functionally important groups, namely, the side chains of the His6 and Phe8 residues.
 IT 177480-67-0P
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); PRP (Properties); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation)
 (conformational and biol. implications of novel cyclic analogs of angiotensin II with cyclization between positions 5 and 7)
 RN 177480-67-0 CAPLUS
 CN L-Phenylalanine, N-methylglycyl-L-arginyl-L-valyl-L-tyrosyl-L- α -glutamyl-L-histidyl-trans-4-amino-L-prolyl-, cyclic (5-7)-peptide (9CI) (CA INDEX NAME)
 Absolute stereochemistry.




L4 ANSWER 245 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
 ACCESSION NUMBER: 1996:341820 CAPLUS
 DOCUMENT NUMBER: 125:33490
 TITLE: Preparation of quinoline-4-carboxamides and related compounds as NK3 antagonists.
 INVENTOR(S): Farina, Carlo; Giardina, Giuseppe; Arnaldo Mari; Grugni, Mario; Raveglia, Luca; Francesco
 PATENT ASSIGNEE(S): Smithkline Beecham Farmaceutici S.P.A., Italy
 SOURCE: PCT Int. Appl., 28 pp.
 CODEN: PIXXD2
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 9602509	A1	19960201	WO 1995-EP2638	19950706
W: JP, US				
RW: AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE				
PRIORITY APPLN. INFO.:			IT 1994-MI1466 19940714	
OTHER SOURCE(S):			MARPAT 125:33490	
GI				




AB Title compds. I; Ar = (substituted) Ph, naphthyl, heterocyclyl; R = (substituted) Ph, heterocyclyl, CHR4R5; R4 = H, alkyl, cycloalkyl, (substituted) Ph, heteroaryl, etc.; R5 = alkyl, (CH2)nAr; n = 0-3; R1 = H, alkyl; R2, R3 = H, alkyl, alkenyl, aryl, carboxamido, sulfonamido, alkoxy, OH, halo, NO2, cyano, hydroxyalkyl, aminoalkyl, acylamino, CO2H, alkylsulfonamino, etc.; X = O, S, H2, NCN, were prepared. Thus, benzylamine, 2-phenylquinoline-4-carbonyl chloride, and K2CO3 were stirred in DMF at 0°-room temperature overnight to give N-benzyl-2-phenylquinoline-4-carboxamide. The latter inhibited binding of 125I-N-Me-Phe7-NKB to guinea pig cortical membranes with IC50 = 620 nM.
 IT 189815-92-7P
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)
 (preparation of quinoline-4-carboxamides and related compds. as NK3 antagonists)
 RN 189815-92-7 CAPLUS
 CN Phenylalanine, N-[(2-phenyl-4-quinolyl)carbonyl]-, methyl ester (9CI) (CA INDEX NAME)




LA	ANSWER 246 OF 261	CAPLUS COPYRIGHT 2004 ACS ON STN
ACCESSION NUMBER:	1996:J39994	CAPLUS
DOCUMENT NUMBER:	125:56105	
TITLE:	The importance of the peptide bond at position 2 in HCO-Met-Leu-Phe-OMe analogs as shown by studies on human neutrophils	
AUTHOR(S):	Cavicchioli, Giorgio; Breveglieri, Angela; Boggiani, Marina; Vertuani, Gianni; Reali, Eva; Spisani, Susanna	
CORPORATE SOURCE:	Department Pharmaceutical Sciences, University Ferrara, Ferrara, Italy	
SOURCE:	Journal of Peptide Science (1996), 2(3), 135-140 CODEN: JPSIIR; ISSN: 1075-2617	
PUBLISHER:	Wiley	
DOCUMENT TYPE:	Journal	
LANGUAGE:	English	
AB	<p>The formylpeptides formyl-methionyl-N-methylleucyl-phenylalanine Me ester [for-Met-(NMe)Leu-Phe-OMe], formyl-methionyl-2-aminoctetralin-2-carboxyl-phenylalanine Me ester [for-Met-Atc-Phe-OMe] 2, formyl-methionyl-1,2,3,4-tetrahydroisoquinoline-3-carboxyl-phenylalanine Me ester [for-Met-Tic-Phe-OMe] and formyl-methionyl-2-aminox-4-methylvaleryl-phenylalanine Me ester [for-Met-OLeu-Phe-OMe] were synthesized in order to investigate the role of the amide bond at position 2 on biol. activities on human neutrophils. Only analog 2, which keeps the NH group at position 2, was found to retain bioactivity for neutrophils (chemotaxis, superoxide formation, and lysozyme release), though sterically encumbered.</p>	
IT	<p>177656-62-1 RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); PRP (Properties); BIOL (Biological study) (amide bond of formyl-Met-Leu-Phe peptide analogs is important for chemotactic activity toward neutrophils)</p>	
RN	177656-62-1 CAPLUS	
CN	L-phenylalanine, N-[[[(3S)-2-[(2S)-2-(formylamino)-4-(methylthio)-1-oxobutyl]-1,2,3,4-tetrahydro-3-isoquinolinyllcarbonyl]-, methyl ester (9CI) (CA INDEX NAME)	

Absolute stereochemistry.



14 ANSWER 247 OF 261 CAPLUS COPYRIGHT 2004 ACS ON STN
ACCESSION NUMBER: 1996:328195 CAPLUS
DOCUMENT NUMBER: 125:323
TITLE: Esters and Amides of 6-(Chloromethyl)-2-oxo-2H-1-benzopyran-3-carboxylic Acid as Inhibitors of α -Chymotrypsin: Significance of the "Aromatic" Nature of the Novel Ester-Type Coumarin for Strong Inhibitory Activity
AUTHOR(S): Pochee, Lionel.; Domet, Caroline; Schynts, Marc; Thierry, Nicole.; Boggetto, Nicole; Pirote, Jean-Beard; Jiang, Kai Y.; Maserel, Bernard; de Tullio, Pascal; et al.
CORPORATE SOURCE: Laboratoire de Chimie Pharmaceutique, Universite de Liege, Liege, B-4000, Belg.
SOURCE: Journal of Medicinal Chemistry (1996), 39(13), 2579-2585
CODEN: JMCMAR; ISSN: 0022-2623
PUBLISHER: American Chemical Society
DOCUMENT TYPE: Journal
LANGUAGE: English
AB A series of esters and amides of 6-(chloromethyl)-2-oxo-2H-1-benzopyran-3-carboxylic acid were synthesized and evaluated in vitro for their inhibitory activity toward bovine α -chymotrypsin and human leukocyte elastase. Both series behaved as time-dependent inhibitors of α -chymotrypsin, but ester-type coumarins were clearly more efficient than the corresponding amides in inactivating the serine protease. The best inactivation was observed with "aromatic" esters, in particular with meta-substituted Ph esters such as m-chlorophenyl 6-(chloromethyl)-2-oxo-2H-1-benzopyran-3-carboxylate, which appears to be one of the most powerful inactivators of α -chymotrypsin yet reported (kinact/KI = 760 000 M⁻¹ s⁻¹ at pH 7.5 and 25°). Usually, the coumarin derivative failed to inhibit significantly human leukocyte elastase. As a result, the reported series of aromatic coumarinic esters behaves as a new chemical family of selective α -chymotrypsin inhibitors.
IT 167770-52-8P
RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); PRP (Properties); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation)
(preparation of esters and amides of 6-(chloromethyl)-2-oxo-2H-1-benzopyran-3-carboxylic acid as inhibitors of α -chymotrypsin)
RN 167770-52-8 CAPLUS
CN L-Phenylalanine, N-[[6-(chloromethyl)-2-oxo-2H-1-benzopyran-3-yl]carbonyl]-L-methyl ester (9CI) (CA INDEX NAME)


Absolute stereochemistry



14 ANSWER 248 OF 261 CAPLUS COPYRIGHT 2004 ACS ON STN
 ACCESSION NUMBER: 1596-313253 CAPLUS
 DOCUMENT NUMBER: 125:80883
 TITLE: Isolation and identification of peptide conformers by reversed-phase high-performance liquid chromatography and NMR at low temperature
 AUTHOR(S): Kalman, Andras; Thuncke, Frank; Schmidt, Ralf; Schiller, Peter W.; Horvath, Csaba
 CORPORATE ENTITY: Department of Chemical Engineering, Yale University, P. O. Box 208286, New Haven, CT, 06520-8286, USA
 SOURCE: Journal of Chromatography, A (1996), 729(1 + 2), 155-171
 CODEN: JCHRAEY; ISSN: 0021-9673
 PUBLISHER: Elsevier
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 AB Peptide conformers with one or more rotationally hindered peptide bonds due to the presence of proline and/or another N-substituted amino acid residue in the molecule separated by reversed-phase chromatog. at low temps., isolated and identified by NMR. The scope of this investigation included the cis-trans isomers of the dipeptides as Leu-Pro, Phe-Pro and Tyr-Pro as well as conformers of opioid peptides containing proline and/or the proline-like Tic (1,2,3,4-tetrahydro-isouquinoline-3-carboxylic acid) residues: Phe-Pro-Phe (β -casomorphin 1-3 fragment), Tyr-Tic-Phe-Phe, Tyr-Pro-Phe-Pro-Gly (β -casomorphin-5), Tyr-Tic-Phe-Phe-Val-Val-Gly-NH₂ and Tyr-Tic-Phe-Gly-Tyr-Pro-Ser-NH₂. Chromatog. with micropellicular and totally porous octadecylated silica stationary phases and aqueous methanol under isocratic conditions resulted in well separated peaks of the rotational isomers at sufficiently low temps. Preparative RP-HPLC was carried out with eluents containing water and methanol, both deuterated, and the effluent fractions containing each isomer were collected for further investigation. The conformational states of the peptide isomers upon separation were conserved by storing the effluent fractions in liquid nitrogen. The Leu-Pro, Phe-Pro, Tyr-Pro and Tyr-Pro-Phe conformers were identified by one- and two-dimensional NMR spectroscopy at -15°C. Upon comparing the NMR spectra of the isomers, for these peptides the retention order of the conformers was unambiguously established: in each case the trans conformer is eluted before the cis conformer. On the basis of NMR data obtained with β -casomorphin-5, which contains 2 proline residues, the elution order of its 4 conformers was established by NMR spectroscopy of the fractions obtained by RP-HPLC at low temperature as trans-trans (least retained), trans-cis, cis-cis and cis-trans (most retained).
 IT 19748-31-7
 RL: ANP (Analyte); PRP (Properties); ANST (Analytical study)
 (Identification of peptide conformers by reversed-phase HPLC and NMR)
 RN 19748-31-7 CAPLUS
 CN Glycine, L-tyrosyl-L-1,2,3,4-tetrahydro-3-isouquinolinecarbonyl-L-phenylalanyl- (9CI) (CA INDEX NAME)


Absolute stereochemistry.

L4 ANSWER 248 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)




L4 ANSWER 249 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
 ACCESSION NUMBER: 1996:290583 CAPLUS
 DOCUMENT NUMBER: 124:343279
 TITLE: Preparation of naphth[2,1-d]isoxazole-3-carboxamide derivatives as antiulcer drugs
 INVENTOR(S): Hasegawa, Yukio; Sato, Michitaka; Hasumi, Koichi; Yamamoto, Norio; Matsui, Teruaki; Shidara, Kazuhiro; Kenjo, Takashi; Myazawa, Katsuhiko; Ogawa, Chisato; Et, Al.
 PATENT ASSIGNEE(S): Teikoku Hormone Mfg Co Ltd, Japan
 SOURCE: Jpn. Kokai Tokkyo Koho, 20 pp.
 CODEN: JKXXAF
 DOCUMENT TYPE: Patent
 LANGUAGE: Japanese
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 08027131	A2	19960130	JP 1994-180457	19940711
PRIORITY APPLN. INFO.:			JP 1994-180457	19940711
OTHER SOURCE(S):		MARPAT 124:343279		



I




II

AB Naphthoisoaxazole derivs. [I; A = CH, CH2, S, O, SO2; R1 = H, alkyl; R2 = hydroxyalkyl, alkoxyalkyl, heterocyclyl containing 1-4 heteroatoms selected from N, S, and O; n = 2-5; R1R2N = heterocyclyl; R3, R4 = H, halo, alkyl, alkoxy, alkenyloxy, OH; when A is CH or CH2, R1 is H] and their salts are prepared for use as antiulcer drugs. Thus, 3-carboxynaphth[2,1-d]isoxazole was treated with PCl5 and then reacted with 5-amino-1H-tetrazole to give 3-(1H-tetrazol-5-ylcarbonyl)naphth[2,1-d]isoxazole (II), which inhibited stress-induced ulcer at 30 mg/kg oral in male rats.

IT 176432-23-8P
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)
 (preparation of naphth[2,1-d]isoxazole-3-carboxamide derivs. as antiulcer drugs)


L4 ANSWER 249 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)
 RN 176432-23-8 CAPLUS
 CN L-Phenylalanine, N-[(4,5-dihydro-7-methoxynaphth[2,1-d]isoxazol-3-yl)carbonyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




L4 ANSWER 250 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
 ACCESSION NUMBER: 1996:257145 CAPLUS
 DOCUMENT NUMBER: 125:34134
 TITLE: Synthesis, structure and stability of novel dimeric peptide-disulfides
 AUTHOR(S): Leban, Johann J.; Spaltenstein, Andrew; Landavazo, Antonio; Chestnut, William; Aulabaugh, Ann; Taylor, Lester C. E.; Daniels, Alejandro J.
 CORPORATE SOURCE: Wellcome Res. Labs., Research Triangle Park, NC, 27709, USA
 SOURCE: International Journal of Peptide & Protein Research (1996), 47(3), 161-6
 CODEN: IJPPC3; ISSN: 0367-8377
 PUBLISHER: Munksgaard
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 AB Oxidation of nonapeptide dithiol H-Ile-Cys-Pro-Cys-Tyr-Arg-Leu-Arg-Tyr-NH2 with K3Fe(CN)6 leads to either monomeric disulfide (4) or antiparallel and parallel dimeric disulfides (3a and 3b) depending upon reaction conditions. When exposed to small amts. of thiols or cyanide in aqueous solution, these three species interconvert to an equilibrium mixture of 2:1:8 (3a:3b:4).
 IT 177582-22-8P
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)
 (preparation, structure and stability of novel dimeric peptide disulfides)
 RN 177582-22-8 CAPLUS
 CN L-Tyrosinamide, L-isoleucyl-L-cysteinyl-L-prolyl-L-cysteinyl-L-tyrosyl-L-arginyl-L-leucyl-L-arginyl-, cyclic (2+4)-disulfide (9CI) (CA INDEX NAME)


PAGE 1-A



PAGE 2-A



PAGE 3-A



ACCESSION NUMBER: 1996:177848 CAPLUS
 DOCUMENT NUMBER: 124:232269
 TITLE: Quinoline derivatives as tachykinin NK3 receptor antagonists
 INVENTOR(S): Farina, Carlo; Giardina, Giuseppe Arnaldo Mari; Grugni, Mario; Raveglia, Luca Francesco
 PATENT ASSIGNEE(S): Smithkline Beecham Farmaceutici S.P.A., Italy
 SOURCE: PCT Int. Appl., 95 pp.
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 9532948	A1	19951207	WO 1995-EP2000	19950523
W: AM, AT, AU, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IS, JP, KE, KG, KP, KZ, LK, LR, LT, LU, LV, MD, MG, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TT				
RW: KE, MW, SD, SZ, UG, AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG				
CA 2191352	AA	19951207	CA 1995-2191352	19950523
CA 2191352	C	20010130		
CA 2257662	AA	19951207	CA 1995-2257662	19950523
AU 9526164	A1	19951221	AU 1995-26164	19950523
AU 699319	B2	19981203		
HU 76286	A2	19970728	HU 1996-3262	19950523
CN 1156451	A	19970806	CN 1995-194338	19950523
CN 1092642	B	20021016		
BR 9507788	A	19970923	BR 1995-7788	19950523
EP 804419	A1	19971105	EP 1995-920894	19950523
EP 804419	B1	20030806		
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI				
JP 10500697	T2	19980120	JP 1996-500287	19950523
RO 114445	B3	19990430	RO 1996-2234	19950523
EP 940391	A2	19990908	EP 1998-204483	19950523
EP 940391	A3	19991110		
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI				
JP 2000026314	A2	20000125	JP 1999-172597	19950523
NZ 329979	A	20000728	NZ 1995-329979	19950523
RU 2155754	C2	20000910	RU 1996-124804	19950523
JP 2002179594	A2	20020626	JP 2001-326622	19950523
SK 282721	B6	20021106	SK 1996-1514	19950523
SK 282722	B6	20021106	SK 1999-47	19950523
CZ 291476	B6	20030312	CZ 1996-3470	19950523
AT 246677	E	20030815	AT 1995-920894	19950523
PL 186075	B1	20031031	PL 1995-317381	19950523
PT 804419	T	20031231	PT 1995-920894	19950523
PL 186665	B1	20040227	PL 1995-341889	19950523
ZA 9504269	A	19960514	ZA 1995-4269	19950525
US 5811553	A	19980922	US 1995-450438	19950525

US 6608083 B1 20030819 US 1995-450437 19950525
 TW 427977 B 20010401 TW 1995-84105319 19950526
 TW 533199 B 20030521 TW 1999-88121625 19950526
 BG 64004 B1 20030930 BG 1996-101008 19961125
 FI 9604712 A 19970123 FI 1996-4712 19961126
 NO 9605036 A 19970124 NO 1996-5036 19961126
 CN 1276211 A 20001213 CN 1999-100978 19990115
 AU 9912162 A1 19990325 AU 1999-12162 19990119
 FI 9900268 A 19990210 FI 1999-268 19990210
 NO 9901813 A 19970124 NO 1999-1813 19990416
 US 2003236281 A1 20031225 US 2001-867133 20010529
 CN 1428145 A 20030709 CN 2002-107941 20020318




PRIORITY APPLN. INFO.:

IT 1994-MI1099 A 19940527
 IT 1995-MI494 A 19950314
 AU 1995-26164 A3 19950523
 CA 1995-2191352 A 19950523
 EP 1995-920894 A3 19950523
 JP 1996-500287 A 19950523
 NZ 1995-287442 A1 19950523
 WO 1995-EP2000 W 19950523
 US 1995-450437 A3 19950525

OTHER SOURCE(S):

MARPAT 124:232269

GI



AB NK3 receptor antagonists I (Ar = (un)substituted Ph, naphthyl, cycloalkadienyl, heteroaryl; R = (un)substituted alkyl, cycloalkyl, (un)substituted Ph, phenylalkyl, or heteroaryl, CO2H and derivs., etc.; R1, R2 = H, alkyl; or R1R2 = (CH2)3-5; or R1R2 = (CH2)2-5; R3, R4 = H, alkyl, alkenyl, aryl, alkoxy, OH, halo, NO2, amino, etc.; R5 = alkyl, cycloalkyl, (un)substituted (hetero)aryl; X = O, S, N(CN)) are useful in treating pulmonary, CNS, and neurodegenerative disorders, etc. Approx. 115 compds. were prepared. For example, amidation of 3-methyl-2-phenylquinoline-4-carbonyl chloride with (R)-α-ethylbenzylamine gave title compound II in 58% yield. II had IC50 of 5.6 nM for displacement of [3H]-senktide from guinea-pig cortical NK3 receptors. Antagonist activity of I was shown by inhibition of senktide-induced contraction of guinea-pig ileum.

IT 174635-51-99

RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)
 (preparation of quinolinecarboxamide derivs. as tachykinin NK3 receptor antagonists)

RN 174635-51-9 CAPLUS

CN Benzeneacetic acid, α-[[2-phenyl-4-quinolinyl]carbonyl]amino]-, methyl ester (9CI) (CA INDEX NAME)

L4 ANSWER 252 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
 ACCESSION NUMBER: 1996:12187 CAPLUS
 DOCUMENT NUMBER: 124:202986
 TITLE: Synthesis of the diastereomers of β -Me-Tyr and B-Me-Phe and their effect on the biological properties of the delta opioid receptor antagonist TIPP
 AUTHOR(S): Marnekens, Els; Tourwe, Dirk; Vanderstichele, Sylvia; Nguyen Thi Diem, Trang; Toth, Geza; Peter, Antal; Chung, Nga N.; Schiller, Peter W.
 CORPORATE SOURCE: Org. Chem., Free Univ., Brussels, B-1050, Belg.
 SOURCE: Letters in Peptide Science (1995), 2(3/4), 190-2
 CODEN: LPSCEM; ISSN: 0929-5666
 PUBLISHER: ESCOM
 DOCUMENT TYPE: Journal
 LANGUAGE: English


AB In order to influence side chain conformations and to increase the μ -agonist properties of the δ -selective opioid receptor δ -antagonist H-Tyr-Tic-Phe-Phe-NH₂ (TIPP; Tic = 1,2,3,4-tetrahydroisoquinoline-3-carboxylic acid), residues Tyr¹, Phe³ and Phe⁴ were replaced by their β -methyl-substituted stereoisomers. Synthesis of β -Me-Tyr was carried out in a stereoselective way. Incorporation of the modified amino acids was performed by solid-phase methods. Receptor binding data and GPI and MVD bioassays were obtained for all stereoisomers, in general showing equal or slightly increased potencies. In the [(R,S)- β -Me-Phe]³ analog, the introduction of the β -Me substituent restores signal transduction.

IT 174147-54-7P
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation)
 (preparation of β -methyltyrosine and β -methylphenylalanine and their effect on the opioid antagonistic activity of TIPP)

RN 174147-54-7 CAPLUS
 CN L-Phenylalaninamide, L-tyrosyl-(3S)-1,2,3,4-tetrahydro-3-isoquinolinecarboxyl- (BR) β -methyl-L-phenylalanyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

L4 ANSWER 252 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)




L4 ANSWER 253 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
 ACCESSION NUMBER: 1996:12183 CAPLUS
 DOCUMENT NUMBER: 124:135903
 TITLE: Towards nonpeptide agonists: design of 'true' peptidomimetics
 AUTHOR(S): Nikiforovich, Gregory V.
 CORPORATE SOURCE: Cent. Mol. Design, Washington Univ., St. Louis, MO, 63130, USA
 SOURCE: Letters in Peptide Science (1995), 2(3/4), 172-6
 CODEN: LPSCEM; ISSN: 0929-5666
 PUBLISHER: ESCOM
 DOCUMENT TYPE: Journal
 LANGUAGE: English

AB This paper outlines the basic strategy to design 'true' peptidomimetics, i.e., nonpeptide compds. that bind to the same receptor site as the parent peptide. Design of highly selective and potent agonist analogs of δ -opioid peptides based on development of the 3D model for the δ -opioid pharmacophore is described. The design employed mol. modeling in combination with synthesis, biol. testing, NMR spectroscopy and x-ray studies. The designed compds. were able to bind the δ -opioid receptors with affinities and selectivities comparable to those for DPDPE, a well-known δ -selective agonist. They also showed moderate δ -agonistic activity.

IT 173555-72-1
 RL: BAC (Biological activity or effector, except adverse); BPR (Biological process); BSU (Biological study, unclassified); PRP (Properties); BIOL (Biological study); PROC (Process)
 (design of δ -opioid agonists from peptidomimetics in relation to conformation)

RN 173555-72-1 CAPLUS
 CN D-Phenylalanine, L-tyrosyl-D-cysteinyglycyl-trans-4-mercapto-L-prolyl-, cyclic (2-4)-disulfide (9CI) (CA INDEX NAME)

Absolute stereochemistry.




L4 ANSWER 254 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
 ACCESSION NUMBER: 1995:998406 CAPLUS
 DOCUMENT NUMBER: 124:203098
 TITLE: Preparation of peptide factor Xa inhibitors as antithrombotics
 INVENTOR(S): Al-Obeidi, Fahad; Lebl, Michal; Ostrem, James A.; Safar, Pavel; Stierandova, Alena; Strop, Peter; Walser, Armin
 PATENT ASSIGNEE(S): Selectide Corp., USA
 SOURCE: PCT Int. Appl., 107 pp.
 CODEN: PIXXD2
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 9529189	A1	19951102	WO 1995-US5268	19950425
W: AM, AU, BB, BG, BR, BY, CA, CN, CZ, EE, FI, GE, HU, JP, KG, KP, KR, KZ, LK, LR, LT, LV, MD, MG, MN, MW, MX, NO, NZ, PL, RO, RU, SI, SK, TJ, TT, UA, UZ, VN				
RW: KE, MW, SD, SZ, UG, AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG				
CA 2186497	AA	19951102	CA 1995-218647	19950425
AU 9523683	A1	19951116	AU 1995-23683	19950425
AU 707653	B2	19990715		
ZA 9503361	A	19960112	ZA 1995-3361	19950425
EP 758341	A1	19970219	EP 1995-917736	19950425
EP 758341	B1	20040324		
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LI, LU, MC, NL, PT, SE				
CN 1147261	A	19970409	CN 1995-192811	19950425
HU 76346	A2	19970828	HU 1996-2954	19950425
JP 10503477	T2	19980331	JP 1995-527853	19950425
RU 2152954	C1	20000720	RU 1996-122647	19950425
EE 3973	B1	20030217	EE 1996-146	19950425
EP 1384725	A2	20040128	EP 2003-21617	19950425
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, MC, NL, PT, IE				
IL 113505	A1	19991231	IL 1995-113505	19950426
TW 409129	B	20001021	TW 1995-84104681	19950511
FI 9604317	A	19961025	FI 1996-4317	19961025
NO 9604553	A	19961227	NO 1996-4553	19961025
LT 4218	B	19970925	LT 1996-151	19961025
LV 11740	B	19971220	LV 1996-410	19961115
US 5849510	A	19981215	US 1997-547794	19971008
PRIORITY APPL. INFO.:			US 1994-233054	A 19940426
			EP 1995-917736	A3 19950425
			US 1995-428404	B1 19950425
			WO 1995-US5268	W 19950425

OTHER SOURCE(S): MARPAT 124:203098
 AB A1-A2-(A3)m-B (m = 0, 1; A1 = R1-R2-R3; A2 = R4-R5-R6; A3 = R7-R8-R9; R1 = (substituted) 1-20 amino acid residues, R1CO, R1R12X; X = N, CH, NCO; R11, R12 = H, alkyl, acyl, aryl, aralkyl, protecting group; R2 = CHR9R100; R9, R100 = H, (substituted) alkyl, aralkyl, heteroalkyl, heteroaryl; R3 = CO, CH2, CHR9CO, etc.; R4 = CH2, imino; R5 = CR20R202; R201, R202 = H, (substituted) alkyl, aryl, aralkyl; R6 = CO, CH2, CHR9CO; R7 = (substituted) R4; R8 = CR210R211; R210, R211 = H, (substituted) alkyl, alkylaryl, heterocyclyl; R9 = CO, CH2, CHR9CO; B = (substituted) 1-20 amino acid residues, amino, OH, alkoxy, acyloxy, etc.; with provisoal, were prepared Thus, Ac-Tyr-Chg-Arg-NH₂ (Chg = cyclohexylglycyl) inhibited

L4 ANSWER 254 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)
 coagulation in human plasma with EC50 = 2.5 µM.
 IT 174132-79-7P
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)
 (preparation of peptide factor Xa inhibitors as antithrombotics)
 RN 174132-79-7 CAPLUS
 CN L-Alaninamide, 4-(aminoiminomethyl)-N-(3-isoquinolylcarbonyl)-L-phenylalanyl-L-2-cyclohexylglycyl-3-(1-methylpyridinium-3-yl)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



L4 ANSWER 255 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
 ACCESSION NUMBER: 1995:911314 CAPLUS
 DOCUMENT NUMBER: 124:87771
 TITLE: Synthesis, tritium labeling and binding characterization of new delta opioid receptor selective antagonists, TIPP and TIPP[w]
 AUTHOR(S): Toth, G.; Nevin, S.; Borsodi, A.; Nguyen, T. M. -D.; Schiller, P.
 CORPORATE SOURCE: Biological Research Center, Hungarian Academy of Sciences, Szeged, H-6701, Hung.
 SOURCE: Synthesis and Applications of Isotopically Labelled Compounds 1994, Proceedings of the International Symposium, 5th, Strasbourg, June 20-24, 1994 (1995), Meeting Date 1994, 141-4. Editor(s): Allen, John; Voges, Rolf. Wiley: Chichester, UK.
 CODEN: 61UMAF
 CONFERENCE
 DOCUMENT TYPE: English
 LANGUAGE: English
 AB A symposium report on the preparation and δ opioid receptor binding affinities of the title tritiated peptide antagonists H-Tyr-Tic-Phe-Phe-OH (TIPP; Tic = 1,2,3,4-tetrahydroisoquinoline-3-carboxylic acid) and its analog TIPP[w] with a pseudopeptide bond between Tic and Phe (H-Tyr-Ticw[CH2NH]Phe-Phe-OH), prepared to obtain a more stable ligand under radioreceptor assay conditions. The two peptides were labeled by tritium using precursors containing 3,5-diiodotyrosine. After catalytic dehalogenation with tritium gas, the crude labeled peptides were purified by HPLC. Specific radioactivity was 1.87 TBq/mmol for TIPP and 1.76 TBq/mmol for TIPP[w]. The tritiated ligands labeled rat brain membrane binding sites with Kd values under the nanomolar range and Bmax values were found to be 82 and 105 fmol/mg protein for TIPP and TIPP[w], resp. Both tritiated ligands proved to be highly selective for δ opioid receptors in binding competition expts.
 IT 172490-62-9P
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation)
 (preparation, tritium labeling and binding characterization of new δ opioid receptor selective antagonist peptides)
 RN 172490-62-9 CAPLUS
 CN L-Phenylalanine, L-tyrosyl-3,5-t2-L-1,2,3,4-tetrahydro-3-isoquinolinecarbonyl-L-phenylalanyl- (9CI) (CA INDEX NAME)


Absolute stereochemistry.

L4 ANSWER 255 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)



L4 ANSWER 256 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
 ACCESSION NUMBER: 1995:804314 CAPLUS
 DOCUMENT NUMBER: 123:228900
 TITLE: Preparation ofazole-fused peptides as substance P antagonists and analgesics.
 INVENTOR(S): Morgan, Barry A.; Gordon, Thomas D.; Hansen, Philip E.; Singh, Jasbir
 PATENT ASSIGNEE(S): Sterling Winthrop Inc., USA
 SOURCE: U.S., 34 pp. Cont. of U.S. Ser. No. 131,706, abandoned.
 CODEN: USXXAM
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 5378803	A	19950103	US 1992-912949	19920710
PRIORITY APPLN. INFO.:		US 1987-131706	19871211	
OTHER SOURCE(S):		MARPAT 123:228900		
GI				



AB Title compds. [I; Q1 = Pro, bond; Q2 = Pro, D-Trp, bond; Q3 = Pro, D-Trp, Phe, (R) (2,3,4,9-tetrahydro)-1H-pyrido[3,4-b]indol-2-yl-3-carboxyl, bond; Q4 = Pro, D-Trp, Phe, bond; Q5 = D-Trp, Phe, bond; Q6 = Leu, Met, bond; Q7 = Phe, N-MePhe, Met, bond; R1 = H, 2, BOC; R2 = H; R3 = Me2CH, Me2CHCH2, H2N(CH2)4, PhCH2, 4-HOC6H4CH2, pyridylmethyl, (1H-indol-3-yl)methyl, R2R3 = atoms to form 2,3,4,9-tetrahydro-1H-pyrido[3,4-b]indol-2,3-diyl, R5 = H, OH or alkali metal salt thereof, MeO, EtO, amino, etc.; X = oxo thia, imido; Y = oxo, H2; n = 0,1; starred center is L or D; with a proviso], were prepared. Thus, title compound (II; starred center has D-configuration), prepared via thionation of 2-D-Trp-NH2 with P2S5 and cyclocondensation of the product with Et 3-chloro-2-oxo-3-phenylpropionate, antagonized substance P in the guinea pig ileum test with pA2 = 7.3, and in the mouse acetylcholine-induced writhing test showed intrathecal ED50 = 0.78 µg/mouse.
 IT 167982-54-9P
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)


L4 ANSWER 256 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)
(prepn. of azole-fused peptides as substance P antagonists and
analgesics)

RN 167982-54-9 CAPLUS
CN 1H-Pyrido[3,4-b]indole-3-carboxamide, 2,3,4,9-tetrahydro-N-[2-[[2-[(1H-indol-3-yl)-1-[5-(1-methylethyl)-4-[[methyl(2-phenylethyl)amino]methyl]-2-thiazolyl]ethyl]amino]-2-oxo-1-(phenylmethyl)ethyl]-, [3R-[3R*(S*(R*))]]-, phosphate (1:2) (9CI) (CA INDEX NAME)

CM 1

CRN 167982-53-8
CMF C47 H51 N7 O2 S

Absolute stereochemistry.



CM 2

CRN 7664-38-2
CMF H3 O4 P



L4 ANSWER 257 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
ACCESSION NUMBER: 1995:801429 CAPLUS

DOCUMENT NUMBER: 123,256711

TITLE: Preparation of gastrin and CCK receptor ligands
INVENTOR(S): Kalindjian, Sarkis Barret; Steel, Katherine Isobel
Mary; Pether, Michael John; Davies, Jonathan Michael
Richard; Low, Caroline Minli Rachel; Hudson, Martin
Lyn; Buck, Ildiko Maria; McDonald, Iain Mair;
Dunstone, David John; Tozer, Matthew John
James Black Foundation Ltd., UK

PATENT ASSIGNEE(S):
SOURCE: PCT Int. Appl., 124 pp.

DOCUMENT TYPE: Patent
LANGUAGE: English

FAMILY ACC. NUM. COUNT: 2
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 9504720	A2	19950216	WO 1994-GB1741	19940809
WO 9504720	A3	19950803		
W: AM, AT, AU, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, ES, FI, GB, GE, HU, JP, KE, KG, KP, KR, KZ, LK, LT, LU, LV, MD, MG, MN, MW, NL, NO, NZ, PL, PT, RO, RU, SD, SE, SI, SK, TJ, TT, UA, US, UZ, VN				
RW: KE, MW, SD, AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG				
AU 9473478	A1	19950228	AU 1994-73478	19940809
AU 682051	B2	19970918		
EP 720601	A1	19960710	EP 1994-922318	19940809
EP 720601	B1	20001025		
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LI, LU, MC, NL, PT, SE				
JP 09502430	T2	19970311	JP 1994-506306	19940809
HU 75301	A2	19970528	HU 1996-70	19940809
AT 197146	E	20001115	AT 1994-922318	19940809
ES 2152989	T3	20010216	ES 1994-922318	19940809
PT 720601	T	20010228	PT 1994-94922318	19940809
PL 181782	B1	20010928	PL 1994-312960	19940809
ZA 9405998	A	19960212	ZA 1994-5998	19940810
GB 2290539	A1	19960103	GB 1995-2503	19950209
WO 9532949	A1	19951207	WO 1995-GB1194	19950525
W: AM, AT, AU, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, ES, FI, GB, GE, HU, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LT, LU, LV, MD, MG, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TT				
RW: KE, MW, SD, SZ, UG, AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG				
AU 9525342	A1	19951221	AU 1995-25342	19950525
EP 763026	A1	19970319	EP 1995-919561	19950525
EP 763026	B1	20030326		
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LI, LU, MC, NL, PT, SE				
JP 10504525	T2	19980506	JP 1995-500483	19950525
AT 235470	E	20030415	AT 1995-919561	19950525
ZA 9504315	A	19961126	ZA 1995-4315	19950526
NO 9600488	A	19960315	NO 1996-488	19960206
FI 9600572	A	19960207	FI 1996-572	19960207
US 5795907	A	19980818	US 1996-583008	19960318

L4 ANSWER 257 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)

US 5912260 A 19990615 US 1996-737725 19961219
US 5919829 A 19990706 US 1998-64849 19980423
GR 3035100 T3 20010330 GR 2000-402788 20001218

PRIORITY APPL. INFO.:

GB 1993-16508 A 19930810

GB 1994-10688 A 19940527

WO 1994-GB1741 W 19940809


GB 1995-2503 A 19950209

WO 1995-GB1194 W 19950525

OTHER SOURCE(S):

MARPAT 123:256711

GI



AB Title compds. [e.g. I; A = atoms to complete a bicyclic ring system; R1 = halo, NH2, cyano, OH, alkyl, CO2H, etc.; 1 of X, W = CO and the other = CO, SO, SO2; Y = NR3R4, hydrocarbyloxy, etc.; R3 = H, hydrocarbyl, etc.; R4 = H, alkyl, (un) esterified CH2CO2H; Z = OH, alkoxy, OPh, (un)substituted NH2, NH2R, etc.; R = H, cyano, alkyl, CH2OH, CO2H, etc.; Z1 = alkylene; m = 0-6] were prepared. Thus, 4-methylphthalic anhydride which was converted in 6 steps to indole-5,6-dicarboxylic anhydride which was amidated by adamantane-1-methylamine and the product amidated by (S)-3,5-(PhH2CO2C)2C6H3NHCOCH(NH2)CH2Ph (preparation given) to give, in 2 addnl. steps, title compound (S)-II the di-N-methyl-D-glucamine salt of which had pK1 of 9.4 for binding at mouse cortex CCKB receptors in vitro.


IT 167991-35-79
RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)
(preparation of gastrin and CCK receptor ligands)

RN 167991-35-7 CAPLUS

CN 1,3-Benzenedicarboxylic acid, 5-[[1-oxo-3-phenyl-2-[[[3-[[[tricyclo[3.3.1.1.3,7]dec-1-ylmethyl]amino]carbonyl]-1H-carbazol-2-yl]carbonyl]amino]propyl]amino]-, (S)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

L4 ANSWER 257 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)




L4 ANSWER 258 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
 ACCESSION NUMBER: 1995:789155 CAPLUS
 DOCUMENT NUMBER: 123:199414
 TITLE: Preparation of peptidylactol derivatives as inhibitors of cathepsin L.
 INVENTOR(S): Sohma, Takashi; Fujisawa, Yukio; Oi, Satoru; Mizoguchi, Junji
 PATENT ASSIGNEE(S): Takeda Chemical Industries, Ltd., Japan
 SOURCE: Eur. Pat. Appl., 54 pp.
 CODEN: EPXXDW
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
EP 641800	A1	19950308	EP 1994-113669	19940901
EP 641800	B1	20020116		
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LI, LU, NL, PT, SE				
NO 9403210	A	19950306	NO 1994-3210	19940830
JP 08104685	A2	19960423	JP 1994-208981	19940901
AT 212036	E	20020215	AT 1994-113669	19940901
CA 2131397	AA	19950304	CA 1994-2131397	19940902
FI 9404040	A	19950304	FI 1994-4040	19940902
AU 9471682	A1	19950316	AU 1994-71682	19940902
AU 678493	B2	19970529		
HU 68717	A2	19950728	HU 1994-2536	19940902
CN 1106001	A	19950802	CN 1994-115669	19940902
US 5496834	A	19960305	US 1994-300738	19940902

PRIORITY APPLN. INFO.:
 JP 1993-219655 A 19930903
 JP 1994-168501 A 19940720
 JP 1994-190385 A 19940812

OTHER SOURCE(S): MARPAT 123:199414
 GI




AB Title compds. (I; Q = 1-2 (substituted) amino acid residues; R3 = (esterified) carboxyl, acyl; A = alkylene; B = H, (substituted) alkyl, acyl), were prepared. Thus, N-benzoyloxycarbonylhomoserine, 1-hydroxybenzotriazole, and 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide were stirred 14 h in DMF at ice temp. room temperature to give 84.3% (S)-3-(N-benzoyloxycarbonylamino)tetrahydrofuran-2-one. This was hydrogenolyzed in EtOH over Pd/C and the product was stirred with BOC-Phe-OH, 1-hydroxybenzotriazole, and 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide in DMF to give 78.3% (S)-3-(N-tert-butoxycarbonylphenylalanylaminotetrahydrofuran-2-one. The latter in THF was treated with DIBAL in PhMe at -72° to give 37.5% title compound (II). I inhibited cathepsin L with IC50 = 6.9 + 10-7-8.0 + 10-9 M, and at 10-30 µM gave 26-82% inhibition of bone resorption in


L4 ANSWER 258 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)
 rat fetuses according to the method of Raisz.

IT 167766-31-6P
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)
 (preparation of peptidylactol derivs. as inhibitors of cathepsin L)
 RN 167766-31-6 CAPLUS
 CN 3-Quinolonecarboxamide, N-[2-oxo-1-(phenylmethyl)-2-[(tetrahydro-2-hydroxy-3-furanyl)amino]ethyl]-, [2R-[2α,3α(S*)]]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




L4 ANSWER 259 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
 ACCESSION NUMBER: 1995:723478 CAPLUS
 DOCUMENT NUMBER: 123:340785
 TITLE: Synthesis of new quinolin-2-carbonyl-N-amino acid derivatives with evaluation of their antimicrobial activity
 AUTHOR(S): Shalaby, A.M.; Kassem, E.M.M.; Farrag, H.A.
 CORPORATE SOURCE: National Research Centre, Cairo, Egypt
 SOURCE: Proceedings of the Pakistan Academy of Sciences (1994), 31(3), 163-73
 CODEN: PKSPAW; ISSN: 0377-2969
 PUBLISHER: Pakistan Academy of Sciences
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 GI



AB 2-Quinolonecarbonyl chloride and 4-(2-quinolonecarbonylamino)benzoyl chloride (I and II; X = bond, R = Cl) reacted with amino acid esters H-X-OMe (X = Gly, L-Val, DL-Met, L-3,5-diiodotyrosine) in THF to give the corresponding amides II (R = OMe), which on hydrolysis gave the free acids II (R = OH). Condensation of II (R = OMe) with N2H4 gave the corresponding hydrazides II (R = NHNH2), which in turn were condensed with aromatic aldehydes R1CHO (R1 = 4-pyridyl, (MeO)3C6H2) to give the Schiff bases II (R = NHN:CHR1). The antimicrobial activity of the prepared compds. was also studied.
 IT 170488-13-8P
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); RCT (Reactant); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation); RACT (Reactant or reagent)
 (preparation and antimicrobial activity of new (quinolonecarbonyl)amino acid derivs.)
 RN 170488-13-8 CAPLUS
 CN L-Tyrosine, 3,5-diiodo-N-(2-quinolinylcarbonyl)-, methyl ester (9CI) (CA INDEX NAME)

Absolute stereochemistry.




L4 ANSWER 260 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN
 ACCESSION NUMBER: 1995:657214 CAPLUS
 DOCUMENT NUMBER: 123:314483
 TITLE: Synthesis and some pharmacological properties of seven new analogs of Asa-D-Tyr(Et)-Phe-Val-Asn-Abu-Pro-Arg-Arg-NH2, a potent linear antagonist of V2 receptors
 AUTHOR(S): Czaja, M.; Konieczna, E.; Wierzbna, T.; Juzwa, W.; Lamnek, B.
 CORPORATE SOURCE: Dep. Chem., Univ. Gdansk, Gdansk, 80-952, Pol.
 SOURCE: Polish Journal of Chemistry (1995), 69(4), 552-8
 CODEN: PJCHDQ; ISSN: 0137-5083
 PUBLISHER: Polish Chemical Society
 DOCUMENT TYPE: Journal
 LANGUAGE: English

AB Seven new analogs of potent V2/V1 antagonist adamantaneacetyl-D-Tyr(Et)-Phe-Val-Asn-Abu-Pro-Arg-Arg-NH2 (I) were synthesized. Four of them were designed by substitution of positions 2 or 3 with L- or D-1,2,3,4-tetrahydroisoquinolinecarboxylic acid. One peptide was designed by replacement of Phe3 residue by L-β-thienylalanine. Two compds. have a disulfide bridge on the C-terminal end of model peptide I. The anti-antidiuretic activity of the analogs was evaluated by their ability to inhibit the antidiuretic effect of endogenous arginine-vasopressin (AVP). The antipressor potency of the peptides was assayed by their ability to inhibit the pressor response to exogenous AVP. The modifications proposed are incompatible with biol. potency, particularly anti-V2 activity. Nevertheless, two of the new analogs are potent vasopressor antagonists.


IT 169824-40-2P
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation)
 (synthesis and anti-antidiuretic and antipressor activities of analogs of potent linear antagonist of V2 receptors)
 RN 169824-40-2 CAPLUS
 CN L-Argininamide, N-[[1,2,3,4-tetrahydro-2-(tricyclo[3.3.1.1.3,7]dec-1-ylacetyl)-3-isoquinolyl]carbonyl]-L-phenylalanyl-L-valyl-L-asparaginyl-L-2-aminobutanoyl-L-prolyl-L-arginyl-, (S)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

PAGE 1-A



PAGE 1-B



L4 ANSWER 261 OF 261 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER:

1995:633575 CAPLUS

DOCUMENT NUMBER:

123:314476

TITLE:

Acid catalysis in the formation of dioxopiperazines from peptides containing tetrahydroisoquinoline-3-carboxylic acid at position 2

AUTHOR(S):

Capasso, Sante; Sica, Filomena; Mazzarella, Lelio; Balboni, Gianfranco; Guerrini, Remo; Salvadori, Severo

CORPORATE SOURCE:

Dep. Chem., Univ. Naples "Federico II, Naples, Italy International Journal of Peptide & Protein Research (1995), 45(6), 567-73

SOURCE:

CODEN: IJPPC3; ISSN: 0367-8377

PUBLISHER:

Munksgaard

DOCUMENT TYPE:

Journal

LANGUAGE:

English

AB

The kinetics of the spontaneous formation of 2,5-dioxopiperazines from peptides containing the Tic (1,2,3,4-tetrahydroisoquinoline-3-carboxylic acid) residue in the 2-position of the sequence has been studied in DMSO and water solution. The reaction is first order in Tic-peptide and subject to general-acid catalysis. Moreover, only the fraction of peptide having the amino terminal group in the deprotonated state reacts with appreciable rate. In pure organic solvent, and in aqueous solution with low buffer concentration, the

degradation reaction of Tic-peptides is very low; at 20° for the peptide H-Tyr-Tic-Phe-Phe-NH₂, in DMSO and in neutral water in the absence of buffer, the half-lives (t_{1/2}) are 3 + 104 and 1.2 + 104 h, resp. The addition of carboxylic acids or buffers to the reaction solns. markedly increases the reaction rate; in 0.01 M HAc in DMSO and in 0.1 M phosphate buffer in water, pH 7.1, t_{1/2} values for the tetrapeptide are 61 and 121 h, resp.

IT

169611-83-0P
RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)
(acid catalysis in formation of dioxopiperazines from peptides containing tetrahydroisoquinolinecarboxylic acid)

RN

169611-83-0 CAPLUS
CN L-Phenylalaninamide, N-[(2-[(1,1-dimethylethoxy)carbonyl]-1,2,3,4-tetrahydro-3-isoquinolinyl)carbonyl]-L-phenylalanyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

